

WEST Search History

DATE: Wednesday, July 21, 2004

09/886,197

Hide?	Set Name	Query	Hit Count
		<i>DB=PGPB,USPT,EPAB,JPAB,DWPI; PLUR=YES; OP=ADJ</i>	
<input type="checkbox"/>	L10	L9 and I1	285
<input type="checkbox"/>	L9	L8 and I2	355
<input type="checkbox"/>	L8	wood\$1 with (protect\$8 or prevent\$8 or preserv\$8)	22884
<input type="checkbox"/>	L7	L6 and I1	181
<input type="checkbox"/>	L6	I2 and L3	324
<input type="checkbox"/>	L5	I1 and L4	13
<input type="checkbox"/>	L4	I2 same L3	51
<input type="checkbox"/>	L3	termite\$3	5495
		<i>DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ</i>	
<input type="checkbox"/>	L2	Imidacloprid or ntn33893 or (ntn 33893) or provado or (Chloro adj2 (pyridylmethyl or pyridinylmethyl or ((pyridinyl or pyridyl) adj methyl)) adj3 (nitroimidazolidinylideneamine or nitroimidazolidinimine or (dihydro adj2 nitro adj2 (imidazoleamine or (imidazol\$2 adj2 amine))) or (nitro adj2 imidazolidin\$2 adj2 (ylideneamine or ideneamine or imine))) or (chloro adj2 pyridylmethyl adj2 (nitroimidazo or nitroimidazolidin) adj2 (lidineylideneamine or (lidine adj2 ylideneamine) or ylideneamine)) or (chloro adj2 pyridinyl adj methyl adj3 dihydro adj2 nitro adj2 (imidazoleamine or (imidazole adj2 amine)))	1205
		<i>DB=PGPB,USPT,EPAB,JPAB,DWPI; PLUR=YES; OP=ADJ</i>	
<input type="checkbox"/>	L1	tebuconazole or (chlorophenyl adj3 ((dimethyl adj6 triazol\$2 adj2 (ylmethylpentan\$2 or methylpentan\$2 or ((ylmethyl or methyl) adj pentan\$2))) or (ethyl adj4 (dimethylethyl or tertbutyl or ("t-butyl")) adj5 triazol\$2 adj2 ethanol)))	1164

END OF SEARCH HISTORY

WEST Search History

DATE: Wednesday, July 21, 2004

Hide?	<u>Set</u> <u>Name</u>	<u>Query</u>	<u>Hit</u> <u>Count</u>
	<i>DB=PGPB,USPT,EPAB,JPAB,DWPI; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L10	L9 and I1	285
<input type="checkbox"/>	L9	L8 and I2	355
<input type="checkbox"/>	L8	wood\$1 with (protect\$8 or prevent\$8 or preserv\$8)	22884
<input type="checkbox"/>	L7	L6 and I1	181
<input type="checkbox"/>	L6	I2 and L3	324
<input type="checkbox"/>	L5	I1 and L4	13
<input type="checkbox"/>	L4	I2 same L3	51
<input type="checkbox"/>	L3	termite\$3	5495
	<i>DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L2	Imidacloprid or ntn33893 or (ntn 33893) or provado or (Chloro adj2 (pyridylmethyl or pyridinylmethyl or ((pyridinyl or pyridyl) adj methyl)) adj3 (nitroimidazolidinylideneamine or nitroimidazolidinimine or (dihydro adj2 nitro adj2 (imidazoleamine or (imidazol\$2 adj2 amine))) or (nitro adj2 imidazolidin\$2 adj2 (ylideneamine or ideneamine or imine)))) or (chloro adj2 pyridylmethyl adj2 (nitroimidazo or nitroimidazolidin) adj2 (lidineylideneamine or (lidine adj2 ylideneamine) or ylideneamine)) or (chloro adj2 pyridinyl adj methyl adj3 dihydro adj2 nitro adj2 (imidazoleamine or (imidazole adj2 amine)))	1205
	<i>DB=PGPB,USPT,EPAB,JPAB,DWPI; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L1	tebuconazole or (chlorophenyl adj3 ((dimethyl adj6 triazol\$2 adj2 (ylmethylpentan\$2 or methylpentan\$2 or ((ylmethyl or methyl) adj pentan\$2))) or (ethyl adj4 (dimethylethyl or tertbutyl or ("t-butyl")) adj5 triazol\$2 adj2 ethanol)))	1164

END OF SEARCH HISTORY

Hit List

[Clear](#) [Generate Collection](#) [Print](#) [Fwd Refs](#) [Bkwd Refs](#)
[Generate OACS](#)

Search Results - Record(s) 11 through 13 of 13 returned.

☐ 11. Document ID: US 6114362 A

Using default format because multiple data bases are involved.

L5: Entry 11 of 13

File: USPT

Sep 5, 2000

US-PAT-NO: 6114362

DOCUMENT-IDENTIFIER: US 6114362 A

TITLE: Compositions for the control of plant pests

DATE-ISSUED: September 5, 2000

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Dutzmann; Stefan	Hilden			DE
Erdelen; Christoph	Leichlingen			DE
Andersch; Wolfram	Bergisch Gladbach			DE
Dehne; Heinz-Wilhelm	Bonn			DE
Hartwig; Jurgen	Leichlingen			DE
Stenzel; Klaus	Dusseldorf			DE
Kramer; Wolfgang	Burscheid			DE

US-CL-CURRENT: [514/341](#); [514/383](#)

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequence	References	Claims	KIMC	Drawings
----------------------	-----------------------	--------------------------	-----------------------	------------------------	--------------------------------	----------------------	---------------------------	--------------------------	----------------------------	------------------------	----------------------	--------------------------

☐ 12. Document ID: US 6022881 A

L5: Entry 12 of 13

File: USPT

Feb 8, 2000

US-PAT-NO: 6022881

DOCUMENT-IDENTIFIER: US 6022881 A

TITLE: Insecticidal composition

DATE-ISSUED: February 8, 2000

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Asai; Takehito	Kawasaki			JP
Okumura; Kenya	Tokyo			JP

Shizawa; Toshiyasu

Yokohama

JP

US-CL-CURRENT: 514/341; 514/404, 514/514, 514/515, 514/531

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequence	Alt. Patent	Claims	KWMC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	------	----------

☐ 13. Document ID: US 5935943 A

L5: Entry 13 of 13

File: USPT

Aug 10, 1999

US-PAT-NO: 5935943

DOCUMENT-IDENTIFIER: US 5935943 A

TITLE: Insecticidal composition

DATE-ISSUED: August 10, 1999

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Asai; Takehito	Kawasaki			JP
Okumura; Kenya	Tokyo			JP
Shizawa; Toshiyasu	Yokohama			JP

US-CL-CURRENT: 514/63; 514/383, 514/511, 514/514, 514/515, 514/721

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequence	Alt. Patent	Claims	KWMC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	------	----------

Clear

Generate Collection

Print

Fwd Refs

Bkwd Refs

Generate OACS

Terms

Documents

L1 and L4

13

Display Format: -

Change Format

[Previous Page](#)[Next Page](#)[Go to Doc#](#)

Hit List

[Clear](#)[Generate Collection](#)[Print](#)[Fwd Refs](#)[Bkwd Refs](#)[Generate OACS](#)

Search Results - Record(s) 1 through 10 of 13 returned.

☐ 1. Document ID: US 20040116484 A1

Using default format because multiple data bases are involved.

L5: Entry 1 of 13

File: PGPB

Jun 17, 2004

PGPUB-DOCUMENT-NUMBER: 20040116484

PGPUB-FILING-TYPE: new

DOCUMENT-IDENTIFIER: US 20040116484 A1

TITLE: Agents for combating plant pests

PUBLICATION-DATE: June 17, 2004

INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY	RULE-47
Erdelen, Christoph	Leichlingen		DE	
Andersch, Wolfram	Bergisch Gladbach		DE	
Stenzel, Klaus	Dusseldorf		DE	
Mauler-Machnik, Astrid	Leichlingen		DE	
Kramer, Wolfgang	Burscheid		DE	

US-CL-CURRENT: 514/365

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequences	Attachments	Claims	KIMC	Draw. De
----------------------	-----------------------	--------------------------	-----------------------	------------------------	--------------------------------	----------------------	---------------------------	---------------------------	-----------------------------	------------------------	----------------------	--------------------------

☐ 2. Document ID: US 20030149080 A1

L5: Entry 2 of 13

File: PGPB

Aug 7, 2003

PGPUB-DOCUMENT-NUMBER: 20030149080

PGPUB-FILING-TYPE: new

DOCUMENT-IDENTIFIER: US 20030149080 A1

TITLE: Agents for controlling plant pests

PUBLICATION-DATE: August 7, 2003

INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY	RULE-47
Erdelen, Christoph	Leichlingen		DE	
Andersch, Wolfram	Bergisch Gladbach		DE	
Stenzel, Klaus	Dusseldorf		DE	

Mauler-Machnik, Astrid Leichlingen DE
Kramer, Wolfgang Burscheid DE

US-CL-CURRENT: 514/342

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequences	Attachments	Claims	KIMC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	-----------	-------------	--------	------	---------

☐ 3. Document ID: US 20030083358 A1

L5: Entry 3 of 13

File: PGPB

May 1, 2003

PGPUB-DOCUMENT-NUMBER: 20030083358
PGPUB-FILING-TYPE: new
DOCUMENT-IDENTIFIER: US 20030083358 A1

TITLE: Agents for combating plant pests

PUBLICATION-DATE: May 1, 2003

INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY	RULE-47
Erdelen, Christoph	Leichlingen		DE	
Andersch, Wolfram	Bergisch Gladbach		DE	
Stenzel, Klaus	Dusseldorf		DE	
Mauler-Machnik, Astrid	Leichlingen		DE	
Kramer, Wolfgang	Burscheid		DE	

US-CL-CURRENT: 514/365

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequences	Attachments	Claims	KIMC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	-----------	-------------	--------	------	---------

☐ 4. Document ID: US 20030027813 A1

L5: Entry 4 of 13

File: PGPB

Feb 6, 2003

PGPUB-DOCUMENT-NUMBER: 20030027813
PGPUB-FILING-TYPE: new
DOCUMENT-IDENTIFIER: US 20030027813 A1

TITLE: Pesticide

PUBLICATION-DATE: February 6, 2003

INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY	RULE-47
Dutzmann, Stefan	Hilden		DE	
Erdelen, Christoph	Leichlingen		DE	
Andersch, Wolfram	Bergisch Gladbach		DE	
Dehne, Heinz-Wilhelm	Bonn		DE	

Hartwig, Jorgen	Leichlingen	DE
Stenzel, Klaus	Dusseldorf	DE
Kramer, Wolfgang	Burscheid	DE

US-CL-CURRENT: [514/217.11](#); [514/329](#), [514/426](#), [514/631](#), [514/634](#)

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequences	Attachments	Claims	KWIC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	-----------	-------------	--------	------	---------

☐ 5. Document ID: US 20020006940 A1

L5: Entry 5 of 13

File: PGPB

Jan 17, 2002

PGPUB-DOCUMENT-NUMBER: 20020006940

PGPUB-FILING-TYPE: new

DOCUMENT-IDENTIFIER: US 20020006940 A1

TITLE: Pesticide

PUBLICATION-DATE: January 17, 2002

INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY	RULE-47
Dutzmann, Stefan	Hilden		DE	
Erdelen, Christoph	Leichlingen		DE	
Andersch, Wolfram	Bergisch Gladbach		DE	
Dehne, Heinz-Wilhelm	Bonn		DE	
Hartwig, Jorgen	Leichlingen		DE	
Stenzel, Klaus	Dusseldorf		DE	
Kramer, Wolfgang	Burscheid		DE	

US-CL-CURRENT: [514/341](#); [514/383](#)

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequences	Attachments	Claims	KWIC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	-----------	-------------	--------	------	---------

☐ 6. Document ID: US 6680325 B2

L5: Entry 6 of 13

File: USPT

Jan 20, 2004

US-PAT-NO: 6680325

DOCUMENT-IDENTIFIER: US 6680325 B2

TITLE: Agents for combating plant pests

DATE-ISSUED: January 20, 2004

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Erdelen; Christoph	Leichlingen			DE
Andersch; Wolfram	Bergisch Gladbach			DE

Sterzel; Klaus	Dusseldorf	DE
Mauler-Machnik; Astrid	Leichlingen	DE
Kramer; Wolfgang	Burscheid	DE

US-CL-CURRENT: 514/269; 514/256, 514/365

Full	Title	Citation	Front	Review	Classification	Date	Reference	Attachments	Claims	KMMC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	-------------	--------	------	---------

☐ 7. Document ID: US 6436976 B1

L5: Entry 7 of 13

File: USPT

Aug 20, 2002

US-PAT-NO: 6436976

DOCUMENT-IDENTIFIER: US 6436976 B1

TITLE: Agents for combating plant pests

DATE-ISSUED: August 20, 2002

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Erdelen; Christoph	Leichlingen			DE
Andersch; Wolfram	Bergisch Gladbach			DE
Stenzel; Klaus	Dusseldorf			DE
Mauler-Machnik; Astrid	Leichlingen			DE
Kramer; Wolfgang	Burscheid			DE

US-CL-CURRENT: 514/365; 514/383

Full	Title	Citation	Front	Review	Classification	Date	Reference	Attachments	Claims	KMMC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	-------------	--------	------	---------

☐ 8. Document ID: US 6436968 B1

L5: Entry 8 of 13

File: USPT

Aug 20, 2002

US-PAT-NO: 6436968

DOCUMENT-IDENTIFIER: US 6436968 B1

TITLE: Agents for controlling plant pests

DATE-ISSUED: August 20, 2002

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Erdelen; Christoph	Leichlingen			DE
Andersch; Wolfram	Bergisch Gladbach			DE
Stenzel; Klaus	Dusseldorf			DE
Mauler-Machnik; Astrid	Leichlingen			DE
Kramer; Wolfgang	Burscheid			DE

US-CL-CURRENT: 514/342; 514/383

Full	Title	Citation	Front	Review	Classification	Date	Reference	Attachments	Claims	KIMC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	-------------	--------	------	----------

☐ 9. Document ID: US 6423726 B1

L5: Entry 9 of 13

File: USPT

Jul 23, 2002

US-PAT-NO: 6423726

DOCUMENT-IDENTIFIER: US 6423726 B1

TITLE: Pesticide

DATE-ISSUED: July 23, 2002

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Dutzmann; Stefan	Hilden			DE
Erdelen; Christoph	Leichlingen			DE
Andersch; Wolfram	Bergisch Gladbach			DE
Dehne; Heinz-Wilhelm	Bonn			DE
Hartwig; Jurgen	Leichlingen			DE
Stenzel; Klaus	Dusseldorf			DE
Kramer; Wolfgang	Burscheid			DE

US-CL-CURRENT: 514/341; 514/478, 514/479

Full	Title	Citation	Front	Review	Classification	Date	Reference	Attachments	Claims	KIMC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	-------------	--------	------	----------

☐ 10. Document ID: US 6297263 B1

L5: Entry 10 of 13

File: USPT

Oct 2, 2001

US-PAT-NO: 6297263

DOCUMENT-IDENTIFIER: US 6297263 B1

TITLE: Pesticide

DATE-ISSUED: October 2, 2001

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Dutzmann; Stefan	Hilden			DE
Erdelen; Christoph	Leichlingen			DE
Andersch; Wolfram	Bergisch Gladbach			DE
Dehne; Heinz-Wilhelm	Bonn			DE
Hartwig; Jurgen	Leichlingen			DE
Stenzel; Klaus	Dusseldorf			DE
Kramer; Wolfgang	Burscheid			DE

US-CL-CURRENT: 514/341; 514/422, 514/427

Full	Title	Citation	Front	Review	Classification	Date	Reference	Dependent	Attorney	Claims	KIMC	Draw D
------	-------	----------	-------	--------	----------------	------	-----------	-----------	----------	--------	------	--------

Clear	Generate Collection	Print	Fwd Refs	Bkwd Refs	Generate OACS
-------	---------------------	-------	----------	-----------	---------------

Terms	Documents
L1 and L4	13

Display Format: [Previous Page](#)[Next Page](#)[Go to Doc#](#)

Hit List

Search Results - Record(s) 1020 through 1029 of 1205 returned.

☐ 1020. Document ID: US 5302619 A

Using default format because multiple data bases are involved.

L2: Entry 1020 of 1205

File: USPT

Apr 12, 1994

US-PAT-NO: 5302619

DOCUMENT-IDENTIFIER: US 5302619 A

TITLE: Aromatic compounds, their production processes and their compositions for the control of insect pests

DATE-ISSUED: April 12, 1994

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Shuto; Akira	Takarazuka			JP
Sakamoto; Noriyasu	Nishinomiya			JP
Kisida; Hiroshi	Takarazuka			JP
Fujimoto; Hiroaki	Toyonaka			JP
Umeda; Kimitoshi	Toyonaka			JP
Matsuo; Noritada	Itami			JP

US-CL-CURRENT: 514/718; 514/717, 568/645, 568/647

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	Discussions	Claims	KWIC	Drawings
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	------	----------

☐ 1021. Document ID: US 5288727 A

L2: Entry 1021 of 1205

File: USPT

Feb 22, 1994

US-PAT-NO: 5288727

DOCUMENT-IDENTIFIER: US 5288727 A

TITLE: Hydrazone compounds, processes for their production, intermediates useful for their production and pesticidal compositions containing them

DATE-ISSUED: February 22, 1994

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Toki; Tadaaki	Kusatsu			JP
Koyanagi; Toru	Kusatsu			JP

Yoshida; Kiyomitsu	Kusatsu	JP
Sasaki; Hiroshi	Kusatsu	JP
Morita; Masayuki	Kusatsu	JP
Yoneda; Tetsuo	Kusatsu	JP

US-CL-CURRENT: 514/632; 514/478, 514/614, 514/639, 560/27, 564/151, 564/226,
564/249

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sanitation	Attachment	Claims	KWIC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	------------	------------	--------	------	----------

☐ 1022. Document ID: US 5262441 A

L2: Entry 1022 of 1205

File: USPT

Nov 16, 1993

US-PAT-NO: 5262441

DOCUMENT-IDENTIFIER: US 5262441 A

TITLE: Aromatic compounds, their production processes and their compositions for
the control of insect pests

DATE-ISSUED: November 16, 1993

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Shuto; Akira	Takarazuka			JP
Sakamoto; Noriyasu	Nishinomiya			JP
Kisida; Hiroshi	Takarazuka			JP
Fujimoto; Hiroaki	Osaka			JP
Umeda; Kimitoshi	Osaka			JP
Matsuo; Noritada	Itami			JP

US-CL-CURRENT: 514/718; 514/464, 514/646, 514/657, 514/712, 514/713, 514/716,
514/717, 514/720, 514/721, 514/741, 514/742

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sanitation	Attachment	Claims	KWIC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	------------	------------	--------	------	----------

☐ 1023. Document ID: US 5260312 A

L2: Entry 1023 of 1205

File: USPT

Nov 9, 1993

US-PAT-NO: 5260312

DOCUMENT-IDENTIFIER: US 5260312 A

**** See image for Certificate of Correction ****

TITLE: Stabilized agrochemical compositions

DATE-ISSUED: November 9, 1993

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
------	------	-------	----------	---------

h e b b cg b cc e

Wada; Yuzuru Tokyo JP
Koyama; Shigeharu Oyama JP

US-CL-CURRENT: 514/342; 514/343, 514/365, 514/370, 514/372

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequence	Attachments	Claims	KWIC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	------	---------

☐ 1024. Document ID: US 5258410 A

L2: Entry 1024 of 1205

File: USPT

Nov 2, 1993

US-PAT-NO: 5258410

DOCUMENT-IDENTIFIER: US 5258410 A

TITLE: Aromatic compounds, their production processes and their compositions for the control of insect pests

DATE-ISSUED: November 2, 1993

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Shuto; Akira	Hyogo			JP
Sakamoto; Noriyasu	Hyogo			JP
Kisida; Hiroshi	Hyogo			JP
Fujimoto; Hiroaki	Osaka			JP
Umeda; Kimitoshi	Osaka			JP

US-CL-CURRENT: 514/721; 568/637, 568/640

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequence	Attachments	Claims	KWIC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	------	---------

☐ 1025. Document ID: US 5252610 A

L2: Entry 1025 of 1205

File: USPT

Oct 12, 1993

US-PAT-NO: 5252610

DOCUMENT-IDENTIFIER: US 5252610 A

TITLE: Aromatic compounds and their compositions for the control of insect pests

DATE-ISSUED: October 12, 1993

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Sakamoto; Noriyasu	Nishinomiya			JP
Shuto; Akira	Takarazuka			JP
Kisida; Hiroshi	Takarazuka			JP
Fujimoto; Hiroaki	Toyonaka			JP
Umeda; Kimitoshi	Toyonaka			JP

US-CL-CURRENT: 514/718; 424/405, 568/637

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	Attachments	Claims	KWIC	Drawings
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	------	----------

☐ 1026. Document ID: US 5250574 A

L2: Entry 1026 of 1205

File: USPT

Oct 5, 1993

US-PAT-NO: 5250574

DOCUMENT-IDENTIFIER: US 5250574 A

TITLE: Aromatic compounds and their compositions for the control of insect pests

DATE-ISSUED: October 5, 1993

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Sakamoto; Noriyasu	Hyogo			JP
Shuto; Akira	Hyogo			JP
Kisida; Hiroshi	Hyogo			JP
Fujimoto; Hiroaki	Osaka			JP
Umeda; Kimitoshi	Osaka			JP

US-CL-CURRENT: 514/721; 424/405, 514/456, 549/362, 549/437, 549/445, 568/637

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	Attachments	Claims	KWIC	Drawings
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	------	----------

☐ 1027. Document ID: US 5243087 A

L2: Entry 1027 of 1205

File: USPT

Sep 7, 1993

US-PAT-NO: 5243087

DOCUMENT-IDENTIFIER: US 5243087 A

TITLE: Pyridine derivatives, their production processes and their compositions for the control of insect pests

DATE-ISSUED: September 7, 1993

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Sakamoto; Noriyasu	Hyogo			JP
Kisida; Hiroshi	Hyogo			JP
Fujimoto; Hiroaki	Osaka			JP
Umeda; Kimitoshi	Osaka			JP
Matsuo; Noritada	Hyogo			JP

US-CL-CURRENT: 568/639; 568/640

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequence	Attachments	Claims	KWMC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	------	---------

☐ 1028. Document ID: US 5232949 A

L2: Entry 1028 of 1205

File: USPT

Aug 3, 1993

US-PAT-NO: 5232949

DOCUMENT-IDENTIFIER: US 5232949 A

TITLE: 3-(4-substituted-2-chlorophenoxy)propionaldoxime ethyl ether compounds,
their production processes and their compositions for the control of insect pests

DATE-ISSUED: August 3, 1993

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Shuto; Akira	Takarazuka			JP
Sakamoto; Noriyasu	Nishinomiya			JP
Kisida; Hiroshi	Takarazuka			JP
Matsuo; Noritada	Itami			JP
Fujimoto; Hiroaki	Toyonaka			JP
Umeda; Kimitoshi	Toyonaka			JP

US-CL-CURRENT: 514/640; 564/256

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequence	Attachments	Claims	KWMC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	------	---------

☐ 1029. Document ID: US 5232701 A

L2: Entry 1029 of 1205

File: USPT

Aug 3, 1993

US-PAT-NO: 5232701

DOCUMENT-IDENTIFIER: US 5232701 A

**** See image for Certificate of Correction ****

TITLE: Boron carbonate and solid acid pesticidal composition

DATE-ISSUED: August 3, 1993

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Ogawa; Masao	Toyonaka			JP
Ohtsubo; Toshiro	Sanda			JP
Tsuda; Shigenori	Kyoto			JP

US-CL-CURRENT: 424/408; 424/409, 424/466, 424/489, 504/358

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequence	Attachments	Claims	KWMC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	------	---------

Clear

Generate Collection

Print

Fwd Refs

Bkwd Refs

Generate OACS

Terms	Documents
Imidacloprid or ntn33893 or (ntn 33893) or provado or (Chloro adj2 (pyridylmethyl or pyridinylmethyl or ((pyridinyl or pyridyl) adj methyl)) adj3 (nitroimidazolidinylideneamine or nitroimidazolidinimine or (dihydro adj2 nitro adj2 (imidazoleamine or (imidazol\$2 adj2 amine))) or (nitro adj2 imidazolidin\$2 adj2 (ylideneamine or ideneamine or imine)))) or (chloro adj2 pyridylmethyl adj2 (nitroimidazo or nitroimidazolidin) adj2 (lidineylideneamine or (lidine adj2 ylideneamine) or ylideneamine)) or (chloro adj2 pyridinyl adj methyl adj3 dihydro adj2 nitro adj2 (imidazoleamine or (imidazole adj2 amine)))	1205

Display Format:

Change Format

[Previous Page](#)[Next Page](#)[Go to Doc#](#)

Hit List

[Clear](#) [Generate Collection](#) [Print](#) [Fwd Refs](#) [Bkwd Refs](#)
[Generate OACS](#)

Search Results - Record(s) 1050 through 1059 of 1205 returned.

☐ 1050. Document ID: JP 10158106 A

Using default format because multiple data bases are involved.

L2: Entry 1050 of 1205

File: JPAB

Jun 16, 1998

PUB-NO: JP410158106A

DOCUMENT-IDENTIFIER: JP 10158106 A

TITLE: INSECTICIDAL/MITICIDAL COMPOSITION

PUBN-DATE: June 16, 1998

INVENTOR-INFORMATION:

NAME

COUNTRY

OBATA, TOKIO

TSUTSUMIUCHI, KIYOSHI

INT-CL (IPC): A01 N 43/54

Full	Title	Citation	Front	Review	Classification	Date	Reference	Send to	Abstract	Claims	KWIC	Draw D
------	-------	----------	-------	--------	----------------	------	-----------	---------	----------	--------	------	--------

☐ 1051. Document ID: JP 08092091 A

L2: Entry 1051 of 1205

File: JPAB

Apr 9, 1996

PUB-NO: JP408092091A

DOCUMENT-IDENTIFIER: JP 08092091 A

TITLE: NON-SYSTEMIC CONTROL OF PARASITE

PUBN-DATE: April 9, 1996

INVENTOR-INFORMATION:

NAME

COUNTRY

DORN, HUBERT DR

HOPKINS, TERENCE DR

INT-CL (IPC): A61 K 31/44; A61 K 31/425; A61 K 31/495; A61 K 31/53; A61 K 31/54;
A61 K 45/00; C07 D 213/61; C07 D 233/64 ; C07 D 277/10; C07 D 279/06; C07 D 401/04;
C07 D 401/06; C07 D 401/06; C07 D 401/06; C07 D 401/06; C07 D 417/06; C07 D 417/06;
C07 D 417/06

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequence	Attachments	Claims	KWMC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	------	----------

☐ 1052. Document ID: JP 08037818 A

L2: Entry 1052 of 1205

File: JPAB

Feb 13, 1996

PUB-NO: JP408037818A

DOCUMENT-IDENTIFIER: JP 08037818 A

TITLE: AGROCHEMICAL-CONTAINING COATED SEED

PUBN-DATE: February 13, 1996

INVENTOR-INFORMATION:

NAME

COUNTRY

HASEGAWA, AKIRA

OKADA, SEIICHI

OKABE, TAKAYUKI

NIIDE, YOJIRO

KOBAYASHI, TOSHIAKI

INT-CL (IPC): A01 C 1/06

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequence	Attachments	Claims	KWMC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	------	----------

☐ 1053. Document ID: JP 07277906 A

L2: Entry 1053 of 1205

File: JPAB

Oct 24, 1995

PUB-NO: JP407277906A

DOCUMENT-IDENTIFIER: JP 07277906 A

TITLE: SYNERGISTIC COMBINED AGENT OF AMMONIUM SALT

PUBN-DATE: October 24, 1995

INVENTOR-INFORMATION:

NAME

COUNTRY

SAGENMUELLER, ALFONS

SCHUBERT, HANS-HERBERT

UZAWA, SHIGERU

SAITO, KENICHI

INT-CL (IPC): A01 N 33/12; A01 N 55/00; A01 N 57/10

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequence	Attachments	Claims	KWMC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	------	----------

☐ 1054. Document ID: JP 05246806 A

L2: Entry 1054 of 1205

File: JPAB

Sep 24, 1993

PUB-NO: JP405246806A
DOCUMENT-IDENTIFIER: JP 05246806 A
TITLE: AGROCHEMICAL MIXTURE

PUBN-DATE: September 24, 1993

INVENTOR-INFORMATION:

NAME

COUNTRY

CARUHELL, PASCAL

MUGNIER, JACQUES

INT-CL (IPC): A01N 43/653

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	Attachments	Claims	Keywords	Drawings
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	----------	----------

☐ 1055. Document ID: GB 2381750 A

L2: Entry 1055 of 1205

File: EPAB

May 14, 2003

PUB-NO: GB002381750A
DOCUMENT-IDENTIFIER: GB 2381750 A
TITLE: Treatment for enhancing joint lubrication

PUBN-DATE: May 14, 2003

INVENTOR-INFORMATION:

NAME

COUNTRY

YERXA, BENJAMIN R

US

COWLEN, MATTHEW S

US

INT-CL (IPC): A61 K 31/465; A61 K 31/4168; A61 K 31/4178; A61 K 31/422; A61 K 31/4427; A61 K 31/4439; A61 K 31/444; A61 K 31/445; A61 K 31/501; A61 K 31/504; A61 P 19/02

EUR-CL (EPC): A61K031/00; A61K031/00, A61K031/4168 , A61K031/4178 , A61K031/422 , A61K031/4427 , A61K031/4439 , A61K031/444 , A61K031/445 , A61K031/465 , A61K031/501 , A61K031/504

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	Attachments	Claims	Keywords	Drawings
------	-------	----------	-------	--------	----------------	------	-----------	----------	-------------	--------	----------	----------

☐ 1056. Document ID: WO 3015515 A1

L2: Entry 1056 of 1205

File: EPAB

Feb 27, 2003

PUB-NO: WO003015515A1
DOCUMENT-IDENTIFIER: WO 3015515 A1
TITLE: FUNGICIDAL ACTIVE SUBSTANCE COMBINATIONS CONTAINING TRIFLOXYSTROBIN

PUBN-DATE: February 27, 2003

INVENTOR-INFORMATION:

NAME	COUNTRY
WACHENDORFF-NEUMANN, ULRIKE	DE
MAULER-MACHNIK, ASTRID	DE
ERDELEN, CHRISTOPH	DE
OHTAKE, HIROHISA	JP

INT-CL (IPC): A01 N 37/36; A01 N 51/00; A01 N 47/40; A01 N 43/40
EUR-CL (EPC): A01N037/50

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	References	Claims	KIMC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	----------	------------	--------	------	----------

☐ 1057. Document ID: EP 1252820 A1

L2: Entry 1057 of 1205

File: EPAB

Oct 30, 2002

PUB-NO: EP001252820A1

DOCUMENT-IDENTIFIER: EP 1252820 A1

TITLE: Pediculicidal and ovacidal treatment compositions and methods for killing head lice and their eggs

PUBN-DATE: October 30, 2002

INVENTOR-INFORMATION:

NAME	COUNTRY
JANSSEN, HERWIG	US
HO, KIE	US
NYSTRAND, GLENN	US
WILLIAMS, DEXTER	US
LAMB, SCOTT C	US

INT-CL (IPC): A01 N 43/22; A01 N 51/00

EUR-CL (EPC): A01N025/02; A01N043/22, A01N051/00

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	References	Claims	KIMC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	----------	------------	--------	------	----------

☐ 1058. Document ID: WO 137668 A2

L2: Entry 1058 of 1205

File: EPAB

May 31, 2001

PUB-NO: WO000137668A2

DOCUMENT-IDENTIFIER: WO 137668 A2

TITLE: FUNGICIDALLY ACTIVE SUBSTANCE COMBINATIONS

PUBN-DATE: May 31, 2001

INVENTOR-INFORMATION:

NAME	COUNTRY
KUCK, KARL-HEINZ	DE
MAULER-MACHNIK, ASTRID	DE

WACHENDORFF-NEUMANN, ULRIKE
KRAEMER, WOLFGANG

DE
DE

INT-CL (IPC): A01 N 51/00; A01 N 47/40
EUR-CL (EPC): A01N051/00

Full	Title	Citation	Front	Review	Classification	Date	Reference	Source	Attachments	Claims	KWIC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	--------	-------------	--------	------	---------

☐ 1059. Document ID: FR 2784011 A1

L2: Entry 1059 of 1205

File: EPAB

Apr 7, 2000

PUB-NO: FR002784011A1

DOCUMENT-IDENTIFIER: FR 2784011 A1

TITLE: Insecticidal compositions contain pyrethroid and chloronicotinyl
insecticide, for the treatment of cotton plants

PUBN-DATE: April 7, 2000

INVENTOR-INFORMATION:

NAME

COUNTRY

COULIBALY, ADAMA

INT-CL (IPC): A01 N 53/12

EUR-CL (EPC): A01N053/00

Full	Title	Citation	Front	Review	Classification	Date	Reference	Source	Attachments	Claims	KWIC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	--------	-------------	--------	------	---------

Clear	Generate Collection	Print	Fwd Refs	Bkwd Refs	Generate OACS
-------	---------------------	-------	----------	-----------	---------------

Terms	Documents
Imidacloprid or ntn33893 or (ntn 33893) or provado or (Chloro adj2 (pyridylmethyl or pyridinylmethyl or ((pyridinyl or pyridyl) adj methyl)) adj3 (nitroimidazolidinylideneamine or nitroimidazolidinimine or (dihydro adj2 nitro adj2 (imidazoleamine or (imidazol\$2 adj2 amine))) or (nitro adj2 imidazolidin\$2 adj2 (ylideneamine or ideneamine or imine))) or (chloro adj2 pyridylmethyl adj2 (nitroimidazo or nitroimidazolidin) adj2 (lidineylideneamine or (lidine adj2 ylideneamine) or ylideneamine)) or (chloro adj2 pyridinyl adj methyl adj3 dihydro adj2 nitro adj2 (imidazoleamine or (imidazole adj2 amine)))	1205

Display Format:

[Previous Page](#)

[Next Page](#)

[Go to Doc#](#)

WEST Search History

Hide Items

Restore

Clear

Cancel

09/886,197

DATE: Wednesday, July 21, 2004

Hide?	<u>Set</u> <u>Name</u>	<u>Query</u>	<u>Hit</u> <u>Count</u>
	<i>DB=PGPB,USPT,EPAB,JPAB,DWPI; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L20	l11 same L18	0
<input type="checkbox"/>	L19	l11 and L18	146
<input type="checkbox"/>	L18	insect\$9 same wood\$1	6165
<input type="checkbox"/>	L17	l11 and L16	115
<input type="checkbox"/>	L16	(424 or 514).clas.	207995
<input type="checkbox"/>	L15	l11 and L14	0
<input type="checkbox"/>	L14	l3 or l4	341
<input type="checkbox"/>	L13	l11 and L12	182
<input type="checkbox"/>	L12	insecticid\$8 and fungicid\$8	20504
<input type="checkbox"/>	L11	(l1 same l2) not l5	182
<input type="checkbox"/>	L10	l5 not l7	1
<input type="checkbox"/>	L9	l7 and L8	0
<input type="checkbox"/>	L8	wood	387229
<input type="checkbox"/>	L7	l5 and L6	11
<input type="checkbox"/>	L6	l2 and l3	14
<input type="checkbox"/>	L5	l1 and L4	12
<input type="checkbox"/>	L4	L2.ab.	170
<input type="checkbox"/>	L3	l1.ab.	182
<input type="checkbox"/>	L2	tebuconazole or (chlorophenyl adj3 ((dimethyl adj6 triazol\$2 adj2 (ylmethylpentan\$2 or methylpentan\$2 or ((ylmethyl or methyl) adj pentan\$2))) or (ethyl adj4 (dimethylethyl or tertbutyl or ("t-butyl")) adj5 triazol\$2 adj2 ethanol)))	1164
	<i>DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L1	Imidacloprid or ntn33893 or (ntn 33893) or provado or (Chloro adj2 (pyridylmethyl or pyridinylmethyl or ((pyridinyl or pyridyl) adj methyl)) adj3 (nitroimidazolidinylideneamine or nitroimidazolidinimine or (dihydro adj2 nitro adj2 (imidazoleamine or (imidazol\$2 adj2 amine))) or (nitro adj2 imidazolidin\$2 adj2 (ylideneamine or ideneamine or imine))) or (chloro adj2 pyridylmethyl adj2 (nitroimidazo or nitroimidazolidin) adj2 (lidineylideneamine or (lidine adj2 ylideneamine) or ylideneamine)) or (chloro adj2 pyridinyl adj methyl adj3 dihydro adj2 nitro adj2 (imidazoleamine or (imidazole adj2 amine)))	1205

END OF SEARCH HISTORY

WEST Search History

Hide Items

Restore

Clear

Cancel

09/886,197

DATE: Wednesday, July 21, 2004

Hide?	Set Name	Query	Hit Count
	<i>DB=PGPB,USPT,EPAB,JPAB,DWPI; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L20	111 same L18	0
<input type="checkbox"/>	L19	111 and L18	146
<input type="checkbox"/>	L18	insect\$9 same wood\$1	6165
<input type="checkbox"/>	L17	111 and L16	115
<input type="checkbox"/>	L16	(424 or 514).clas.	207995
<input type="checkbox"/>	L15	111 and L14	0
<input type="checkbox"/>	L14	l3 or l4	341
<input type="checkbox"/>	L13	111 and L12	182
<input type="checkbox"/>	L12	insecticid\$8 and fungicid\$8	20504
<input type="checkbox"/>	L11	(l1 same l2) not l5	182
<input type="checkbox"/>	L10	l5 not l7	1
<input type="checkbox"/>	L9	l7 and L8	0
<input type="checkbox"/>	L8	wood	387229
<input type="checkbox"/>	L7	l5 and L6	11
<input type="checkbox"/>	L6	l2 and l3	14
<input type="checkbox"/>	L5	l1 and L4	12
<input type="checkbox"/>	L4	L2.ab.	170
<input type="checkbox"/>	L3	l1.ab.	182
<input type="checkbox"/>	L2	tebuconazole or (chlorophenyl adj3 ((dimethyl adj6 triazol\$2 adj2 (ylmethylpentan\$2 or methylpentan\$2 or ((ylmethyl or methyl) adj pentan\$2))) or (ethyl adj4 (dimethylethyl or tertbutyl or ("t-butyl")) adj5 triazol\$2 adj2 ethanol)))	1164
	<i>DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ</i>		
<input type="checkbox"/>	L1	Imidacloprid or ntn33893 or (ntn 33893) or provado or (Chloro adj2 (pyridylmethyl or pyridinylmethyl or ((pyridinyl or pyridyl) adj methyl)) adj3 (nitroimidazolidinylideneamine or nitroimidazolidinimine or (dihydro adj2 nitro adj2 (imidazoleamine or (imidazol\$2 adj2 amine))) or (nitro adj2 imidazolidin\$2 adj2 (ylideneamine or ideneamine or imine))) or (chloro adj2 pyridylmethyl adj2 (nitroimidazo or nitroimidazolidin) adj2 (lidineylideneamine or (lidine adj2 ylideneamine) or ylideneamine)) or (chloro adj2 pyridinyl adj methyl adj3 dihydro adj2 nitro adj2 (imidazoleamine or (imidazole adj2 amine)))	1205

END OF SEARCH HISTORY

WEST Search History

DATE: Wednesday, July 21, 2004

Hide?	Set Name	Query	Hit Count
		<i>DB=PGPB,USPT,EPAB,JPAB,DWPI; PLUR=YES; OP=ADJ</i>	
<input type="checkbox"/>	L9	17 and L8	0
<input type="checkbox"/>	L8	wood	387229
<input type="checkbox"/>	L7	15 and L6	11
<input type="checkbox"/>	L6	12 and 13	14
<input type="checkbox"/>	L5	11 and L4	12
<input type="checkbox"/>	L4	L2.ab.	170
<input type="checkbox"/>	L3	11.ab.	182
<input type="checkbox"/>	L2	tebuconazole or (chlorophenyl adj3 ((dimethyl adj6 triazol\$2 adj2 (ylmethylpentan\$2 or methylpentan\$2 or ((ylmethyl or methyl) adj pentan\$2))) or (ethyl adj4 (dimethylethyl or tertbutyl or ("t-butyl")) adj5 triazol\$2 adj2 ethanol)))	1164
		<i>DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ</i>	
<input type="checkbox"/>	L1	Imidacloprid or ntn33893 or (ntn 33893) or provado or (Chloro adj2 (pyridylmethyl or pyridinylmethyl or ((pyridinyl or pyridyl) adj methyl)) adj3 (nitroimidazolidinylideneamine or nitroimidazolidinimine or (dihydro adj2 nitro adj2 (imidazoleamine or (imidazol\$2 adj2 amine))) or (nitro adj2 imidazolidin\$2 adj2 (ylideneamine or ideneamine or imine))) or (chloro adj2 pyridylmethyl adj2 (nitroimidazo or nitroimidazolidin) adj2 (lidineylideneamine or (lidine adj2 ylideneamine) or ylideneamine)) or (chloro adj2 pyridinyl adj methyl adj3 dihydro adj2 nitro adj2 (imidazoleamine or (imidazole adj2 amine)))	1205

END OF SEARCH HISTORY

Hit List

Clear

Generate Collection

Print

Fwd Refs

Bkwd Refs

Generate OACS

Search Results - Record(s) 1 through 10 of 11 returned.☐ 1. Document ID: US 20020006940 A1**Using default format because multiple data bases are involved.**

L7: Entry 1 of 11

File: PGPB

Jan 17, 2002

PGPUB-DOCUMENT-NUMBER: 20020006940

PGPUB-FILING-TYPE: new

DOCUMENT-IDENTIFIER: US 20020006940 A1

TITLE: Pesticide

PUBLICATION-DATE: January 17, 2002

INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY	RULE-47
Dutzmann, Stefan	Hilden		DE	
Erdelen, Christoph	Leichlingen		DE	
Andersch, Wolfram	Bergisch Gladbach		DE	
Dehne, Heinz-Wilhelm	Bonn		DE	
Hartwig, Jurgen	Leichlingen		DE	
Stenzel, Klaus	Dusseldorf		DE	
Kramer, Wolfgang	Burscheid		DE	

US-CL-CURRENT: [514/341](#); [514/383](#)

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequences	Attachments	Claims	KWIC	Drawings
------	-------	----------	-------	--------	----------------	------	-----------	-----------	-------------	--------	------	----------

☒ 2. Document ID: US 6423726 B1

L7: Entry 2 of 11

File: USPT

Jul 23, 2002

US-PAT-NO: 6423726

DOCUMENT-IDENTIFIER: US 6423726 B1

TITLE: Pesticide

DATE-ISSUED: July 23, 2002

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Dutzmann; Stefan	Hilden			DE
Erdelen; Christoph	Leichlingen			DE

h e b b g e e e f e f g e e f b e

Andersch; Wolfram	Bergisch Gladbach	DE
Dehne; Heinz-Wilhelm	Bonn	DE
Hartwig; Jurgen	Leichlingen	DE
Stenzel; Klaus	Dusseldorf	DE
Kramer; Wolfgang	Burscheid	DE

US-CL-CURRENT: 514/341; 514/478, 514/479

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	Claims	KWIC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	----------	--------	------	----------

☐ 3. Document ID: US 20020110574 A1, US 6488949 B2

L7: Entry 3 of 11

File: DWPI

Aug 15, 2002

DERWENT-ACC-NO: 2003-110440

DERWENT-WEEK: 200310

COPYRIGHT 2004 DERWENT INFORMATION LTD

TITLE: Composition for inhibiting phase separation in low viscous aqueous pesticide suspension, comprises preset amount of pesticide, wetting agent, thickener, antimicrobial, anti-freeze agent, hydrophobic silica and water

INVENTOR: HUDSON, D C; SHAFER, J G

PRIORITY-DATA: 2001US-0759797 (January 12, 2001), 1998US-0086075 (May 28, 1998), 1999US-0228904 (January 11, 1999), 2000US-0506655 (February 17, 2000), 2002US-0071539 (February 8, 2002)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
<u>US 20020110574 A1</u>	August 15, 2002		007	A01N025/00
<u>US 6488949 B2</u>	December 3, 2002		000	A01N025/00

INT-CL (IPC): A01 N 25/00; A01 N 43/64; A01 N 43/72; A01 N 43/78; A61 K 31/425; A61 K 31/44; A61 K 31/535

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	Claims	KWIC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	----------	--------	------	----------

☐ 4. Document ID: AU 200214045 A, WO 200237964 A1

L7: Entry 4 of 11

File: DWPI

May 21, 2002

DERWENT-ACC-NO: 2002-471526

DERWENT-WEEK: 200260

COPYRIGHT 2004 DERWENT INFORMATION LTD

TITLE: Synergistic pesticidal composition useful for controlling pests comprises combination of N-cyanomethyl-4-(trifluoromethyl)-nicotinamide and at least one other compound e.g. azamethiphos or chlorfenvinphos

INVENTOR: ANGST, M; MAIENFISCH, P ; RINDLISBACHER, A

PRIORITY-DATA: 2000CH-0002189 (November 10, 2000)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
AU 200214045 A	May 21, 2002		000	A01N043/40
WO 200237964 A1	May 16, 2002	E	030	A01N043/40

INT-CL (IPC): A01 N 43/40; A01 N 61/00; A01 N 61:00; A01 N 43/40

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	Claims	KWIC	Draw D
------	-------	----------	-------	--------	----------------	------	-----------	----------	--------	------	--------

□ 5. Document ID: EP 1289366 B1, DE 10103832 A1, WO 200184931 A1, AU 200172383 A, EP 1289366 A1, CZ 200203724 A3, BR 200110699 A, KR 2002093063 A, CN 1429074 A, HU 200302025 A2, JP 2003532654 W, US 20030229087 A1, ZA 200208114 A

L7: Entry 5 of 11

File: DWPI

Jun 30, 2004

DERWENT-ACC-NO: 2002-042659

DERWENT-WEEK: 200444

COPYRIGHT 2004 DERWENT INFORMATION LTD

TITLE: Synergistic combination of fungicides for use in plant protection, comprises 2-(pyrimidinyl-oxy-phenyl)-2-(methoxyimino)-N-methyl-acetamide derivative and e.g. spiroxamine, quinoxifen or tebuconazole

INVENTOR: GAYER, H; MAULER-MACHNIK, A ; WACHENDORFF-NEUMANN, U ; WACHENDORFF-NEUMANN, U

PRIORITY-DATA: 2000DE-1022951 (May 11, 2000), 2002US-0275500 (November 6, 2002)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
EP 1289366 B1	June 30, 2004	G	000	A01N043/54
DE 10103832 A1	November 15, 2001		106	A01N043/54
WO 200184931 A1	November 15, 2001	G	000	A01N043/54
AU 200172383 A	November 20, 2001		000	A01N043/54
EP 1289366 A1	March 12, 2003	G	000	A01N043/54
CZ 200203724 A3	March 12, 2003		000	A01N043/54
BR 200110699 A	March 18, 2003		000	A01N043/54
KR 2002093063 A	December 12, 2002		000	A01N043/54
CN 1429074 A	July 9, 2003		000	A01N043/54
HU 200302025 A2	September 29, 2003		000	A01N043/54
JP 2003532654 W	November 5, 2003		167	A01N043/54
US 20030229087 A1	December 11, 2003		000	A01N043/58
ZA 200208114 A	December 31, 2003		162	A01N000/00

INT-CL (IPC): A01 N 0/00; A01 N 31/10; A01 N 37/18; A01 N 37/20; A01 N 37/22; A01 N 37/24; A01 N 37/34; A01 N 37/50; A01 N 37/52; A01 N 39/02; A01 N 43/08; A01 N 43/30; A01 N 43/34; A01 N 43/36; A01 N 43/38; A01 N 43/40; A01 N 43/42; A01 N 43/50; A01 N 43/52; A01 N 43/54; A01 N 43/58; A01 N 43/60; A01 N 43/653; A01 N 43/707; A01 N 43/78; A01 N 43/84; A01 N 43/86; A01 N 43/88; A01 N 43/90; A01 N 43/54; A01 N 47/04; A01 N 47/10; A01 N 47/12; A01 N 47/14; A01 N 47/18; A01 N 47/34; A01 N 47/36; A01 N 47/38; A01 N 47/44; A01 N 51/00; A01 N 53/08; A01 N 53/12; A01 N 55/00; A01 N 57/10; A01 N 57/14; A01 N 61/00; A01 N 61:00; A01 N

43/54; A01 N 61:00

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequence	Drawings	Claims	KMC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	----------	----------	--------	-----	----------

☐ 6. Document ID: MX 2002010331 A1, DE 10019758 A1, WO 200180641 A2, AU 200160189 A, EP 1276375 A2, BR 200110116 A, KR 2002087985 A, US 20030158151 A1, CN 1436044 A, JP 2003531154 W, ZA 200207474 A, HU 200302686 A2

L7: Entry 6 of 11

File: DWPI

May 1, 2003

DERWENT-ACC-NO: 2002-035268

DERWENT-WEEK: 200415

COPYRIGHT 2004 DERWENT INFORMATION LTD

TITLE: Fungicidal combinations containing known methoxyimino-acetic acid amide derivatives useful for the control of phytopathogenic fungi

INVENTOR: ASSMANN, L; GAYER, H ; HEINEMANN, U ; KRAEMER, W ; KRUEGER, B ; NEUMANN, U W ; SEITZ, T ; WACHENDORFF-NEUMANN, U ; KRAMER, W ; KRUGER, B

PRIORITY-DATA: 2000DE-1019758 (April 20, 2000)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
<u>MX 2002010331 A1</u>	May 1, 2003		000	A01N037/50
<u>DE 10019758 A1</u>	October 25, 2001		040	A01N035/10
<u>WO 200180641 A2</u>	November 1, 2001	G	000	A01N043/00
<u>AU 200160189 A</u>	November 7, 2001		000	A01N043/00
<u>EP 1276375 A2</u>	January 22, 2003	G	000	A01N037/50
<u>BR 200110116 A</u>	February 11, 2003		000	A01N043/00
<u>KR 2002087985 A</u>	November 23, 2002		000	A01N043/653
<u>US 20030158151 A1</u>	August 21, 2003		000	A01N055/00
<u>CN 1436044 A</u>	August 13, 2003		000	A01N037/50
<u>JP 2003531154 W</u>	October 21, 2003		095	A01N037/50
<u>ZA 200207474 A</u>	November 26, 2003		096	A01N000/00
<u>HU 200302686 A2</u>	November 28, 2003		000	A01N043/00

INT-CL (IPC): A01 N 0/00; A01 N 35/10; A01 N 37/50; A01 N 39/02; A01 N 43/00; A01 N 43/40; A01 N 43/42; A01 N 43/54; A01 N 43/653; A01 N 43/78; A01 N 43/828; A01 N 43/88; A01 N 47/10; A01 N 55/00; A01 N 57/10; A01 N 59/16; A01 N 59/20; A01 N 61/00; A01 N 37/50; A01 N 59/20; A01 N 61:00

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequence	Drawings	Claims	KMC	Draw. De
------	-------	----------	-------	--------	----------------	------	-----------	----------	----------	--------	-----	----------

☐ 7. Document ID: US 20010008873 A1, US 6379687 B2

L7: Entry 7 of 11

File: DWPI

Jul 19, 2001

DERWENT-ACC-NO: 2001-557096

DERWENT-WEEK: 200310

COPYRIGHT 2004 DERWENT INFORMATION LTD

TITLE: Composition for inhibiting phase separation and non-uniform distribution of

h e b b g e e e f e f g e e f b e

active ingredient in low viscosity water-based pesticide suspension, includes hydrophobic silica

INVENTOR: HUDSON, D C; SHAFER, J G

PRIORITY-DATA: 2001US-0759797 (January 12, 2001), 1998US-0086075 (May 28, 1998), 1999US-0228904 (January 11, 1999), 2000US-0506655 (February 17, 2000)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
US 20010008873 A1	July 19, 2001		007	A01N043/72
US 6379687 B2	April 30, 2002		000	A01N025/00

INT-CL (IPC): A01 N 25/00; A01 N 43/72; A01 N 43/78; A61 K 31/425; A61 K 31/44; A61 K 31/535

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	Claims	Summary	Drawings
------	-------	----------	-------	--------	----------------	------	-----------	----------	--------	---------	----------

8. Document ID: TW 521994 A, DE 19939841 A1, WO 200030440 A2, AU 200010460 A, BR 9915518 A, EP 1130963 A2, CZ 200101749 A3, KR 2001085954 A, CN 1326316 A, HU 200104483 A2, MX 2001005029 A1, ZA 200103143 A, AU 752441 B, JP 2002530297 W, US 6559136 B1, US 20030161896 A1

L7: Entry 8 of 11

File: DWPI

Mar 1, 2003

DERWENT-ACC-NO: 2000-367125

DERWENT-WEEK: 200365

COPYRIGHT 2004 DERWENT INFORMATION LTD

TITLE: Synergistic fungicide combination for use in plant protection contains 4,6-diphenoxy-5-halo-pyrimidine derivative and e.g. tebuconazole, fenpropimorph, azoxystrobin, carbendazim or folpet

INVENTOR: GAYER, H; MAUIER-MACHNIK, A ; WACHENDORFF-NEUMANN, U ; MAULER-MACHNIK, A ; WACHENDORF-NEUMANN, U

PRIORITY-DATA: 1998DE-1053559 (November 20, 1998)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
TW 521994 A	March 1, 2003		000	A01N043/54
DE 19939841 A1	May 25, 2000		017	A01N043/54
WO 200030440 A2	June 2, 2000	G	000	A01N000/00
AU 200010460 A	June 13, 2000		000	
BR 9915518 A	July 17, 2001		000	A01N043/54
EP 1130963 A2	September 12, 2001	G	000	A01N043/54
CZ 200101749 A3	December 12, 2001		000	A01N043/54
KR 2001085954 A	September 7, 2001		000	A01N043/54
CN 1326316 A	December 12, 2001		000	A01N043/54
HU 200104483 A2	March 28, 2002		000	A01N043/54
MX 2001005029 A1	August 1, 2001		000	A01N000/00000
ZA 200103143 A	June 26, 2002		061	A01N000/00

AU 752441 B	September 19, 2002	000	A01N043/54
JP 2002530297 W	September 17, 2002	041	A01N043/54
US 6559136 B1	May 6, 2003	000	A01N043/653
US 20030161896 A1	August 28, 2003	000	A01N065/00

B1 , US 20030161896 A1 INT-CL (IPC): A01 N 0/00; A01 N 0/00000; A01 N 37/18; A01 N 37/28; A01 N 37/34; A01 N 43/54; A01 N 43/64; A01 N 43/653; A01 N 43/80; A01 N 55/10; A01 N 59/02; A01 N 65/00

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	Claims	Summary	Drawings
------	-------	----------	-------	--------	----------------	------	-----------	----------	--------	---------	----------

□ 9. Document ID: TW 505504 A, DE 19716257 A1, WO 9847367 A1, ZA 9803236 A, AU 9875220 A, EP 975219 A1, CZ 9903696 A3, CN 1252690 A, SK 9901435 A3, BR 9809100 A, NZ 500367 A, HU 200001682 A2, AU 727186 B, MX 9909480 A1, KR 2001006063 A, US 6306850 B1, JP 2001520665 W, EP 975219 B1, DE 59803337 G, ES 2172143 T3, US 20020173529 A1

L7: Entry 9 of 11

File: DWPI

Oct 11, 2002

DERWENT-ACC-NO: 1998-558428

DERWENT-WEEK: 200341

COPYRIGHT 2004 DERWENT INFORMATION LTD

TITLE: Synergistic fungicidal composition for use in plant protection - contains substituted hydroxypropyl-triazole:thione derivative and conventional fungicidal compounds.

INVENTOR: DUTZMANN, S; JAUTELAT, M ; STENZEL, K

PRIORITY-DATA: 1997DE-1016257 (April 18, 1997)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
<u>TW 505504 A</u>	October 11, 2002		000	A01N043/653
<u>DE 19716257 A1</u>	October 22, 1998		044	A01N043/653
<u>WO 9847367 A1</u>	October 29, 1998	G	000	A01N043/653
<u>ZA 9803236 A</u>	December 30, 1998		070	A01N000/00
<u>AU 9875220 A</u>	November 13, 1998		000	A01N043/653
<u>EP 975219 A1</u>	February 2, 2000	G	000	A01N043/653
<u>CZ 9903696 A3</u>	February 16, 2000		000	A01N043/653
<u>CN 1252690 A</u>	May 10, 2000		000	A01N043/653
<u>SK 9901435 A3</u>	June 12, 2000		000	A01N043/653
<u>BR 9809100 A</u>	August 1, 2000		000	A01N043/653
<u>NZ 500367 A</u>	September 29, 2000		000	A01N043/653
<u>HU 200001682 A2</u>	September 28, 2000		000	A01N043/653
<u>AU 727186 B</u>	December 7, 2000		000	A01N043/653
<u>MX 9909480 A1</u>	February 1, 2000		000	A01N043/653
<u>KR 2001006063 A</u>	January 15, 2001		000	A01N043/653
<u>US 6306850 B1</u>	October 23, 2001		000	A61K031/535
<u>JP 2001520665 W</u>	October 30, 2001		063	A01N043/653
<u>EP 975219 B1</u>	March 13, 2002	G	000	A01N043/653

DE 59803337 G	April 18, 2002	000	A01N043/653
ES 2172143 T3	September 16, 2002	000	A01N043/653
US 20020173529 A1	November 21, 2002	000	A01N043/64

B1 , JP 2001520665 W INT-CL (IPC): A01 N 0/00; A01 N 37/18; A01 N 37/24; A01 N 37/26; A01 N 37/34; A01 N 37/50; A01 N 41/02; A01 N 43/30; A01 N 43/36; A01 N 43/54; A01 N 43/64; A01 N 43/653; A01 N 43/828; A01 N 43/84; A01 N 43/88; A01 N 47/04; A01 N 47/12; A01 N 47/30; A01 N 47/32; A01 N 47/44; A01 N 51/00; A01 N 53/12; A01 N 55/02; A01 N 57/12; A61 K 31/535

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	Claims	KWIC	Draw Ds
------	-------	----------	-------	--------	----------------	------	-----------	----------	--------	------	---------

☐ 10. Document ID: DE 19716256 A1, US 20020072535 A1, WO 9847370 A1, ZA 9803235 A, AU 9875221 A, EP 975221 A1, CZ 9903697 A3, CN 1252692 A, BR 9809763 A, TW 385232 A, NZ 500368 A, AU 727180 B, HU 200002361 A2, MX 9909479 A1, JP 2001505924 W, KR 2001006064 A, US 6297236 B1

L7: Entry 10 of 11

File: DWPI

Oct 22, 1998

DERWENT-ACC-NO: 1998-558427

DERWENT-WEEK: 200243

COPYRIGHT 2004 DERWENT INFORMATION LTD

TITLE: Synergistic fungicidal composition for use in plant protection - contains a known 1-isoxazolyl-sulphonyl-2-halo-dioxolo:benzimidazole derivative and specified conventional fungicide compounds

INVENTOR: ASSMANN, L; DUTZMANN, S ; MAULER-MACHNIK, A ; STENZEL, K ; ASMANN, L ; ABMANN, L

PRIORITY-DATA: 1997DE-1016256 (April 18, 1997), 1999US-0402908 (October 13, 1999)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
DE 19716256 A1	October 22, 1998		034	A01N043/90
US 20020072535 A1	June 13, 2002		000	A01N043/52
WO 9847370 A1	October 29, 1998	G	000	A01N043/90
ZA 9803235 A	December 30, 1998		054	A01N000/00
AU 9875221 A	November 13, 1998		000	A01N043/90
EP 975221 A1	February 2, 2000	G	000	A01N043/90
CZ 9903697 A3	February 16, 2000		000	A01N043/90
CN 1252692 A	May 10, 2000		000	A01N043/90
BR 9809763 A	June 20, 2000		000	A01N043/90
TW 385232 A	March 21, 2000		000	A01N043/90
NZ 500368 A	September 29, 2000		000	A01N043/90
AU 727180 B	December 7, 2000		000	A01N043/90
HU 200002361 A2	November 28, 2000		000	A01N043/90
MX 9909479 A1	February 1, 2000		000	A01N043/90
JP 2001505924 W	May 8, 2001		046	A01N043/90
KR 2001006064 A	January 15, 2001		000	A01N043/90
US 6297236 B1	October 2, 2001		000	A61K031/535

2001006064 A , US 6297236 B1 INT-CL (IPC): A01 N 0/00; A01 N 37/10; A01 N 37/20;
A01 N 37/22; A01 N 37/24; A01 N 43/00; A01 N 43/30; A01 N 43/50; A01 N 43/52; A01 N
43/54; A01 N 43/64; A01 N 43/653; A01 N 43/76; A01 N 43/80; A01 N 43/82; A01 N
43/828; A01 N 43/84; A01 N 43/88; A01 N 43/90; A01 N 47/04; A01 N 47/10; A01 N
47/12; A01 N 47/14; A01 N 47/34; A01 N 47/44; A01 N 53/12; A01 N 55/02; A01 N
57/12; A61 K 31/535

Full	Title	Citation	Front	Review	Classification	Date	Reference	Abstract	IPC Class	Claims	KWIC	Draw D
------	-------	----------	-------	--------	----------------	------	-----------	----------	-----------	--------	------	--------

[Clear](#)[Generate Collection](#)[Print](#)[Fwd Refs](#)[Bkwd Refs](#)[Generate OACS](#)

Terms	Documents
L5 and L6	11

Display Format: [Change Format](#)[Previous Page](#)[Next Page](#)[Go to Doc#](#)

Hit List

Clear	Generate Collection	Print	Fwd Refs	Bkwd Refs
Generate OACS				

Search Results - Record(s) 11 through 11 of 11 returned.

☐ 11. Document ID: IL 130188 A, DE 19739982 A1, WO 9825465 A1, ZA 9711024 A, AU 9856559 A, CZ 9902086 A3, EP 944318 A1, CN 1239866 A, BR 9714390 A, HU 200000504 A2, US 6191128 B1, AU 729713 B, MX 9905065 A1, KR 2000069059 A, JP 2001505886 W, US 6303598 B1, US 6372737 B1, RU 2192743 C2, US 6509343 B1, EP 944318 B1, DE 59709805 G, MX 207803 B, ES 2192708 T3

Using default format because multiple data bases are involved.

L7: Entry 11 of 11

File: DWPI

May 12, 2004

DERWENT-ACC-NO: 1998-349826

DERWENT-WEEK: 200441

COPYRIGHT 2004 DERWENT INFORMATION LTD

TITLE: Synergistic fungicide combination for plant protection - comprising 4,6-di:phenoxy-5-fluoro-pyrimidine derivative and, e.g. tebuconazole, triadimenol, mancozeb, folpet or metalaxyl

INVENTOR: DUTZMANN, S; HEINEMANN, U ; STENZEL, K

PRIORITY-DATA: 1997DE-1005159 (February 11, 1997), 1996DE-1051217 (December 10, 1996)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
<u>IL 130188 A</u>	May 12, 2004		000	A01N043/84
<u>DE 19739982 A1</u>	June 18, 1998		045	A01N043/88
<u>WO 9825465 A1</u>	June 18, 1998	G	000	A01N043/88
<u>ZA 9711024 A</u>	August 26, 1998		067	A01N000/00
<u>AU 9856559 A</u>	July 3, 1998		000	A01N043/88
<u>CZ 9902086 A3</u>	September 15, 1999		000	A01N043/88
<u>EP 944318 A1</u>	September 29, 1999	G	000	A01N043/88
<u>CN 1239866 A</u>	December 29, 1999		000	A01N043/88
<u>BR 9714390 A</u>	May 16, 2000		000	A01N043/88
<u>HU 200000504 A2</u>	June 28, 2000		000	A01N043/88
<u>US 6191128 B1</u>	February 20, 2001		000	A61K031/535
<u>AU 729713 B</u>	February 8, 2001		000	A01N043/88
<u>MX 9905065 A1</u>	January 1, 2000		000	A01N043/88
<u>KR 2000069059 A</u>	November 25, 2000		000	A01N043/88
<u>JP 2001505886 W</u>	May 8, 2001		059	A01N043/88
<u>US 6303598 B1</u>	October 16, 2001		000	A61K031/535
<u>US 6372737 B1</u>	April 16, 2002		000	A01N043/40
<u>RU 2192743 C2</u>	November 20, 2002		000	A01N043/88
<u>US 6509343 B1</u>	January 21, 2003		000	A01N055/02

EP 944318 B1	April 9, 2003	G	000	A01N043/88
DE 59709805 G	May 15, 2003		000	A01N043/88
MX 207803 B	May 14, 2002		000	A01N043/88
ES 2192708 T3	October 16, 2003		000	A01N043/88

6303598 B1 , US 6372737 B1 INT-CL (IPC): A01 N 0/00; A01 N 43/40; A01 N 43/54; A01 N 43/64; A01 N 43/84; A01 N 43/88; A01 N 55/02; A61 K 31/515; A61 K 31/535; A61 K 31/555; C07 D 413/12; A01 N 35:06; A01 N 37:22; A01 N 37:24; A01 N 37:50; A01 N 43/88 ; A01 N 43:30; A01 N 43:36; A01 N 43:40; A01 N 43:54; A01 N 43:653; A01 N 43:84; A01 N 47:04; A01 N 47:12 ; A01 N 47:30; A01 N 47:34; A01 N 47:44; A01 N 51:00; A01 N 53:00; A01 N 55:02; A01 N 57:12; A01 N 43/88; A01 N 47:04; A01 N 47:12; A01 N 47:30; A01 N 47:34; A01 N 47:44; A01 N 51:00; A01 N 53:00; A01 N 55:02; A01 N 57:12; A01 N 43/88; A01 N 47:04; A01 N 47:12; A01 N 47:30; A01 N 47:34; A01 N 47:44; A01 N 51:00; A01 N 53:00; A01 N 55:02; A01 N 57:12; A01 N 43/88; A01 N 47:04; A01 N 47:12; A01 N 47:30; A01 N 47:34; A01 N 47:44; A01 N 51:00; A01 N 53:00; A01 N 55:02; A01 N 57:12; A01 N 37:22; A01 N 37:24; A01 N 37:50; A01 N 43/88; A01 N 43:30; A01 N 43:36; A01 N 43:40; A01 N 43:54; A01 N 43:653; A01 N 43:84; A01 N 47:04; A01 N 47:12; A01 N 47:30; A01 N 47:34; A01 N 47:44; A01 N 51:00; A01 N 53:00; A01 N 55:02; A01 N 57:12; A01 N 35:06; A01 N 37:22; A01 N 37:24; A01 N 37:50; A01 N 43/88; A01 N 43:30; A01 N 43:36; A01 N 43:40; A01 N 43:54; A01 N 43:653; A01 N 43:84; A01 N 47:04; A01 N 47:12; A01 N 47:30; A01 N 47:34; A01 N 53:12; A01 N 55:02; A01 N 57:12; A01 N 35:06; A01 N 37:22; A01 N 37:24; A01 N 37:50; A01 N 43/88; A01 N 43:30; A01 N 43:36; A01 N 43:40; A01 N 43:54; A01 N 43:653; A01 N 43:84; A01 N 47:04; A01 N 47:12; A01 N 47:30; A01 N 47:34; A01 N 47:44; A01 N 51:00; A01 N 53:00; A01 N 55:02; A01 N 57:12

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequences	Abstracts	Claims	KeyC	Draw D
------	-------	----------	-------	--------	----------------	------	-----------	-----------	-----------	--------	------	--------

Clear	Generate Collection	Print	Fwd Refs	Bkwd Refs	Generate OACS
-------	---------------------	-------	----------	-----------	---------------

Terms	Documents
L5 and L6	11

Display Format:

[Previous Page](#)

[Next Page](#)

[Go to Doc#](#)

[First Hit](#) [Previous Doc](#) [Next Doc](#) [Go to Doc#](#)

☐ [Generate Collection](#) [Print](#)

L7: Entry 1 of 11

File: PGPB

Jan 17, 2002

PGPUB-DOCUMENT-NUMBER: 20020006940
PGPUB-FILING-TYPE: new
DOCUMENT-IDENTIFIER: US 20020006940 A1

TITLE: Pesticide

PUBLICATION-DATE: January 17, 2002

INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY	RULE-47
Dutzmann, Stefan	Hilden		DE	
Erdelen, Christoph	Leichlingen		DE	
Andersch, Wolfram	Bergisch Gladbach		DE	
Dehne, Heinz-Wilhelm	Bonn		DE	
Hartwig, Jurgen	Leichlingen		DE	
Stenzel, Klaus	Dusseldorf		DE	
Kramer, Wolfgang	Burscheid		DE	

APPL-NO: 09/ 911263 [\[PALM\]](#)
DATE FILED: July 23, 2001

RELATED-US-APPL-DATA:

Application 09/911263 is a division-of US application 09/585227, filed June 1, 2000, UNKNOWN

FOREIGN-APPL-PRIORITY-DATA:

COUNTRY	APPL-NO	DOC-ID	APPL-DATE
DE	P 44 26 753.3	1994DE-P 44 26 753.3	July 28, 1994

INT-CL: [07] [A01 N 43/40](#), [A01 N 43/64](#)

US-CL-PUBLISHED: 514/341; 514/383

US-CL-CURRENT: [514/341](#); [514/383](#)

ABSTRACT:

A composition includes synergistic amounts of:

(a) a compound formula (I) 1

wherein X represents .dbd.CH-- or .dbd.N--; E represents an electron-withdrawing radical; R represents optionally substituted hetarylalkyl; A represents hydrogen, alkyl, or a bifunctional group which is linked to the radical Z; Z represents alkyl, --NH-alkyl, --N(alkyl).sub.2 or a bifunctional group which is linked to the radical A, and

(b) a fungicidal active compound, excluding cyclopropylcarboxamide derivatives and azolylmethylcycloalkanes. The combination of imidacloprid and a fungicidal active compound selected from the group consisting of tebuconazole and compounds of the formula 2

is excluded.

[Previous Doc](#)

[Next Doc](#)

[Go to Doc#](#)

[First Hit](#) [Fwd Refs](#)[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

Generate Collection

Print

L7: Entry 2 of 11

File: USPT

Jul 23, 2002

DOCUMENT-IDENTIFIER: US 6423726 B1

TITLE: Pesticide

Abstract Text (2):

wherein X represents .dbd.CH-- or .dbd.N--; E represents an electron-withdrawing radical; R represents optionally substituted hetarylalkyl; A represents hydrogen, alkyl, or a bifunctional group which is linked to the radical Z; Z represents alkyl, --NH-alkyl, --N(alkyl).sub.2 or a bifunctional group which is linked to the radical A, and (b) a fungicidal active compound, excluding cyclopropylcarboxamide derivatives and azolymethylcycloalkanes. The combination of imidacloprid and a fungicidal active compound selected from the group consisting of tebuconazole and compounds of the formula ##STR2##

Brief Summary Text (8):

Mixtures of inter alia imidacloprid- and fungicidal active compounds for use in material protection and against termites, but not for use against plant-damaging pests, are already known (EP-OS (European Published Specification) (Nit 259)). Mixtures of imidacloprid and azolymethylcycloalkanes, in particular triticonazole, are known from EP-OS (European Published Specification) 545 834.

Brief Summary Text (62):

In the examples that follow, imidacloprid is employed as active compound of the formula (I). The fungicidal active compounds also used are stated in the examples.

Detailed Description Text (7):

Mixtures of imidacloprid with tebuconazole, captan, euparen M, bitertanol, triazoxide, thiram, fludioxonil exhibit a pronounced increase in activity as compared with treatment using the individual compounds.

Detailed Description Text (14):

Mixtures of imidacloprid with euparen, guazatine, triadimenol, difenconazole, fenpiclonil exhibit a pronounced increase in activity as compared with treatment using the individual compounds.

Detailed Description Text (20):

Mixtures of imidacloprid with anilazine, benomyl, bitertanol, captan, diclofluanid, mancozeb, maneb, metalaxyl, prochloraz, procymidone, sulphate, tolylfluanid, triadimefon, triadimenol exhibit a pronounced increase in activity as compared with treatment using the individual compounds.

Detailed Description Text (26):

Mixtures of imidacloprid with bitertanol, fenpropimorph, prochloraz, tebuconazole exhibit a pronounced increase in activity as compared with treatment using the individual compounds.

Detailed Description Text (31):

Mixtures of imidacloprid with procymidone, tolylfluanid, tebuconazole exhibit a pronounced increase in activity as compared with treatment using the individual compounds.

Detailed Description Text (38):

Mixtures of imidacloprid with fenpropidin, triadimenol exhibit a pronounced increase in activity as compared with treatment using the individual compounds.

CLAIMS:

1. A composition comprising synergistic amounts of: imidacloprid; and the compound of formula XXXVIII ##STR17##
2. The composition of claim 1, wherein the composition comprises 0.1 to 10 parts by weight of compound XXXVIII per part by weight of imidacloprid.
3. The composition of claim 1, comprising 0.3 to 3 parts by weight of compound XXXVIII per part by weight of imidacloprid.
4. A process for preparing composition of claim 1, comprises mixing imidacloprid and compound XXXVIII with at least one ingredient selected from extenders, surface-active substances and mixtures thereof.
8. The process of claim 5, wherein the weight ratio of imidacloprid to compound XXXVIII is from 1:0.1 to 1:10 and the composition is applied at a concentration sufficient for controlling plant disease.

[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

[First Hit](#) [Previous Doc](#) [Next Doc](#) [Go to Doc#](#)

☐ [Generate Collection](#) [Print](#)

L7: Entry 5 of 11

File: DWPI

Jun 30, 2004

DERWENT-ACC-NO: 2002-042659

DERWENT-WEEK: 200444

COPYRIGHT 2004 DERWENT INFORMATION LTD

TITLE: Synergistic combination of fungicides for use in plant protection, comprises 2-(pyrimidinyloxy-phenyl)-2-(methoxyimino)-N-methyl-acetamide derivative and e.g. spiroxamine, quinoxifen or tebuconazole

INVENTOR: GAYER, H; MAULER-MACHNIK, A ; WACHENDORFF-NEUMANN, U ; WACHENDORF-NEUMANN, U

PATENT-ASSIGNEE: BAYER AG (FARB), BAYER CROPSCIENCE AG (FARB), BAYER LANDWIRTSCHAFTLICHE KULTURPFLANZEN (FARB), BAYER CROPSCIENCE GMBH (FARB), GAYER H (GAYEI), MAULER-MACHNIK A (MAULI), WACHENDORFF-NEUMANN U (WACHI)

PRIORITY-DATA: 2000DE-1022951 (May 11, 2000), 2002US-0275500 (November 6, 2002)

[Search Selected](#)[Search ALL](#)[Clear](#)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	PAGES	MAIN-IPC
<input type="checkbox"/> EP 1289366 B1	June 30, 2004	G	000	A01N043/54
<input type="checkbox"/> DE 10103832 A1	November 15, 2001		106	A01N043/54
<input type="checkbox"/> WO 200184931 A1	November 15, 2001	G	000	A01N043/54
<input type="checkbox"/> AU 200172383 A	November 20, 2001		000	A01N043/54
<input type="checkbox"/> EP 1289366 A1	March 12, 2003	G	000	A01N043/54
<input type="checkbox"/> CZ 200203724 A3	March 12, 2003		000	A01N043/54
<input type="checkbox"/> BR 200110699 A	March 18, 2003		000	A01N043/54
<input type="checkbox"/> KR 2002093063 A	December 12, 2002		000	A01N043/54
<input type="checkbox"/> CN 1429074 A	July 9, 2003		000	A01N043/54
<input type="checkbox"/> HU 200302025 A2	September 29, 2003		000	A01N043/54
<input type="checkbox"/> JP 2003532654 W	November 5, 2003		167	A01N043/54
<input type="checkbox"/> US 20030229087 A1	December 11, 2003		000	A01N043/58
<input type="checkbox"/> ZA 200208114 A	December 31, 2003		162	A01N000/00

DESIGNATED-STATES: AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR AE
AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK DM DZ EE ES FI GB GD
GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MA MD MG MK MN MW
MX MZ NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW AT
BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ NL OA PT SD SE SL SZ

TR TZ UG ZW AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT RO SE
SI TR

APPLICATION-DATA:

PUB-NO	APPL-DATE	APPL-NO	DESCRIPTOR
EP 1289366B1	April 30, 2001	2001EP-0951466	
EP 1289366B1	April 30, 2001	2001WO-EP04844	
EP 1289366B1		WO 200184931	Based on
DE 10103832A1	January 29, 2001	2001DE-1003832	
WO 200184931A1	April 30, 2001	2001WO-EP04844	
AU 200172383A	April 30, 2001	2001AU-0072383	
AU 200172383A		WO 200184931	Based on
EP 1289366A1	April 30, 2001	2001EP-0951466	
EP 1289366A1	April 30, 2001	2001WO-EP04844	
EP 1289366A1		WO 200184931	Based on
CZ 200203724A3	April 30, 2001	2001WO-EP04844	
CZ 200203724A3	April 30, 2001	2002CZ-0003724	
CZ 200203724A3		WO 200184931	Based on
BR 200110699A	April 30, 2001	2001BR-0010699	
BR 200110699A	April 30, 2001	2001WO-EP04844	
BR 200110699A		WO 200184931	Based on
KR2002093063A	October 22, 2002	2002KR-0714175	
CN 1429074A	April 30, 2001	2001CN-0809303	
HU 200302025A2	April 30, 2001	2001WO-EP04844	
HU 200302025A2	April 30, 2001	2003HU-0002025	
HU 200302025A2		WO 200184931	Based on
JP2003532654W	April 30, 2001	2001JP-0581607	
JP2003532654W	April 30, 2001	2001WO-EP04844	
JP2003532654W		WO 200184931	Based on
US20030229087A1	April 30, 2001	2001WO-EP04844	
US20030229087A1	November 6, 2002	2002US-0275500	
ZA 200208114A	October 9, 2002	2002ZA-0008114	

INT-CL (IPC): A01 N 0/00; A01 N 31/10; A01 N 37/18; A01 N 37/20; A01 N 37/22; A01 N 37/24; A01 N 37/34; A01 N 37/50; A01 N 37/52; A01 N 39/02; A01 N 43/08; A01 N 43/30; A01 N 43/34; A01 N 43/36; A01 N 43/38; A01 N 43/40; A01 N 43/42; A01 N 43/50; A01 N 43/52; A01 N 43/54; A01 N 43/58; A01 N 43/60; A01 N 43/653; A01 N 43/707; A01 N 43/78; A01 N 43/84; A01 N 43/86; A01 N 43/88; A01 N 43/90; A01 N 43/54; A01 N 47/04; A01 N 47/10; A01 N 47/12; A01 N 47/14; A01 N 47/18; A01 N 47/34; A01 N 47/36; A01 N 47/38; A01 N 47/44; A01 N 51/00; A01 N 53/08; A01 N 53/12; A01 N 55/00; A01 N 57/10; A01 N 57/14; A01 N 61/00; A01 N 61/00; A01 N 43/54; A01 N 61/00

ABSTRACTED-PUB-NO: DE 10103832A

BASIC-ABSTRACT:

NOVELTY - Active agent combination comprises:

(a) (2E/Z)-2-(2-((6-(3-chloro-2-methylphenoxy)-5-fluoro-4-pyrimidinyl)-oxy-)-phenyl)-2-(methoxyimino)-N-methylethanamide (I); and

(b) one or more of 82 specific fungicides, e.g. spiroxamine, quinoxyfen, tebuconazole, fenpropidin, imidacloprid, sulfur, copper oxychloride, captan or benomyl.

DETAILED DESCRIPTION - Active agent combination comprises:

(a) (2E/Z)-2-(2-((6-(3-chloro-2-methylphenoxy)-5-fluoro-4-pyrimidinyl)-oxy-)-phenyl)-2-(methoxyimino)-N-methylethanamide of formula (I), and

(b) at least one of: (1) spiroxamine, (2) quinoxyfen, (3) tebuconazole, (4) fenpropidin, (5) fenpropimorph, (6) fenoxanil, (7) chlorothalonitrile, (8) triadimefon, (9) triadimenol, (10) epiconazole, (11) metconazole, (12) fluquinconazole, (13) cyproconazole, (14) penconazole, (15) kresoxim-methyl, (16) azoxystrobin, (17) cyprodinil, (18) iminoctadine triacetate, (19) flusilazole, (20) prochloraz, (21) propiconazole, (22) bitertanol, (23) imidacloprid, (24) dichlofluanid, (25) tolylfluanid, (26) metalaxyl, (27) fenpiclonil, (28) difenconazole, (29) fludioxonil, (30) carbendazim, (31) fuberidazole, (32) enilconazole, (33) triazoxide, (34) cyfluthrin, (35) guazatine, (36) acibenzolar-S-methyl, (37) pencycuron, (38) flutolanil, (39) tricyclazole, (40) propineb, (41) procymidone, (42) mancozeb, (43) folpet, (44) dimetomorph, (45) cymoxanil, (46) fosetyl-Al, (47) famoxadone, (48) pyrimethanil, (49) mepanipyrim, (50) iprovalicarb, (51) fenhexamid, (52) carpropamid, (53) fluazinam, (54) captan, (55) quinomethionate, (56) fenamidone, (57) chlothianidin, (58) thiacloprid, (59) thiamethoxam, (60) acetamiprid, (61) picoxystrobin, (62) sulfur, (63) copper oxychloride, (64) iprodione, (65) vinclozolin, (66) phtalid, (67) edifenphos, (68) pyroquilon, (69) 2-(2-(1-chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl)-3-thioxo-3,4-dihydro-2H-1,2,4-triazole, (70) 2-chloro-5,6-(difluoromethylenedioxy)-1-(3,5-dimethylisoxazol-4-ylsulfonyl)-imidazole, (71) 4-(2-chlorophenoxy)-6-(2-(1-(5,6-dihydro-1,4,2-dioxazin-3-yl)-1-(methoxyimino)-methyl)-phenoxy)-5-fluoropyrimidine, (72) zoxamide, (73) cyamidosulfamid, (74) silthiofam, (75) trifloxystrobin, (76) N-methyl-2-(methoxyimino)-2-(2-((1-(3-trifluoromethyl-phenyl)-ethoxy)-)-iminomethyl)-phenyl)-acetamide, (77) 2-(2-((2-phenyl-2-methoxyimino-1-methyl)-1-iminooxymethyl)-phenyl)-2-methoxyimino-N-methylacetamide, (78) 2-(2-((2-(4-fluorophenyl)-2-methoxyimino-1-methylethyl)-1-iminooxymethyl)-phenyl)-2-methoxyimino-N-methylacetamide, (79) 2-(4-methoxy-3-(1-methylethoxy)-1,4-diazabuta-1,3-dienyl-oxymethyl)-phenyl-2-methoxyimino-N-methylacetamide, (80) methyl N-(2-(1-(4-chlorophenyl)-pyrazol-3-yloxy)-methyl)-phenyl)-N-methoxycarbamate, (81) 2,4-dihydro-5-methoxy-2-methyl-4-(2-(((1-(3-trifluoromethyl-phenyl)-ethylidene)-amino)-oxy)-methyl)-phenyl)-3H-1,2,4-triazol-3-one and/or (82) benomyl.

An INDEPENDENT CLAIM is included for the preparation of a fungicidal composition by mixing the combination of (a) and (b) with extenders and/or surfactants.

ACTIVITY - Fungicide.

In tests for protection of barley against Erysiphe graminis, the degree of effectiveness was 80% for (I) at 12.5 g/ha, 0% for imidacloprid at 12.5 g/ha and 100% for a combination of (I) and imidacloprid at 6.25 g/ha each.

MECHANISM OF ACTION - None given.

USE - The combination of (a) and (b) is used for controlling fungi (claimed), specifically plant pathogenic fungi such as Plasmadiophoromycetes, Oomycetes, Chytridiomycetes, Zygomycetes, Ascomycetes, Basidiomycetes and Deuteromycetes, especially Phytophthora infestans or Plasmopara viticola.

ADVANTAGE - The combination of (I) (a known fungicide described in DE19646407-A)

with one or more of the conventional fungicides (b) has a strongly synergistic effect, and is thus effective at low application rates. The combinations have very strong activity against a broad spectrum of fungi and are well tolerated by plants.

ABSTRACTED-PUB-NO: DE 10103832A
EQUIVALENT-ABSTRACTS:

CHOSEN-DRAWING: Dwg.0/0

DERWENT-CLASS: C03

CPI-CODES: C05-A01B; C05-A03A; C05-B01B; C05-B01N; C05-B02A3; C06-A02; C06-D04;
C06-D05; C06-D06; C06-D09; C06-D13; C06-E01; C06-E05; C06-F03; C06-F05; C07-A03;
C07-A04; C07-D02; C07-D04C; C07-D05; C07-D08; C07-D12; C07-D13; C07-E03; C07-E04;
C07-F01; C10-A08; C10-A12A; C10-A15; C10-A17; C10-A18; C10-D02; C14-A06;

Previous Doc

Next Doc

Go to Doc#

[First Hit](#) [Previous Doc](#) [Next Doc](#) [Go to Doc#](#)

[Generate Collection](#)[Print](#)

L7: Entry 9 of 11

File: DWPI

Oct 11, 2002

DERWENT-ACC-NO: 1998-558428

DERWENT-WEEK: 200341

COPYRIGHT 2004 DERWENT INFORMATION LTD

TITLE: Synergistic fungicidal composition for use in plant protection - contains substituted hydroxypropyl-triazole:thione derivative and conventional fungicidal compounds.

Basic Abstract Text (1):

A fungicidal composition contains:A) 2-[2-(1-chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl]-2,4-dihydro-[1,2,4]-triazole-3-thione (I); and B) one or more compounds from a triazole derivative (II), tebuconazole (III), an aniline derivative of formula $R_1-p-C_6H_4-CH(S-CCl_2-F)-SO_2-N(Me)_2$ (IV), N-[1-(4-chlorophenyl)-ethyl]-2,2-dichloro-1-ethyl-3-methyl-cyclopro-panecarboxamide (V), propineb (VI), thiocarbamate(s) (VII), teldor (VIII), a compound of formula (Me) $2CH-O-CO-NH-CH(CHMe_2)-CO-NH-CH(Me)-pC_6H_4-Me$ (IX), bendicar (X), spiroxamin (XI), azoxystrobin (XII), kresoxime methyl (XIII), compound (XIV), procymidone (XV), a pyrimidine derivative (XVI), chlorothalonil (XVII), dimetomorph (XVIII), folpet (XIX), fosetyl-Al (XX), phenylpyrrole derivative (XXI), imidacloprid (XXII), pencycuron (XXIII), benzamide derivative (XXIV) and guanidine derivative of formula $R_5-NH-(CH_2)_8-[NR_5-(CH_2)_8]_m-NH-R_5 \times (2+m) MeCOOH$ (XXV). X = Cl or phenyl; Y = CO or CHOH; $R_1 = H$ or methyl; M = Zn and/or Mn; $R_2 = methyl$ or cyclopropyl; $R_3, R_4 = Cl$ or $R_3 + R_4 = -O-CF_2-O-$; m = 0-5; $R_5 = H$ (17-23%) or $-C(NH)-NH_2$ (77-83%).

Equivalent Abstract Text (1):

A fungicidal composition contains:A) 2-[2-(1-chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl]-2,4-dihydro-[1,2,4]-triazole-3-thione (I); and B) one or more compounds from a triazole derivative (II), tebuconazole (III), an aniline derivative of formula $R_1-p-C_6H_4-CH(S-CCl_2-F)-SO_2-N(Me)_2$ (IV), N-[1-(4-chlorophenyl)-ethyl]-2,2-dichloro-1-ethyl-3-methyl-cyclopro-panecarboxamide (V), propineb (VI), thiocarbamate(s) (VII), teldor (VIII), a compound of formula (Me) $2CH-O-CO-NH-CH(CHMe_2)-CO-NH-CH(Me)-pC_6H_4-Me$ (IX), bendicar (X), spiroxamin (XI), azoxystrobin (XII), kresoxime methyl (XIII), compound (XIV), procymidone (XV), a pyrimidine derivative (XVI), chlorothalonil (XVII), dimetomorph (XVIII), folpet (XIX), fosetyl-Al (XX), phenylpyrrole derivative (XXI), imidacloprid (XXII), pencycuron (XXIII), benzamide derivative (XXIV) and guanidine derivative of formula $R_5-NH-(CH_2)_8-[NR_5-(CH_2)_8]_m-NH-R_5 \times (2+m) MeCOOH$ (XXV). X = Cl or phenyl; Y = CO or CHOH; $R_1 = H$ or methyl; M = Zn and/or Mn; $R_2 = methyl$ or cyclopropyl; $R_3, R_4 = Cl$ or $R_3 + R_4 = -O-CF_2-O-$; m = 0-5; $R_5 = H$ (17-23%) or $-C(NH)-NH_2$ (77-83%).

Equivalent Abstract Text (5):

A fungicidal composition contains:A) 2-[2-(1-chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl]-2,4-dihydro-[1,2,4]-triazole-3-thione (I); and B) one or more compounds from a triazole derivative (II), tebuconazole (III), an aniline derivative of formula $R_1-p-C_6H_4-CH(S-CCl_2-F)-SO_2-N(Me)_2$ (IV), N-[1-(4-chlorophenyl)-ethyl]-2,2-dichloro-1-ethyl-3-methyl-cyclopro-panecarboxamide (V), propineb (VI), thiocarbamate(s) (VII), teldor (VIII), a compound of formula (Me) $2CH-O-CO-NH-CH(CHMe_2)-CO-NH-CH(Me)-pC_6H_4-Me$ (IX), bendicar (X), spiroxamin (XI), azoxystrobin (XII), kresoxime methyl (XIII), compound (XIV), procymidone (XV), a pyrimidine derivative (XVI), chlorothalonil (XVII), dimetomorph (XVIII), folpet

(XIX), fosetyl-Al (XX), phenylpyrrole derivative (XXI), imidacloprid (XXII), pencycuron (XXIII), benzamide derivative (XXIV) and guanidine derivative of formula $R_5-NH-(CH_2)_8-[NR_5-(CH_2)_8]_m-NH-R_5 \times (2+m) MeCOOH$ (XXV). X = Cl or phenyl; Y = CO or CHOH; R1 = H or methyl; M = Zn and/or Mn; R2 = methyl or cyclopropyl; R3, R4 = Cl or R3 + R4 = -O-CF₂-O-; m = 0-5; R5 = H (17-23%) or -C(NH)-NH₂ (77-83%).

[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

[First Hit](#) [Previous Doc](#) [Next Doc](#) [Go to Doc#](#)

☐ [Generate Collection](#) [Print](#)

L7: Entry 10 of 11

File: DWPI

Oct 22, 1998

DERWENT-ACC-NO: 1998-558427

DERWENT-WEEK: 200243

COPYRIGHT 2004 DERWENT INFORMATION LTD

TITLE: Synergistic fungicidal composition for use in plant protection - contains a known 1-isoxazolyl-sulphonyl-2-halo-dioxolo:benzimidazole derivative and specified conventional fungicide compounds

Basic Abstract Text (1):

A fungicidal composition contains: A) a 1-(3,5-dimethyl-isoxazole-4-sulpho-nyl)-2-halo-6,6-difluoro-[1,3]-dioxolo-[4,5-f]-benzimidazole (I) [see WO9706171]; and B) one or more compounds selected from e.g. a triazole derivative of formula (II) [see DE2201063, DE2324010]; tebuconazole (III) [see EP40345]; an aniline derivative of formula R1-paraC6H4-N(S-CCl2F)-SO- 2-N(CH3)2 (IV); N-[1-(4-chlorophenyl)-ethyl]-2,2-dichloro-1-ethyl-3-methyl- -cyclopropanecarboxamide (V) [see EP341475]; propineb (VI), teldor (VIII) [see EP339418]; a compound of formula (CH3)2CH-O-CO-NH-CH(CH3)2-CO-NH- -CH(CH3)-paraC6H4-CH3 (IX) [see EP472996]; bendicar (X) [see EP313512]; spiroxamin (XI) [see EP281842]; azoxystrobin (XII) [see EP382375]; kresoxime methyl (XIII) [see EP515901]; [see DE19602095]; cymoxanil (XV); [see EP270111, EP310550]; metalaxyl (XVII), dimetomorph (XVIII), folpet (XIX), fosetyl-Al (XX), [see WO9616048]; imidacloprid (XXII), famoxadone (XXIII) [see EP393911]; [see EP600629]; and penconazole (XXVI). X = Cl or phenyl; Y = C(O) or CH(OH); R1 = H or methyl; [Those compounds not given a patent number reference may be found in 'The Pesticide Manual', 9th Edition].

Equivalent Abstract Text (1):

A fungicidal composition contains: A) a 1-(3,5-dimethyl-isoxazole-4-sulpho-nyl)-2-halo-6,6-difluoro-[1,3]-dioxolo-[4,5-f]-benzimidazole (I) [see WO9706171]; and B) one or more compounds selected from e.g. a triazole derivative of formula (II) [see DE2201063, DE2324010]; tebuconazole (III) [see EP40345]; an aniline derivative of formula R1-paraC6H4-N(S-CCl2F)-SO- 2-N(CH3)2 (IV); N-[1-(4-chlorophenyl)-ethyl]-2,2-dichloro-1-ethyl-3-methyl- -cyclopropanecarboxamide (V) [see EP341475]; propineb (VI), teldor (VIII) [see EP339418]; a compound of formula (CH3)2CH-O-CO-NH-CH(CH3)2-CO-NH- -CH(CH3)-paraC6H4-CH3 (IX) [see EP472996]; bendicar (X) [see EP313512]; spiroxamin (XI) [see EP281842]; azoxystrobin (XII) [see EP382375]; kresoxime methyl (XIII) [see EP515901]; [see DE19602095]; cymoxanil (XV); [see EP270111, EP310550]; metalaxyl (XVII), dimetomorph (XVIII), folpet (XIX), fosetyl-Al (XX), [see WO9616048]; imidacloprid (XXII), famoxadone (XXIII) [see EP393911]; [see EP600629]; and penconazole (XXVI). X = Cl or phenyl; Y = C(O) or CH(OH); R1 = H or methyl; [Those compounds not given a patent number reference may be found in 'The Pesticide Manual', 9th Edition].

Equivalent Abstract Text (5):

A fungicidal composition contains: A) a 1-(3,5-dimethyl-isoxazole-4-sulpho-nyl)-2-halo-6,6-difluoro-[1,3]-dioxolo-[4,5-f]-benzimidazole (I) [see WO9706171]; and B) one or more compounds selected from e.g. a triazole derivative of formula (II) [see DE2201063, DE2324010]; tebuconazole (III) [see EP40345]; an aniline derivative of formula R1-paraC6H4-N(S-CCl2F)-SO- 2-N(CH3)2 (IV); N-[1-(4-chlorophenyl)-ethyl]-2,2-dichloro-1-ethyl-3-methyl- -cyclopropanecarboxamide (V) [see EP341475]; propineb (VI), teldor (VIII) [see EP339418]; a compound of formula (CH3)2CH-O-CO-

NH-CH(CH(CH₃)₂)-CO-NH- -CH(CH₃)-paraC₆H₄-CH₃ (IX) [see EP472996]; bendicar (X) [see EP313512]; spiroxamin (XI) [see EP281842]; azoxystrobin (XII) [see EP382375]; kresoxime methyl (XIII) [see EP515901]; [see DE19602095]; cymoxanil (XV); [see EP270111, EP310550]; metalaxyl (XVII), dimetomorph (XVIII), folpet (XIX), fosetyl-Al (XX), [see WO9616048]; imidacloprid (XXII), famoxadone (XXIII) [see EP393911]; [see EP600629]; and penconazole (XXVI). X = Cl or phenyl; Y = C(O) or CH(OH); R₁ = H or methyl; [Those compounds not given a patent number reference may be found in ''The Pesticide Manual'', 9th Edition].

[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

[First Hit](#) [Previous Doc](#) [Next Doc](#) [Go to Doc#](#)
End of Result Set

☐ [Generate Collection](#) [Print](#)

L7: Entry 11 of 11

File: DWPI

May 12, 2004

DERWENT-ACC-NO: 1998-349826

DERWENT-WEEK: 200441

COPYRIGHT 2004 DERWENT INFORMATION LTD

TITLE: Synergistic fungicide combination for plant protection - comprising 4,6-di:phenoxy-5-fluoro-pyrimidine derivative and, e.g. tebuconazole, triadimenol, mancozeb, folpet or metalaxyl

Basic Abstract Text (3):

(B) at least 1 of antracol (propineb), euparen (dichlofluanid), euparen M (tolylfluanid), bitertanol, tebuconazole (II), triadimefon, triadimenol, imidacloprid, sumisclex, mancozeb, folpet (phaltan), dimetomorph, cymoxanil, metalaxyl, aliette (fosetyl-Al), famoxadone, pyrimethanil, cyprodinyl, mepanipyrim, kresoximmethyl, azoxystrobin, epoxiconazole, metconazole, fluquinconazole, fludioxonil, fenpiclonil, guazatine, bion, (2-methyl-1(((1-(4-methylphenyl)ethyl) amino)carbonyl)-propyl)-carboxylic acid 1-methylethyl ester, 8-t-butyl-2-(N-ethyl-N-n-propyl-amino)-methyl-1-, 4-dioxa-spiro-(5,4)-decane, 2,3-dichloro-4-(1-methylcyclohexylcarbonylami- no)-phenol, N-(R)-(1-(4-chlorophenyl)-ethyl)-2,2-dichloro-1-ethyl-3t-methy- 1-1r-cyclopropane-carboxamide, fluazinam, captan, monceren (pencycuron) and fenipiclonil.

Equivalent Abstract Text (3):

(B) at least 1 of antracol (propineb), euparen (dichlofluanid), euparen M (tolylfluanid), bitertanol, tebuconazole (II), triadimefon, triadimenol, imidacloprid, sumisclex, mancozeb, folpet (phaltan), dimetomorph, cymoxanil, metalaxyl, aliette (fosetyl-Al), famoxadone, pyrimethanil, cyprodinyl, mepanipyrim, kresoximmethyl, azoxystrobin, epoxiconazole, metconazole, fluquinconazole, fludioxonil, fenpiclonil, guazatine, bion, (2-methyl-1(((1-(4-methylphenyl)ethyl) amino)carbonyl)-propyl)-carboxylic acid 1-methylethyl ester, 8-t-butyl-2-(N-ethyl-N-n-propyl-amino)-methyl-1-, 4-dioxa-spiro-(5,4)-decane, 2,3-dichloro-4-(1-methylcyclohexylcarbonylami- no)-phenol, N-(R)-(1-(4-chlorophenyl)-ethyl)-2,2-dichloro-1-ethyl-3t-methy- 1-1r-cyclopropane-carboxamide, fluazinam, captan, monceren (pencycuron) and fenipiclonil.

Equivalent Abstract Text (10):

(B) at least 1 of antracol (propineb), euparen (dichlofluanid), euparen M (tolylfluanid), bitertanol, tebuconazole (II), triadimefon, triadimenol, imidacloprid, sumisclex, mancozeb, folpet (phaltan), dimetomorph, cymoxanil, metalaxyl, aliette (fosetyl-Al), famoxadone, pyrimethanil, cyprodinyl, mepanipyrim, kresoximmethyl, azoxystrobin, epoxiconazole, metconazole, fluquinconazole, fludioxonil, fenpiclonil, guazatine, bion, (2-methyl-1(((1-(4-methylphenyl)ethyl) amino)carbonyl)-propyl)-carboxylic acid 1-methylethyl ester, 8-t-butyl-2-(N-ethyl-N-n-propyl-amino)-methyl-1-, 4-dioxa-spiro-(5,4)-decane, 2,3-dichloro-4-(1-methylcyclohexylcarbonylami- no)-phenol, N-(R)-(1-(4-chlorophenyl)-ethyl)-2,2-dichloro-1-ethyl-3t-methy- 1-1r-cyclopropane-carboxamide, fluazinam, captan, monceren (pencycuron) and fenipiclonil.

Equivalent Abstract Text (17):

(B) at least 1 of antracol (propineb), euparen (dichlofluanid), euparen M

(tolylfluanid), bitertanol, tebuconazole (II), triadimefon, triadimenol, imidacloprid, sumisclex, mancozeb, folpet (phaltan), dimetomorph, cymoxanil, metalaxyl, aliette (fosetyl-Al), famoxadone, pyrimethanil, cyprodinyl, mepaniprim, kresoximmethyl, azoxystrobin, epoxiconazole, metconazole, fluquinconazole, fludioxonil, fenpiclonil, guazatine, bion, (2-methyl-1(((1-(4-methylphenyl)ethyl) amino)carbonyl)-propyl)-carboxylic acid 1-methylethyl ester, 8-t-butyl-2-(N-ethyl-N-n-propyl-amino)-methyl-1-, 4-dioxa-spiro-(5,4)-decane, 2,3-dichloro-4-(1-methylcyclohexylcarbonylami- no)-phenol, N-(R)-(1-(4-chlorophenyl)-ethyl)-2,2-dichloro-1-ethyl-3t-methy- 1-1r-cyclopropane-carboxamide, fluazinam, captan, monceren (pencycuron) and fenipiclonil.

[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

Hit List

Clear	Generate Collection	Print	Fwd Refs	Bkwd Refs
Generate OACS				

Search Results - Record(s) 1 through 1 of 1 returned.

☒ 1. Document ID: US 5200421 A

Using default format because multiple data bases are involved.

L10: Entry 1 of 1

File: USPT

Apr 6, 1993

US-PAT-NO: 5200421

DOCUMENT-IDENTIFIER: US 5200421 A

**** See image for Certificate of Correction ****

TITLE: Microbicidal active compound combinations

DATE-ISSUED: April 6, 1993

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Ludwig; Georg-Wilhelm	Krefeld			DE
Exner; Otto	Ratingen			DE
Paulus; Wilfried	Krefeld			DE
Buchel; Karl-Heinz	Burscheid			DE
Holmwood; Graham	Wuppertal			DE

US-CL-CURRENT: 514/383; 514/478, 514/479

Full	Title	Citation	Front	Review	Classification	Date	Reference	Supplements	Attachments	Claims	KWIC	Draw De
------	-------	----------	-------	--------	----------------	------	-----------	-------------	-------------	--------	------	---------

Clear	Generate Collection	Print	Fwd Refs	Bkwd Refs	Generate OACS
-------	---------------------	-------	----------	-----------	---------------

Terms	Documents
L5 not L7	1

Display Format:

[Previous Page](#)

[Next Page](#)

[Go to Doc#](#)

[First Hit](#) [Fwd Refs](#)[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

End of Result Set



Generate Collection

Print

L10: Entry 1 of 1

File: USPT

Apr 6, 1993

DOCUMENT-IDENTIFIER: US 5200421 A

**** See image for Certificate of Correction ****

TITLE: Microbicidal active compound combinations

Abstract Text (1):

New synergistic microbicidal active compound combinations, and the use thereof, comprising an azole fungicide, such as tebuconazole, and an iodopropargyl derivative, such as IPBC, if appropriate with the addition of other active compounds.

Brief Summary Text (100):

Nitroimino and Nitromethylenes, such as 1-[6-chloro-3-pyridinyl)-methyl]-4,5-dihydro-N-nitro-1H-imidazol-2-amin (Imidacloprid)

[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

09886197

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s imidacloprid
L2 1376 IMIDACLOPRID

=> s l2 and termites
2129 TERMITES
L3 22 L2 AND TERMITES

=> s l2 and wood
145525 WOOD
L4 25 L2 AND WOOD

=> s l3 and l4
L5 6 L3 AND L4

=> d l5 1-6 ibib hitstr abs

L5 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:695680 CAPLUS

DOCUMENT NUMBER: 137:228094

TITLE: Termiticidal baits for eliminating termite colonies

INVENTOR(S): Brode, Philip Frederick, III; Garrett, Garry Steven; Laughlin, Leo Timothy; Matthews, Randall Stryker; Barker, Dale Edwin; Kinne, Daniel James; Miller, Christopher Miles; Probst, Timothy Robert; McKibben, Gary Eugene

PATENT ASSIGNEE(S): The Procter & Gamble Company, USA

SOURCE: PCT Int. Appl., 61 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002069704	A2	20020912	WO 2002-US6200	20020301
WO 2002069704	A3	20021114		
WO 2002069704	C1	20031231		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 2002172658	A1	20021121	US 2001-799184	20010305
US 6716421	B2	20040406		
US 2003017187	A1	20030123	US 2002-172855	20020617
US 2003124166	A1	20030703	US 2002-173527	20020617
US 2003124164	A1	20030703	US 2002-268356	20021010
WO 2003105580	A1	20031224	WO 2003-US17713	20030605
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,			


LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, OM, PH, PL, PT, RO,
 RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ,
 VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
 CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
 NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
 GW, ML, MR, NE, SN, TD, TG

WO 2003106395 A1 20031224 WO 2003-US17714 20030605
 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
 CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GM,
 HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
 LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, OM, PH, PL, PT, RO,
 RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ,
 VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
 CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
 NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
 GW, ML, MR, NE, SN, TD, TG

WO 2004032625 A2 20040422 WO 2003-US32092 20031007
 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
 CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
 GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
 LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,
 PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ,
 UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU,
 TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
 CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
 NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
 GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.: US 2001-799184 A 20010305
 US 2002-172855 A 20020617
 US 2002-173527 A 20020617
 US 2002-268356 A 20021010

OTHER SOURCE(S): MARPAT 137:228094
 GI



AB This invention relates to devices, kits, and methods for eliminating termite colonies. The kits, devices, and methods employ a termiticidal bait matrix contain (a) a termiticide (I, X = nil, a hydrocarbon group, O or NR₈, R₉ where R₈ and R₉ are H or a hydrocarbon group; X₁ = CH, a carbon atom or a heteroatom; R, R₁, R₂, R₃ = H or OH and if R₄ and R₅ are O and R₆ and R₇ are H then R, R₁, R₂ and R₃ may be C₁-6; R₄ and R₅ are H, O or N; R₉ and R₁₀ are nil, C₁-6, and amides) selected such that the termiticide causes death to about 50 to about 100% of **termites** within about 24 to about 84 days after the **termites** begin to ingest the termiticide or the bait matrix comprising the termiticide, (b) a cellulose containing material, and (c) water. The termiticidal bait matrix can be used in a bait station installed in the ground. The kits are suitable to be

used by consumers in their homes.

L5 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:547214 CAPLUS
 DOCUMENT NUMBER: 137:105178
 TITLE: Termiticides containing 2-pyridinethiol-1-oxide salts and **wood** and polymers containing the termiticides
 INVENTOR(S): Nishimoto, Koichi; Sato, Toshio; Suga, Mamoru
 PATENT ASSIGNEE(S): Yoshitomi Fine Chemicals Ltd., Japan
 SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.
 CODEN: JKXXAF
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002205906	A2	20020723	JP 2001-337124	20010926
PRIORITY APPLN. INFO.:			JP 2000-381082	A 20001108

AB The termiticides, which are effective on **termites**, bark beetles, etc., and environmentally safe, contain (a) ≥ 1 selected from Cu, Zn, and Na salts of 2-pyridine-1-oxide and optionally (b) ≥ 1 selected from pyrethroids, nicotinoids, organophosphorus compds., isocyanuric acid compds., carbamates, acetamiprid, and inorg. boric acid compds. **Wood** and polymers containing the termiticides are also claimed. A **wood** block was coated with DMSO solution containing Cu pyrrithione and **imidacloprid** and dried at room temperature for ≥ 20 days. The **wood** block.

L5 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:720924 CAPLUS
 DOCUMENT NUMBER: 135:340463
 TITLE: Chemical prevention of colony foundation by *Cryptotermes brevis* (Isoptera: Kalotermitidae) in attic modules
 AUTHOR(S): Scheffrahn, Rudolf H.; Busey, Philip; Edwards, Jeffrey K.; Krecek, Jan; Maharajh, Boudanath; Su, Nan-Yao
 CORPORATE SOURCE: Ft. Lauderdale Research and Education Center, University of Florida, Fort Lauderdale, FL, 33314, USA
 SOURCE: Journal of Economic Entomology (2001), 94(4), 915-919
 CODEN: JEENAI; ISSN: 0022-0493
 PUBLISHER: Entomological Society of America
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Disodium octaborate tetrahydrate (DOT) dust, DOT aqueous solution, **imidacloprid** dust, and amorphous silica gel dust with synergized 1% pyrethrins were applied on **wood** surfaces to simulated attic modules. Modules (30 by 30 cm) with and without fiberglass insulation were exposed to dispersal flights of *Cryptotermes brevis* (Walker) in May and June of 1998 and 1999. Six months after flights, modules were disassembled and inspected for nuptial chamber location and contents. During both years, air and water control treatments contained 22.2 ± 9.94 (mean \pm SD) nuptial chambers, 7.5 ± 5.7 live imagos, and 2.0 ± 1.4 chambers with brood. This survivorship indicated that the attic modules performed well as a colonizing platform for *C. brevis*. *C. brevis* dealates preferred constructing nuptial chambers in the crevices at the bases or tops of the modules instead of internal crevices. Modules treated in 1998 and 1999 with DOT or silica dusts contained no live

termites, whereas zero of five modules treated with **imidacloprid** dust in 1998 and two of 20 modules treated with **imidacloprid** dust in 1999 contained single live incipient colonies. In 1998, 15% DOT solution, applied as a postconstruction treatment, yielded significantly fewer chambers and live **termites** than controls, but was not as effective as dusts in preventing successful colonization. In 1999, the DOT solution, applied as a construction-phase treatment, was equally as effective in preventing colonization as the dust treatments during that year. Results indicate that dust formulations of DOT, silica gel, and **imidacloprid** can be used to prevent drywood termite colonization in existing building voids and attics. Where the entire **wood** framing is exposed to treatment, such as during building construction, aqueous DOT solution can be equally effective as dusts in preventing colonization by *C. brevis*.

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:467897 CAPLUS
DOCUMENT NUMBER: 133:85609
TITLE: Termiticidal baits comprising nonhygroscopic agents in hygroscopic containers
INVENTOR(S): Minakawa, Fumiyasu; Uchida, Yuki
PATENT ASSIGNEE(S): Yuko Chemical Industries Co., Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000189031	A2	20000711	JP 1998-369335	19981225
PRIORITY APPLN. INFO.:			JP 1998-369335	19981225

AB A nonhygroscopic agent for controlling **termites** (e.g. diflubenzuron) is housed in a container which is made of an edible hygroscopic material (cellulosic cloth, polyvinyl alc. film). Thus, sulfluramid 0.001 and pine oil 1% (attractant) were dissolved in polyethylene glycol, and the solution was made to soaked into **wood** flour at a 25/100 weight ratio. The agent was heat sealed in an envelope (15 + 7 cm) made of nonwoven fabric of cellulose fibers with 1% by weight added pine oil to obtain a bait with satisfactory attractiveness to *Reticulitermes*.

L5 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:125767 CAPLUS
DOCUMENT NUMBER: 130:178773
TITLE: Composition for the control of **wood** -destroying insects, especially **termites**
INVENTOR(S): Anderson, John-phillip-evans; Keuken, Oliver
PATENT ASSIGNEE(S): Bayer A.-G., Germany
SOURCE: Eur. Pat. Appl., 21 pp.
CODEN: EPXXDW
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 896791	A2	19990217	EP 1998-114187	19980729
EP 896791	A3	20000112		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
DE 19734665	A1	19990218	DE 1997-19734665	19970811
TW 505500	B	20021011	TW 1998-87112592	19980731
US 6264968	B1	20010724	US 1998-128818	19980804
ZA 9807118	A	19990209	ZA 1998-7118	19980807
JP 11124302	A2	19990511	JP 1998-234861	19980807
AU 9879895	A1	19990218	AU 1998-79895	19980811
AU 768390	B2	20031211		
BR 9803138	A	19991221	BR 1998-3138	19980811
PRIORITY APPLN. INFO.:		DE 1997-19734665 A 19970811		
AB The title compns. (no examples) comprise an insecticide, preferably imidacloprid , incorporated into an organic natural and/or synthetic carrier. Optional ingredients are insect attractants and microbicides.				

L5 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1993:54353 CAPLUS
DOCUMENT NUMBER: 118:54353
TITLE: Imidazolidine derivatives and related compounds as industrial insecticides and wood preservatives
INVENTOR(S): Tsuboi, Shinichi; Sone, Shinzaburo; Obinata, Toru; Exner, Otto; Schwamborn, Michael
PATENT ASSIGNEE(S): Nihon Bayer Agrochem K. K., Japan
SOURCE: Eur. Pat. Appl., 15 pp.
CODEN: EPXXDW
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 511541	A1	19921104	EP 1992-106384	19920414
EP 511541	B1	19960904		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
JP 05032505	A2	19930209	JP 1991-350751	19911212
JP 3162450	B2	20010425		
JP 2001031511	A2	20010206	JP 2000-233512	19911212
AU 9213908	A1	19921029	AU 1992-13908	19920330
AU 645672	B2	19940120		
AT 142077	E	19960915	AT 1992-106384	19920414
ES 2090400	T3	19961016	ES 1992-106384	19920414
BR 9201534	A	19921201	BR 1992-1534	19920427
US 6323224	B1	20011127	US 1995-543351	19951016
US 2001051643	A1	20011213	US 2001-886197	20010621
PRIORITY APPLN. INFO.:		JP 1991-125172 A 19910427		
		JP 1991-350751 A 19911212		
		US 1992-872279 B1 19920422		
		US 1995-543351 A3 19951016		
OTHER SOURCE(S):		MARPAT 118:54353		
GI				

09886197



AB The imidazolidine derivs. and related compds. I (X = NH, S; Y = CH, N; Z = 2-chloro-5-pyridyl, 2-chloro-5-thiazolyl; R = H, Me; n = 0, 1) are industrial insecticides and wood preservatives. Wood impregnated with 0.32 ppm imidacloprid was lethal to termites (Coptotermes formosanus) for ≥ 3 wk.

=> d his

(FILE 'HOME' ENTERED AT 16:06:53 ON 21 JUL 2004)

FILE 'STNGUIDE' ENTERED AT 16:07:06 ON 21 JUL 2004

FILE 'REGISTRY' ENTERED AT 16:08:17 ON 21 JUL 2004

L1 0 S IMIDALCLOPRID

FILE 'CAPLUS' ENTERED AT 16:09:11 ON 21 JUL 2004

L2 1376 S IMIDACLOPRID

L3 22 S L2 AND TERMITES

L4 25 S L2 AND WOOD

L5 6 S L3 AND L4

=> d l3 1-22 ibib hitstr abs

L3 ANSWER 1 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:892507 CAPLUS

DOCUMENT NUMBER: 139:360411

TITLE: Naphthalenic compounds as termite bait toxicants

INVENTOR(S): Rojas, Maria Guadalupe; Morales-Ramos, Juan A.; Green, Frederick, III

PATENT ASSIGNEE(S): The United States of America, as Represented by the Secretary of Agriculture, USA

SOURCE: PCT Int. Appl., 17 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003092375	A2	20031113	WO 2003-US13457	20030430
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,			

CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG

US 6691453 B1 20040217 US 2002-135224 20020430

PRIORITY APPLN. INFO.: US 2002-135224 A 20020430

AB A matrix suitable to be used as baits and attractants for **termites** comprises cellulose, naphthalenic compds., water, and potentially other termite-preferred nutrients. Methods of monitoring the presence of **termites** using such matrixes and methods of controlling **termites** using such matrixes to deliver termite toxicants (e.g., streptomycin sulfate or **imidacloprid**) are also provided. Thus, N-hydroxynaphthalimide sodium salt (I) was incorporated into a bait matrix containing lecithin, ergosterol, EtOH, yeast hydrolyzate, and cellulose. I at 500 ppm was sufficient to induce mortality of Formosan subterranean termite (*Coptotermes formosanus*) within .apprx.2 mo without any repellency to **termites**.

L3 ANSWER 2 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:690211 CAPLUS

DOCUMENT NUMBER: 139:334278

TITLE: Evaluation of chemical control measures for **termites** in maize

AUTHOR(S): Riekert, H. F.; Van den Berg, J.

CORPORATE SOURCE: ARC-Grain Crops Institute, Potchefstroom, 2520, S. Afr.

SOURCE: South African Journal of Plant and Soil (2003), 20(1), 1-5

CODEN: SAJSEV; ISSN: 0257-1862

PUBLISHER: Forum Press International

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Field trails were conducted from the 1994/95 to 2000/2001 growing seasons to evaluate various insecticides for preventative and corrective control of the fungus-growing **termites**, *Microtermes* sp., *Odontotermes* sp. and *Allodotermes* sp. in maize. The incidence of lodged maize plants was used as criteria for insecticide efficacy. Carbofuran GR, **imidacloprid** WS, chlorpyrifos GR and fipronil GR were evaluated as preventative treatments. Corrective treatments in the form of spray applications of the systemic insecticides carbosulfan EC, benfuracarb EC and **imidacloprid** SL were also evaluated. Treatments were applied to the basal 25 cm of maize stems and to the soil surface surrounding plants. **Imidacloprid** spray applications generally provided good control of **termites**. The optimum plant growth stage for **imidacloprid** application was during the pre-flowering stage, 6 to 10 wk after plant emergence. Pre-flowering applications were usually more effective in limiting damage than post-flowering applications. The granular insecticide, fipronil, showed promise for termite control. Chlorpyrifos granules, applied as a side dressing four weeks after plant emergence, significantly reduced lodging. Two novel control methods (fishmeal and diesel fuel) on the soil surface resulted in suppression of termite damage and subsequent reduction in lodging of plants. In the majority of trials total yields (lodged and upright plants) did not differ over insecticide treatments. However, the proportion of the total yield that had to be hand-harvested from lodged plants ranged from 0 to 41%, and was significantly higher in ineffective treatments. This resulted in increased production costs and uneconomic maize production

REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 3 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

09886197

ACCESSION NUMBER: 2003:177184 CAPLUS
DOCUMENT NUMBER: 138:333176
TITLE: Effect of **imidacloprid** tree treatments on the occurrence of formosan subterranean **termites**, *Coptotermes formosanus* Shiraki (Isoptera: Rhinotermitidae), in independent monitors
AUTHOR(S): Osbrink, Weste L. A.; Lax, Alan R.
CORPORATE SOURCE: Southern Regional Research Center, USDA-ARS, New Orleans, LA, 70124, USA
SOURCE: Journal of Economic Entomology (2003), 96(1), 117-125
CODEN: JEENAI; ISSN: 0022-0493
PUBLISHER: Entomological Society of America
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Periodic sampling of 87 independent monitors, initially active with the Formosan subterranean termite, *Coptotermes formosanus* Shiraki, was conducted. Monitors, located in eight sectors adjacent to seven buildings, were various distances (1-46 m) from 57 trees treated with 0.1% **imidacloprid** foam. **Termites** collected from six of the eight sectors showed latent mortality attributed to **imidacloprid** intoxication at all monitor-tree distances. Approx. 6 mo after treatment, termite populations had recovered in these sectors. Another sector showed termite population suppression for ~15 mo, followed by recovery. **Imidacloprid** tree treatments did not control *C. formosanus* populations in independent monitors adjacent to the treatments.
REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 4 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:54728 CAPLUS
DOCUMENT NUMBER: 138:333162
TITLE: Comparative evaluation of chemical and botanical insecticides against **termites**
AUTHOR(S): Singh, S. K.; Singh, G.
CORPORATE SOURCE: Indian Institute of Pulses Research, Kanpur, 208024, India
SOURCE: Entomon (2002), 27(2), 153-160
CODEN: ENTOD5; ISSN: 0377-9335
PUBLISHER: Association for Advancement of Entomology
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Insecticides viz., **Imidacloprid** 17.8 SL, chlorpyrifos 20 EC, lindane 20 EC, endosulfan 35 EC, cypermethrin 10 EC and phorate 10G and neem manure were tested against **termites** in pots. **Imidacloprid** 0.012% was effective up to 3 mo but at 0.008 and 0.004% were effective up to 2 mo only. Chlorpyrifos at 0.04% was effective up to 2 mo but at 0.02 and 0.03% were effective up to one month only. Lindane at 0.03 and 0.04% and endosulfan at 0.08% were effective up to one month. All the above insecticides gave above 50% corrected mortality. Lindane 0.02%, endosulfan 0.07%, neem manure 50 g per pot, phorate 0.1 g a.i. per pot and cypermethrin 0.0025% were found least effective. Among botanical insecticides, Nimbecidine and Nemactin were effective up to two months while Rakshak, Multineem, Neemgourd and Vanguard were effective for short time up to one month. Field trial was conducted in mango orchards of Upeda, Ghaziabad and Rohenda, Bulandshahar, Uttar Pradesh, India. **Imidacloprid** 0.012%, chlorpyrifos 0.04% and lindane 0.04% were found most effective and gave 100% reduction in termite population up to five months. **Imidacloprid** 0.004%, chlorpyrifos 0.02%, lindane 0.02%, lindane 1.3% dust @ 100 g per tree and neem manure 500 g per tree were found less effective.

09886197

REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 5 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 2003:14144 CAPLUS
DOCUMENT NUMBER: 138:40461
TITLE: Manufacture of additive-containing prefoamed polymer particles
INVENTOR(S): Maeda, Tadanobu
PATENT ASSIGNEE(S): Mitsubishi Chemical Foam Plastic Corp., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 7 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003001627	A2	20030108	JP 2001-193245	20010626
PRIORITY APPLN. INFO.:			JP 2001-193245	20010626

AB The particles are manufactured by heating expandable polymer particles in a prefoaming apparatus under stirring and adding plastic additives to the prefoaming polymer particles. Thus, 600 g Styropor JF 200 (polystyrene expandable particle) was prefoamed, mixed with 0.1 part imidacloprid at expansion ratio 2, and further expanded to give prefoamed particles (expansion ratio 50), which were molded to give a plastic foam molding with compressive strength at 5% strain (JIS A 9511) 118 kPa, bending strength 273 kPa, d. 20.3 g/L, and reduced damage caused by termites.

L3 ANSWER 6 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 2002:695680 CAPLUS
DOCUMENT NUMBER: 137:228094
TITLE: Termiticidal baits for eliminating termite colonies
INVENTOR(S): Brode, Philip Frederick, III; Garrett, Garry Steven; Laughlin, Leo Timothy; Matthews, Randall Stryker; Barker, Dale Edwin; Kinne, Daniel James; Miller, Christopher Miles; Probst, Timothy Robert; McKibben, Gary Eugene
PATENT ASSIGNEE(S): The Procter & Gamble Company, USA
SOURCE: PCT Int. Appl., 61 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002069704	A2	20020912	WO 2002-US6200	20020301
WO 2002069704	A3	20021114		
WO 2002069704	C1	20031231		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH,

CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR,
 BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

US 2002172658 A1 20021121 US 2001-799184 20010305
 US 6716421 B2 20040406
 US 2003017187 A1 20030123 US 2002-172855 20020617
 US 2003124166 A1 20030703 US 2002-173527 20020617
 US 2003124164 A1 20030703 US 2002-268356 20021010
 WO 2003105580 A1 20031224 WO 2003-US17713 20030605

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
 CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GM,
 HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
 LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, OM, PH, PL, PT, RO,
 RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ,
 VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
 CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
 NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
 GW, ML, MR, NE, SN, TD, TG

WO 2003106395 A1 20031224 WO 2003-US17714 20030605


W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
 CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GM,
 HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
 LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, OM, PH, PL, PT, RO,
 RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ,
 VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
 CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
 NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
 GW, ML, MR, NE, SN, TD, TG

WO 2004032625 A2 20040422 WO 2003-US32092 20031007

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
 CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
 GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
 LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,
 PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ,
 UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU,
 TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
 CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
 NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
 GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.: US 2001-799184 A 20010305
 US 2002-172855 A 20020617
 US 2002-173527 A 20020617
 US 2002-268356 A 20021010

OTHER SOURCE(S): MARPAT 137:228094
 GI



AB This invention relates to devices, kits, and methods for eliminating

termite colonies. The kits, devices, and methods employ a termiticidal bait matrix contain (a) a termiticide (I, X = nil, a hydrocarbon group, O or NR₈, R₉ where R₈ and R₉ are H or a hydrocarbon group; X₁ = CH, a carbon atom or a heteroatom; R, R₁, R₂, R₃ = H or OH and if R₄ and R₅ are O and R₆ and R₇ are H then R, R₁, R₂ and R₃ may be C₁-6; R₄ and R₅ are H, O or N; R₉ and R₁₀ are nil, C₁-6, and amides) selected such that the termiticide causes death to about 50 to about 100% of **termites** within about 24 to about 84 days after the **termites** begin to ingest the termiticide or the bait matrix comprising the termiticide, (b) a cellulose containing material, and (c) water. The termiticidal bait matrix can be used in a bait station installed in the ground. The kits are suitable to be used by consumers in their homes.

L3 ANSWER 7 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:547214 CAPLUS
DOCUMENT NUMBER: 137:105178
TITLE: Termiticides containing 2-pyridinethiol-1-oxide salts and wood and polymers containing the termiticides
INVENTOR(S): Nishimoto, Koichi; Sato, Toshio; Suga, Mamoru
PATENT ASSIGNEE(S): Yoshitomi Fine Chemicals Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002205906	A2	20020723	JP 2001-337124	20010926
PRIORITY APPLN. INFO.: JP 2000-381082 A 20001108				

AB The termiticides, which are effective on **termites**, bark beetles, etc., and environmentally safe, contain (a) ≥ 1 selected from Cu, Zn, and Na salts of 2-pyridine-1-oxide and optionally (b) ≥ 1 selected from pyrethroids, nicotinoids, organophosphorus compds., isocyanuric acid compds., carbamates, acetamiprid, and inorg. boric acid compds. Wood and polymers containing the termiticides are also claimed. A wood block was coated with DMSO solution containing Cu pyrithione and **imidacloprid** and dried at room temperature for ≥ 20 days. The wood block.

L3 ANSWER 8 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:317186 CAPLUS
DOCUMENT NUMBER: 136:365273
TITLE: Effect of insecticide treatments against **termites** on yield and quality of sugarcane
AUTHOR(S): Singh, Manager; Singh, N. B.
CORPORATE SOURCE: Sugarcane Research Institute, Shahjahanpur, 242 001, India
SOURCE: Sugar Cane International (2002), (March/April), 27-29
CODEN: SCINFQ; ISSN: 1468-6031
PUBLISHER: Agra Europe (London) Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English
AB In a field experiment in 1995-97 at three sites in Uttar Pradesh sugarcane cv. Cos 767 setts were treated with several insecticides for the control of **termites**. Mean cane yields were highest with treatment with 0.20% solution **imidacloprid** 70 ws (77.8 t/ha), 2.5 kg ai/ha phorate 10 G (76.1 t), 2.5 kg ai/ha chlorpyrifos 15 G (73.9 t) and 1 kg ai/ha chlorpyrifos 20 EC (73.5 t) compared with the control yield of 54.4 t.

Cane juice sucrose content was highest with 0.20% solution
imidacloprid 70 WS (17.53%) compared with the control of 14.96%.

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 9 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:720924 CAPLUS
 DOCUMENT NUMBER: 135:340463
 TITLE: Chemical prevention of colony foundation by
Cryptotermes brevis (Isoptera: Kalotermitidae) in
 attic modules
 AUTHOR(S): Scheffrahn, Rudolf H.; Busey, Philip; Edwards, Jeffrey
 K.; Krecek, Jan; Maharajh, Boudanath; Su, Nan-Yao
 CORPORATE SOURCE: Ft. Lauderdale Research and Education Center,
 University of Florida, Fort Lauderdale, FL, 33314, USA
 SOURCE: Journal of Economic Entomology (2001), 94(4), 915-919
 CODEN: JEENAI; ISSN: 0022-0493
 PUBLISHER: Entomological Society of America
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Disodium octaborate tetrahydrate (DOT) dust, DOT aqueous solution,
imidacloprid dust, and amorphous silica gel dust with synergized
 1% pyrethrins were applied on wood surfaces to simulated attic modules.
 Modules (30 by 30 cm) with and without fiberglass insulation were exposed
 to dispersal flights of *Cryptotermes brevis* (Walker) in May and June of
 1998 and 1999. Six months after flights, modules were disassembled and
 inspected for nuptial chamber location and contents. During both years,
 air and water control treatments contained 22.2 ± 9.94 (mean \pm SD)
 nuptial chambers, 7.5 ± 5.7 live imagoes, and 2.0 ± 1.4 chambers with
 brood. This survivorship indicated that the attic modules performed well
 as a colonizing platform for *C. brevis*. *C. brevis* dealates preferred
 constructing nuptial chambers in the crevices at the bases or tops of the
 modules instead of internal crevices. Modules treated in 1998 and 1999
 with DOT or silica dusts contained no live **termites**, whereas
 zero of five modules treated with **imidacloprid** dust in 1998 and
 two of 20 modules treated with **imidacloprid** dust in 1999
 contained single live incipient colonies. In 1998, 15% DOT solution, applied
 as a postconstruction treatment, yielded significantly fewer chambers and
 live **termites** than controls, but was not as effective as dusts
 in preventing successful colonization. In 1999, the DOT solution, applied as
 a construction-phase treatment, was equally as effective in preventing
 colonization as the dust treatments during that year. Results indicate
 that dust formulations of DOT, silica gel, and **imidacloprid** can
 be used to prevent drywood termite colonization in existing building voids
 and attics. Where the entire wood framing is exposed to treatment, such
 as during building construction, aqueous DOT solution can be equally effective

as

dusts in preventing colonization by *C. brevis*.

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 10 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:336305 CAPLUS
 DOCUMENT NUMBER: 135:1645
 TITLE: Effects of sublethal exposure to **imidacloprid**
 on subsequent behavior of subterranean termite
Reticulitermes virginicus (Isoptera: Rhinotermitidae)
 AUTHOR(S): Thorne, Barbara L.; Breisch, Nancy L.
 CORPORATE SOURCE: Department of Entomology, University of Maryland,
 College Park, MD, 20742, USA

09886197

SOURCE: Journal of Economic Entomology (2001), 94(2), 492-498
CODEN: JEENAI; ISSN: 0022-0493
PUBLISHER: Entomological Society of America
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Expts. were conducted to determine whether subterranean **termites**, *Reticulitermes virginicus* (Banks), previously exposed to sublethal doses of **imidacloprid** (Premise), and allowed to recover for 1 wk, demonstrated behavioral aversion to a subsequent exposure. Worker **termites** experiencing a previous sublethal but debilitating exposure to **imidacloprid**-treated sand (either 10 or 100 ppm for 4 h) showed no apparent aversion to a second encounter with **imidacloprid**-treated sand under conditions of this experiment. If these laboratory results hold in the field and **termites** traveling through a zone of soil treated with **imidacloprid** are impaired but subsequently recover, they will be just as likely as their naive nestmates to reenter the treated area if their travels take them through the nonrepellent application a second time. Thus, a sublethal exposure to **imidacloprid** can affect termite tunneling behavior. Many worker **termites** that received an initial 4-h exposure to 100 ppm **imidacloprid**-treated sand died, but those that survived tunneled significantly less than did their naive nestmates, as did some **termites** exposed to 10 ppm **imidacloprid**.

REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 11 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:283714 CAPLUS

DOCUMENT NUMBER: 134:276894

TITLE: Nonedible foraging matrix insert for subterranean termite control

INVENTOR(S): Koehler, Philip G.; Oi, Faith M.

PATENT ASSIGNEE(S): University of Florida, USA; United States of America, as Represented by the Secretary of Agriculture

SOURCE: PCT Int. Appl., 30 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001026456	A1	20010419	WO 2000-US6591	20000314
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
AU 758489	B2	20030320	AU 2000-37432	20000314
AU 2000037432	A5	20010423		
JP 2004500043	T2	20040108	JP 2001-529256	20000314

PRIORITY APPLN. INFO.: US 1999-159266P P 19991013
WO 2000-US6591 W 20000314

AB A several step process starts with taking a tube with a removal cap at one end, such as a two to four inch PVC tube, and filling the inner chamber

with a food source such as rolled cardboard. The tube is then placed with its open end adjacent to a termite population, so that live **termites** can then enter the entrance/exit of the tube to reach the food source. Once **termites** are inside the tube, the cap is removed from the tube, and a nonedible foraging matrix, such as a disk of loose soil and or sand that is treated with a slow acting and nonrepellent toxicant, is placed between the food source in the chamber and the termite entrance/exit of the chamber. Slow-acting and non-repellent toxicants can be fipronil, chlorfenapyr, **imidacloprid**, and chlorpyrifos. The **termites** are then forced to pass through and disperse the slow-acting and non-repellent toxicant on soil particles or other nonedible foraging matrixes through their tunnels and living space in order to kill **termites**. **Termites** that contact tunnels and living space contaminated with the treated nonedible foraging matrix particles die over time.

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 12 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:573349 CAPLUS

DOCUMENT NUMBER: 133:248356

TITLE: Feeding inhibition and mortality in *Reticulitermes flavipes* (Isoptera: Rhinotermitidae) after exposure to **imidacloprid**-treated soils

AUTHOR(S): Ramakrishnan, Rathna; Suiter, Daniel R.; Nakatsu, Cindy H.; Bennett, Gary W.

CORPORATE SOURCE: Center for Urban & Industrial Pest Management, Department of Entomology, Purdue University, West Lafayette, IN, 47907, USA

SOURCE: Journal of Economic Entomology (2000), 93(2), 422-428
CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Feeding inhibition and mortality of *Reticulitermes flavipes* (Kollar) exposed to sand, sandy loam, loam, and silty clay loam soils treated with several concns. of **imidacloprid** were studied using bioassay techniques under laboratory conditions. Termite workers stopped feeding after exposure to treated soils. Differences in feeding reduction varied among the soil types. Based on the magnitude of the F-statistics, the effect of **imidacloprid** on the reduction of termite feeding was greatest in sand followed by sandy loam, loam, and silty clay loam soils. Soil properties such as organic matter content, silt and clay proportions, pH, and cation exchange capacity were suggested to affect the bioavailability of **imidacloprid**. Similar soil effects on mortality were observed in **termites** continuously exposed to treated soil for 21 days. In 3 of 4 soils tested, susceptibility to **imidacloprid** was not affected by the source of the **termites** tested.

REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 13 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:467897 CAPLUS

DOCUMENT NUMBER: 133:85609

TITLE: Termiticidal baits comprising nonhygroscopic agents in hygroscopic containers

INVENTOR(S): Minakawa, Fumiyasu; Uchida, Yuki

PATENT ASSIGNEE(S): Yuko Chemical Industries Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKXXAF

09886197

DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000189031	A2	20000711	JP 1998-369335	19981225
PRIORITY APPLN. INFO.:			JP 1998-369335	19981225

AB A nonhygroscopic agent for controlling **termites** (e.g. diflubenzuron) is housed in a container which is made of an edible hygroscopic material (cellulosic cloth, polyvinyl alc. film). Thus, sulfluramid 0.001 and pine oil 1% (attractant) were dissolved in polyethylene glycol, and the solution was made to soaked into wood flour at a 25/100 weight ratio. The agent was heat sealed in an envelope (15 + 7 cm) made of nonwoven fabric of cellulose fibers with 1% by weight added pine oil to obtain a bait with satisfactory attractiveness to *Reticulitermes*.

L3 ANSWER 14 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:52160 CAPLUS

DOCUMENT NUMBER: 132:133596

TITLE: Degradation of bifenthrin, chlorpyrifos and **imidacloprid** in soil and bedding materials at termiticidal application rates

AUTHOR(S): Baskaran, Sundaram; Kookana, Rai S.; Naidu, Ravendra

CORPORATE SOURCE: CSIRO Land and Water, Glen Osmond, 5064, Australia

SOURCE: Pesticide Science (1999), 55(12), 1222-1228

CODEN: PSSCBG; ISSN: 0031-613X

PUBLISHER: John Wiley & Sons Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Organophosphorus, pyrethroid and chloronicotinyl insecticides have been used to control **termites** in building structures in recent years. The degradation behavior of three insecticides (bifenthrin, chlorpyrifos and **imidacloprid**) at termiticidal application rates was studied under standard laboratory conditions (25°C, 60% field moisture capacity and darkness) for 24 mo. The study was carried out on one soil and two bedding materials (sand-dolomite and quarry sand), which are commonly used under housing in Australia. Expts. were also conducted to examine the effect of soil moisture on the degradation of these insecticides. Insecticide residues in the samples collected at different days after application were measured by HPLC. The rate of degradation of bifenthrin and **imidacloprid** insecticides was adequately described by a first-order kinetic model ($r^2=0.93-0.97$). However, chlorpyrifos degradation was biphasic, showing an initial faster degradation followed by a slower rate. Therefore, the degradation data during the slower phase only (after a two-month period) followed the first-order law ($r^2=0.95$). Soil moisture had little effect on degradation of **imidacloprid** and bifenthrin. Among the three insecticides, bifenthrin and **imidacloprid** were most stable and chlorpyrifos the least. Chlorpyrifos showed a major loss (75-90%) of residue during the 24 mo incubation period. In the bedding materials, simultaneous accumulation of the primary metabolite of chlorpyrifos, TCP (3,5,6-trichloro-2-pyridinol) was observed. Hydrolysis appeared to have caused the observed rapid loss of chlorpyrifos, especially in the highly alkaline bedding materials (sand-dolomite and quarry sand).

REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 15 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

09886197

ACCESSION NUMBER: 1999:799698 CAPLUS
DOCUMENT NUMBER: 132:9953
TITLE: Termite control
INVENTOR(S): De Villiers, Vivian; Van der Westhuizen, M. C.;
Robbertse, Ernest
PATENT ASSIGNEE(S): Bayer A.-G., Germany
SOURCE: S. African, 16 pp.
CODEN: SFXXAB
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
ZA 9711701	A	19980706	ZA 1997-11701	19971230
AP 1174	A	20030630	AP 1998-1424	19981228
W: BW, GH, GM, KE, LS, MW, SD, SZ, UG, ZM, ZW				
BR 9805735	A	20010424	BR 1998-5735	19981229
PRIORITY APPLN. INFO.:		ZA 1997-11701	A	19971230

AB Agonists or antagonists of nicotinergeric acetylcholine receptors of insects are used for the control of harvester **termites**, i.e. Hodotermitidae. **Imidacloprid** is the preferred active ingredient. The bait formulations comprise lucerne or grass particles.

L3 ANSWER 16 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:797191 CAPLUS
DOCUMENT NUMBER: 132:60446
TITLE: **Imidacloprid**-enhanced Reticulitermes flavipes (Isoptera: Rhinotermitidae) susceptibility to the entomopathogen Metarhizium anisopliae
AUTHOR(S): Ramakrishnan, Rathna; Suiter, Daniel R.; Nakatsu, Cindy H.; Humber, Richard A.; Bennett, Gary W.
CORPORATE SOURCE: Center for Urban & Industrial Pest Management, Department of Entomology, Purdue University, West Lafayette, IN, 47907, USA
SOURCE: Journal of Economic Entomology (1999), 92(5), 1125-1132
CODEN: JEENAI; ISSN: 0022-0493
PUBLISHER: Entomological Society of America
DOCUMENT TYPE: Journal
LANGUAGE: English
AB The effects of **imidacloprid** and the entomopathogen Metarhizium anisopliae (Metsch.) Sorokin on the eastern subterranean termite, Reticulitermes flavipes (Kollar), were evaluated in a 4 + 3 factorial experiment in both sterile and nonsterile loam soil. **Termites** were not susceptible to M. anisopliae when assays were conducted in nonsterile soil, but were highly susceptible in sterile soil. Termite mortality after 21 days of continuous exposure to 104 conidia per g soil was 0 and 41.6% in nonsterile and sterile soil, resp. **Termites** were significantly more susceptible to sterile soil containing 107 conidia per g than to the same soil containing 104 conidia per g.
In continuous exposure assays, **termites** were highly susceptible to **imidacloprid**-treated (5,10, and 20 ppm) nonsterile and sterile soil containing no exptl. introduced M. anisopliae. Exposure of **termites** to **imidacloprid** enhanced their susceptibility to introduced M. anisopliae in nonsterile and sterile soil. Native entomopathogens recovered from **termites** exposed to **imidacloprid**-treated, nonsterile soil (i.e., no introduced M.

anisopliae) included *Conidiobolus coronatus* (Constantin) Batko, *Cunninghamella echinulata* Thaxter, *Fusarium* spp., *Aspergillus* spp., and a naturally occurring strain of *M. anisopliae* variety *majus*.

REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 17 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:125767 CAPLUS

DOCUMENT NUMBER: 130:178773

TITLE: Composition for the control of wood-destroying insects, especially **termites**

INVENTOR(S): Anderson, John-phillip-evans; Keuken, Oliver

PATENT ASSIGNEE(S): Bayer A.-G., Germany

SOURCE: Eur. Pat. Appl., 21 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 896791	A2	19990217	EP 1998-114187	19980729
EP 896791	A3	20000112		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
DE 19734665	A1	19990218	DE 1997-19734665	19970811
TW 505500	B	20021011	TW 1998-87112592	19980731
US 6264968	B1	20010724	US 1998-128818	19980804
ZA 9807118	A	19990209	ZA 1998-7118	19980807
JP 11124302	A2	19990511	JP 1998-234861	19980807
AU 9879895	A1	19990218	AU 1998-79895	19980811
AU 768390	B2	20031211		
BR 9803138	A	19991221	BR 1998-3138	19980811

PRIORITY APPLN. INFO.: DE 1997-19734665 A 19970811

AB The title comps. (no examples) comprise an insecticide, preferably **imidacloprid**, incorporated into an organic natural and/or synthetic carrier. Optional ingredients are insect attractants and microbicides.

L3 ANSWER 18 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:54407 CAPLUS

DOCUMENT NUMBER: 130:206253

TITLE: Control of the termite *Heterotermes tenuis* (Hagen) using Termitrap baits impregnated with insecticides associated with the entomopathogenic fungus *Beauveria bassiana* (Bals.) Vuill.

AUTHOR(S): Almeida, Jose E. M.; Alves, Sergio B.; Moino, Alcides, Jr.; Lopes, E. Rogerio B.

CORPORATE SOURCE: Laboratorio de Controle Biologico, Centro Experimental, Instituto Biologico, Campinas, 13001-970, Brazil

SOURCE: Anais da Sociedade Entomologica do Brasil (1998), 27(4), 639-644

CODEN: ASENBI; ISSN: 0301-8059

PUBLISHER: Sociedade Entomologica do Brasil

DOCUMENT TYPE: Journal

LANGUAGE: Portuguese

AB The control of *H. tenuis* was evaluated using the bait/trap Termitrap impregnated with insecticides in low concns., associated to *B. bassiana* isolate 634 (from *Solenopsis invicta*), in sugarcane (*Saccharum*

officinatum). The treatments consisted of: **imidacloprid** 0,01%; **imidacloprid** 0,01% + *B. bassiana*; WG 0,003%; WG 0,003% + *B. bassiana*; *B. bassiana*; and untreated control. Each treatment was replicated five times. The insecticides were impregnated on baits by immersion in water, their concns. being calculated according to the weight of the bait, and the *B. bassiana* was applied as pure conidia (109 conidia/bait). The evaluations were made after 15, 30, 41, 63, 86 e 136 days, by assigning indexes to populations levels. All treatments significantly reduced termite populations when compared to the control. It took longer for *B. bassiana* alone to reduced *H. tenuis* population. The treatments with **imidacloprid** and WG were the most efficient in the control of **termites** in sugarcane. The baits/traps did not repel the **termites**.

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 19 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1996:411657 CAPLUS

TITLE: **Imidacloprid** - chemical synergist for microbial control agents of **termites**.

AUTHOR(S): Boucias, D. G.

CORPORATE SOURCE: Department Entomology & Nematology, University Florida, Gainesville, FL, 32611-0620, USA

SOURCE: Book of Abstracts, 212th ACS National Meeting, Orlando, FL, August 25-29 (1996), AGRO-019. American Chemical Society: Washington, D. C. CODEN: 63BFAF

DOCUMENT TYPE: Conference; Meeting Abstract

LANGUAGE: English

AB Our research has determined that the neurotoxin, **imidacloprid**, at sublethal concns., can significantly alter the behavioral patterns of insects. For example, the subterranean termite, *Reticulotermis flavipes* possesses social behaviors (grooming, tunnel construction) which serve as the primary line of defense against pathogenic and opportunistic microorganisms. These behaviors, in combination with the resident microflora, confer a high degree of disease resistance upon these social insects. Exposure to low dosages of **imidacloprid** produces a long term disruption of these social behaviors resulting in the onset of epizootics initiated by either resident or introduced microbes. Related studies on other nonsocial insects (cockroaches, weevils) have supported the results found with **termites**. At sublethal concns., **imidacloprid** acted as a behavioral modifying agent significantly increasing the host insects susceptibility to microbial control agents.

L3 ANSWER 20 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1995:648220 CAPLUS

DOCUMENT NUMBER: 123:27832

TITLE: Odorless insect repellents against **termites**

INVENTOR(S): Ueda, Masayoshi; Muto, Yutaka

PATENT ASSIGNEE(S): Japan Carlit Co Ltd, Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----

09886197

JP 07089803 A2 19950404 JP 1993-258961 19930924
PRIORITY APPLN. INFO.: JP 1993-258961 19930924
OTHER SOURCE(S): MARPAT 123:27832
GI



AB An odorless insect repellent contains a repellent, a solvent and surfactant, or preservative; the solvent being I (R1, R2 = H, C1-2 alkyl; R3 = C1-3 alkyl). The active repellent may be chlorpyrifos, phoxim, pyridaphenthion, allethrin, carbaril, **imidacloprid**, etc. For example, an odorless emulsion was prepared by combining dimethylpropylnaphthalene, chlorpyrifos, Sorpol-3006K and Sorpol-3008K.

L3 ANSWER 21 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1995:187187 CAPLUS

DOCUMENT NUMBER: 122:25815

TITLE: **Imidacloprid** - a new systemic insecticide.

AUTHOR(S): Elbert, A.; Becker, B.; Hartwig, J.; Erdelen, C.

CORPORATE SOURCE: Geschäftsbereich Pflanzenschutz
Entwicklung/Insektizide, Bayer AG, Leverkusen, 5090,
Germany

SOURCE: Pflanzenschutz-Nachrichten Bayer (German Edition)
(1991), 44(2), 113-36

CODEN: PNBAT; ISSN: 0340-1723

PUBLISHER: Bayer AG

DOCUMENT TYPE: Journal

LANGUAGE: German


AB The biol. profile of **Imidacloprid** (I) was defined on the basis of the results of exhaustive laboratory expts. and greenhouse trials. I is extremely effective against sucking insects, such as rice leafhoppers, aphids, thrips and mealybugs, and very effective against whitefly. It is also effective against some species of biting insects, such as paddy stem borers and Colorado beetle, but it has no effect on nematodes or spider mites. At comparatively high doses it kills adult insects and has ovicidal effects. I is a nicotinic acetylcholine receptor stimulator. Its mechanism of action differs from that of conventional insecticides. It therefore gives excellent control of all resistant populations investigated hitherto. I has a pos. temperature coefficient. After foliar application, it has a good residual action, it is highly photostable and it shows satisfactory resistance to rain. I is active after oral ingestion and by direct contact, but it is not active in the vapor phase. The LD95 after oral ingestion by *Myzus persicae* is .apprx.2 pg/aphid. After topical application it is .apprx.160 pg/aphid. It has not been possible to demonstrate recovery of injured aphids, or antifeeding effects. I has a faster action against aphids than oxydemeton-Me. After foliar application, I shows good translaminar and acropetal translocation, so it is also likely to provide effective control of pests with a furtive lifestyle, and protect the parts of the plant which regenerate after treatment. By virtue of its good contact action and powerful systemic action after uptake through the root system, I can be applied to soil and used as a seed dressing. It gives excellent control of pests such as onion maggots, *Diabrotica*, wire worms, **termites** and fire ants

which live in the soil, and of insects such as aphids which live above ground level. It has a good residual action after application to the soil and when it is used as a seed dressing. The compatibility of I with plants is good after use as a seed dressing, as a soil treatment and after foliar application. By virtue of its biol. properties, I is likely to have a wide range of uses for controlling economically important pests of rice, cotton, cereals, maize, sugar beet, potatoes, vegetables, citrus fruit, pome and stone fruit and other crops.

L3 ANSWER 22 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1993:54353 CAPLUS
 DOCUMENT NUMBER: 118:54353
 TITLE: Imidazolidine derivatives and related compounds as industrial insecticides and wood preservatives
 INVENTOR(S): Tsuboi, Shinichi; Sone, Shinzaburo; Obinata, Toru; Exner, Otto; Schwamborn, Michael
 PATENT ASSIGNEE(S): Nihon Bayer Agrochem K. K., Japan
 SOURCE: Eur. Pat. Appl., 15 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 511541	A1	19921104	EP 1992-106384	19920414
EP 511541	B1	19960904		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
JP 05032505	A2	19930209	JP 1991-350751	19911212
JP 3162450	B2	20010425		
JP 2001031511	A2	20010206	JP 2000-233512	19911212
AU 9213908	A1	19921029	AU 1992-13908	19920330
AU 645672	B2	19940120		
AT 142077	E	19960915	AT 1992-106384	19920414
ES 2090400	T3	19961016	ES 1992-106384	19920414
BR 9201534	A	19921201	BR 1992-1534	19920427
US 6323224	B1	20011127	US 1995-543351	19951016
US 2001051643	A1	20011213	US 2001-886197	20010621
PRIORITY APPLN. INFO.:			JP 1991-125172	A 19910427
			JP 1991-350751	A 19911212
			US 1992-872279	B1 19920422
			US 1995-543351	A3 19951016

OTHER SOURCE(S): MARPAT 118:54353
 GI



AB The imidazolidine derivs. and related compds. I (X = NH, S; Y = CH, N; Z = 2-chloro-5-pyridyl, 2-chloro-5-thiazolyl; R = H, Me; n = 0, 1) are industrial insecticides and wood preservatives. Wood impregnated with 0.32 ppm imidacloprid was lethal to termites

09886197

(Coptotermes formosanus) for ≥ 3 wk.

=> d 14 1-25 ibib hitstr abs

L4 ANSWER 1 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 2003:143339 CAPLUS
DOCUMENT NUMBER: 138:189635
TITLE: UV-protecting aqueous **wood** preservatives
with low hiding power
INVENTOR(S): Fukuoka, Naohiko; Onishi, Isamu
PATENT ASSIGNEE(S): Chemipro Kasei Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003055119	A2	20030226	JP 2001-250908	20010821

PRIORITY APPLN. INFO.: JP 2001-250908 20010821

AB Title preservatives contain UV absorbers, insecticides, **wood** preservatives, and binders. Thus, an aqueous **wood** preservative containing 2-(2'-hydroxy-3',5'-di-tert-amylphenyl)benzotriazole, ethofenprox, 3-iodo-2-propynyl butylcarbamate, SN Defoamer 318 (silicone emulsion) and Rikabond ES 1 (acrylic copolymer emulsion) was applied on lumber and left for 18 mo to show no fungi formation, no discoloration on the coated surface, and yellowing ΔE 7.7.

L4 ANSWER 2 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 2002:964915 CAPLUS
DOCUMENT NUMBER: 138:12164
TITLE: Barrier preventing **wood** pest access to wooden structures
INVENTOR(S): Van Voris, Peter; Cataldo, Dominic A.; Burton, Frederick G.; Leong, Henry; Stonich, Derek; Lin, K. C.; McClellan, William D.; Bowdle, Kurt W.
PATENT ASSIGNEE(S): USA
SOURCE: U.S. Pat. Appl. Publ., 33 pp., Cont.-in-part of U.S. Ser. No. 353,494.
CODEN: USXXCO
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 3
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2002192259	A1	20021219	US 2001-5804	20011203
US 5985304	A	19991116	US 1998-30690	19980225

PRIORITY APPLN. INFO.: US 1998-30690 A1 19980225
US 1999-353494 A2 19990713
US 2000-251112P P 20001203
US 2000-251141P P 20001204

AB A multi-layer **wood** pest barrier having a prolonged lifetime is given. The lifetime can be as long as the life of a building or structure to be protected. The lifetime protection is achieved by binding at least one pesticide within a continuous or discontinuous polymer matrix layer

thereby reducing release of the pesticide from the matrix. The release rate of the pesticide from the matrix can be controlled by the use of a carrier such as carbon black. The release of the pesticide from the barrier can be further controlled by inclusion of addnl. layers which can make the barrier nonreleasing.

L4 ANSWER 3 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:868829 CAPLUS
DOCUMENT NUMBER: 137:354591
TITLE: Carrier composition of fungicides and insecticides for protective treatment of wood
INVENTOR(S): Rodriguez Ramos, Rafael
PATENT ASSIGNEE(S): Spain
SOURCE: PCT Int. Appl., 28 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: Spanish
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002090068	A1	20021114	WO 2001-ES175	20010507
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
SI 21088	C	20030630	SI 2001-20039	20010507
BR 2001012150	A	20030701	BR 2001-12150	20010507
EP 1391278	A1	20040225	EP 2001-929660	20010507
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
NO 2002006272	A	20030219	NO 2002-6272	20021230
BG 107440	A	20030930	BG 2003-107440	20030106
HR 2003000076	A1	20030430	HR 2003-76	20030206
US 2003162781	A1	20030828	US 2003-371740	20030221
US 6673836	B2	20040106		

PRIORITY APPLN. INFO.: WO 2001-ES175 A 20010507

AB The carrier comprises toluene (40-70%), xylene (6-40%), benzophenone (3-18%), butylglycol (2-9%), cetyl acetate (1-7%) and methanol (0.3-4%) and insecticides and fungicides. The insecticides and fungicides are selected from Chlorpyrifos, Fipronil, Silafluofen, Acetamiprid, Etofenprox, tri-Pr isocyanate, Fenobucarb, Hexaflumuron, Fenitrothion, Esfenvalerate, Imidacloprid, Diflubenzuron, λ -cyhalothrin, Propioconazole, and mixts.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 4 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:695680 CAPLUS
DOCUMENT NUMBER: 137:228094
TITLE: Termiticidal baits for eliminating termite colonies
INVENTOR(S): Brode, Philip Frederick, III; Garrett, Garry Steven; Laughlin, Leo Timothy; Matthews, Randall Stryker; Barker, Dale Edwin; Kinne, Daniel James; Miller,

Christopher Miles; Probst, Timothy Robert; McKibben,
 Gary Eugene
 PATENT ASSIGNEE(S): The Procter & Gamble Company, USA
 SOURCE: PCT Int. Appl., 61 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002069704	A2	20020912	WO 2002-US6200	20020301
WO 2002069704	A3	20021114		
WO 2002069704	C1	20031231		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 2002172658	A1	20021121	US 2001-799184	20010305
US 6716421	B2	20040406		
US 2003017187	A1	20030123	US 2002-172855	20020617
US 2003124166	A1	20030703	US 2002-173527	20020617
US 2003124164	A1	20030703	US 2002-268356	20021010
WO 2003105580	A1	20031224	WO 2003-US17713	20030605
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
WO 2003106395	A1	20031224	WO 2003-US17714	20030605
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
WO 2004032625	A2	20040422	WO 2003-US32092	20031007
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,			

09886197

CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG


PRIORITY APPLN. INFO.:

US 2001-799184 A 20010305
US 2002-172855 A 20020617
US 2002-173527 A 20020617
US 2002-268356 A 20021010

OTHER SOURCE(S):

MARPAT 137:228094

GI



I

AB This invention relates to devices, kits, and methods for eliminating termite colonies. The kits, devices, and methods employ a termiticidal bait matrix contain (a) a termiticide (I, X = nil, a hydrocarbon group, O or NR₈, R₉ where R₈ and R₉ are H or a hydrocarbon group; X₁ = CH, a carbon atom or a heteroatom; R, R₁, R₂, R₃ = H or OH and if R₄ and R₅ are O and R₆ and R₇ are H then R, R₁, R₂ and R₃ may be C₁-6; R₄ and R₅ are H, O or N; R₉ and R₁₀ are nil, C₁-6, and amides) selected such that the termiticide causes death to about 50 to about 100% of termites within about 24 to about 84 days after the termites begin to ingest the termiticide or the bait matrix comprising the termiticide, (b) a cellulose containing material, and (c) water. The termiticidal bait matrix can be used in a bait station installed in the ground. The kits are suitable to be used by consumers in their homes.

L4 ANSWER 5 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:547214 CAPLUS

DOCUMENT NUMBER: 137:105178

TITLE: Termiticides containing 2-pyridinethiol-1-oxide salts and **wood** and polymers containing the termiticides

INVENTOR(S): Nishimoto, Koichi; Sato, Toshio; Suga, Mamoru

PATENT ASSIGNEE(S): Yoshitomi Fine Chemicals Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002205906	A2	20020723	JP 2001-337124	20010926

PRIORITY APPLN. INFO.: JP 2000-381082 A 20001108

AB The termiticides, which are effective on termites, bark beetles, etc., and environmentally safe, contain (a) ≥ 1 selected from Cu, Zn, and Na salts of 2-pyridine-1-oxide and optionally (b) ≥ 1 selected from pyrethroids, nicotinoids, organophosphorus compds., isocyanuric acid compds., carbamates, acetamiprid, and inorg. boric acid compds. **Wood** and polymers containing the termiticides are also claimed. A

wood block was coated with DMSO solution containing Cu pyrithione and **imidacloprid** and dried at room temperature for ≥ 20 days. The **wood** block.

L4 ANSWER 6 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:429963 CAPLUS
DOCUMENT NUMBER: 137:29419
TITLE: The use of Confidor S in the float, a new tobacco seedlings production system in the South of Brazil
AUTHOR(S): Leal, R. S.
CORPORATE SOURCE: Bayer S.A. Desenvolvimento Tecnico de Produtos, Sao Paulo-SP, 04779-900, Brazil
SOURCE: Pflanzenschutz-Nachrichten Bayer (German Edition) (2001), 54(3), 337-352
CODEN: PNBAT; ISSN: 0340-1723
PUBLISHER: Bayer AG
DOCUMENT TYPE: Journal
LANGUAGE: English

AB A float system for tobacco seedlings was introduced in the southern region of Brazil as an alternative to the Me bromide based fumigants used on tobacco seedbeds. Seedlings are cultivated on Styrofoam trays, which are filled with a special substrate on cellulose basis. After that, the trays are placed in a pool of water with a black plastic film and bricks or **wood** outlining the whole system. The following advantages were achieved: production of healthier and protected seedlings for transplantation, more uniform and productive crops, more comfortable work conditions, no seedbed sterilization with Me bromide and frequent irrigation, no controlling of mollusks in the seedbeds, seedling transplantation is less dependent on rain levels. To adopt plant protection to the new system, the insecticide mixture Confidor S 51 WP (500 g/kg of **imidacloprid** + 10 g/kg of cyfluthrin) was developed. The product is applied by watering the tobacco seedlings about 24 h before the definitive transplanting to the crops. The same excellent level of efficacy and residual effect in the control of pests was achieved with Confidor S compared to Confidor 70 WG. The addition of cyfluthrin broadened the spectrum of efficacy and controls *Agrotis ypsilon*. The addition of Confidor S to the float system to tobacco crops resulted in a series of benefits in the management of pests: protection since the initial stages of the cultivation against pests which are difficult to control, reduction in the number of sprays of the transplanted crops, less interference in the environment due to the reduction of the treated area, long-lasting protection, economical use of manpower, and less risk to the farmer.

REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 7 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:118039 CAPLUS
DOCUMENT NUMBER: 136:130232
TITLE: Preparation of **imidacloprid** microemulsion
INVENTOR(S): Zhou, Benxin
PATENT ASSIGNEE(S): Nuopuxin Agrochemistry Co., Ltd., Shenzhen, Peop. Rep. China
SOURCE: Faming Zhuanli Shenqing Gongkai Shuomingshu, 8 pp.
CODEN: CNXXEV
DOCUMENT TYPE: Patent
LANGUAGE: Chinese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
CN 1299594	A	20010620	CN 2001-100514	20010105

PRIORITY APPLN. INFO.: CN 2001-100514 20010105

AB The title microemulsion comprises **imidacloprid** 1-50, emulsifier 5-30, solubilizer 5-30, synergist 5-10, stabilizing agent 5-10, and water 20-80%. The solutizer is selected from one, two or three of benzyl alc., ethanol, isopropanol, n-butanol, n-pentanol, acetone, cyclohexanone and dimethylformamide; the emulsifier from two or three of emulsifier No 201, 500#, 602, 2201, 700#, Tween-80 and Tx-10; the synergist from one of octachlorodipropyl ether, azone or piperonyl butoxide; and the stabilizing agent from ethanediol, polyethylene glycol, urea or glycerin. The insecticide is prepared by mixing raw material and homogenizing.

L4 ANSWER 8 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:767469 CAPLUS

DOCUMENT NUMBER: 135:299970

TITLE: Insecticides containing salicylate esters for wood preservation

INVENTOR(S): Sato, Toshio; Nakamura, Norihiko; Goto, Shinji

PATENT ASSIGNEE(S): Yoshitomi Fine Chemical K. K., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 26 pp.
CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2001294506	A2	20011023	JP 2000-112664	20000413

PRIORITY APPLN. INFO.: JP 2000-112664 20000413

OTHER SOURCE(S): MARPAT 135:299970

AB The insecticides, which are especially useful for controlling termite and not toxic to humans, livestock, or environment, contain 2-OHC6H4CO2W1R1 [R1 = (un)substituted Ph, C2-12 (hydroxy)alkyl, C2-12 (hydroxy)alkenyl, C2-12 (hydroxy)alkynyl, W1 = bond, C1-6 alkylene, C2-6 alkenylene, C2-6 alkynylene]. The salicylates also serve as enhancers for com. available insecticides, showing synergistic effect. Thus, quartz sand treated with Ph salicylate showed 100% termiticidal activity.

L4 ANSWER 9 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:720924 CAPLUS

DOCUMENT NUMBER: 135:340463

TITLE: Chemical prevention of colony foundation by *Cryptotermes brevis* (Isoptera: Kalotermitidae) in attic modules

AUTHOR(S): Scheffrahn, Rudolf H.; Busey, Philip; Edwards, Jeffrey K.; Krecek, Jan; Maharajh, Boudanath; Su, Nan-Yao

CORPORATE SOURCE: Ft. Lauderdale Research and Education Center, University of Florida, Fort Lauderdale, FL, 33314, USA

SOURCE: Journal of Economic Entomology (2001), 94(4), 915-919
CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Disodium octaborate tetrahydrate (DOT) dust, DOT aqueous solution, **imidacloprid** dust, and amorphous silica gel dust with synergized 1% pyrethrins were applied on wood surfaces to simulated attic modules. Modules (30 by 30 cm) with and without fiberglass insulation

were exposed to dispersal flights of *Cryptotermes brevis* (Walker) in May and June of 1998 and 1999. Six months after flights, modules were disassembled and inspected for nuptial chamber location and contents. During both years, air and water control treatments contained 22.2 ± 9.94 (mean \pm SD) nuptial chambers, 7.5 ± 5.7 live imagos, and 2.0 ± 1.4 chambers with brood. This survivorship indicated that the attic modules performed well as a colonizing platform for *C. brevis*. *C. brevis* dealates preferred constructing nuptial chambers in the crevices at the bases or tops of the modules instead of internal crevices. Modules treated in 1998 and 1999 with DOT or silica dusts contained no live termites, whereas zero of five modules treated with **imidacloprid** dust in 1998 and two of 20 modules treated with **imidacloprid** dust in 1999 contained single live incipient colonies. In 1998, 15% DOT solution, applied as a postconstruction treatment, yielded significantly fewer chambers and live termites than controls, but was not as effective as dusts in preventing successful colonization. In 1999, the DOT solution, applied as a construction-phase treatment, was equally as effective in preventing colonization as the dust treatments during that year. Results indicate that dust formulations of DOT, silica gel, and **imidacloprid** can be used to prevent drywood termite colonization in existing building voids and attics. Where the entire **wood** framing is exposed to treatment, such as during building construction, aqueous DOT solution can be equally effective as dusts in preventing colonization by *C. brevis*.

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 10 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:161372 CAPLUS

DOCUMENT NUMBER: 134:189458

TITLE: Fast-drying preservative composition for **wood** and leather

INVENTOR(S): Narayanan, Kolazi S.; Jon, Domingo I.; Prettypaul, Donald

PATENT ASSIGNEE(S): ISP Investments Inc., USA

SOURCE: U.S., 3 pp.

CODEN: USXXAM

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6197098	B1	20010306	US 1999-464758	19991216
WO 2001043547	A1	20010621	WO 2000-US33425	20001208
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 2001022571	A5	20010625	AU 2001-22571	20001208
PRIORITY APPLN. INFO.:			US 1999-464758	A 19991216
			WO 2000-US33425	W 20001208

AB The invention relates to a fast-drying preservative composition having enhanced penetration for the treatment and preservation of **wood**, leather

09886197

and similar natural products, which comprises 10-50 weight % of a concentrate comprising a petroleum distillate boiling >40 and the balance a C2-4 aliphatic alc. containing 0-85 weight % of mineral spirit as a diluent to provide a sprayable composition The petroleum distillate concentrate comprises: (a) 0.5-7 weight % nitrogen- or sulfur-containing biocide and (b) 20-55 weight % solvent, consisting of: (i) butyrolactone containing 0-85 weight % N-methylpyrrolidone and/or 0-85 weight % C2-4 aliphatic alc. or (ii) N-methylpyrrolidone containing 0-85 weight % C2-4 alc. The biocide is **imidacloprid**, a guanidine, nicotine, a salicylate, etc. Pyroligneous acid can be optionally added as a stabilizer.

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 11 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:1168 CAPLUS
DOCUMENT NUMBER: 134:41726
TITLE: Controlled-release pesticide and fertilizer briquettes
INVENTOR(S): Moore, William Percy, Jr.
PATENT ASSIGNEE(S): Lesco, Inc., USA
SOURCE: Eur. Pat. Appl., 9 pp.
CODEN: EPXXDW
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1063215	A2	20001227	EP 2000-303118	20000413
EP 1063215	A3	20020925		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
US 6225258	B1	20010501	US 1999-344083	19990625
SE 2000001520	A	20001226	SE 2000-1520	20000427
FI 2000001363	A	20001226	FI 2000-1363	20000607
NO 2000003322	A	20001227	NO 2000-3322	20000623
JP 2001048705	A2	20010220	JP 2000-189238	20000623

PRIORITY APPLN. INFO.: US 1999-344083 A 19990625

AB An attrition- and shatter-resistant plant nutrient/pesticide briquette composition which slowly releases the nutrients and of biol. active materials over long periods of time, comprises slow-release plant nutrient particles, pesticide sorption particles, liquid systemic pesticide sorbed on the pesticide sorption particles to reduce pesticide leachability, and an adhesive coating the slow-release plant nutrient and pesticide sorption particles. The composition is formed into briquettes by pressing into dies at elevated pressures and temps. A six-step method is provided for the preparation of the slow-releasing briquettes from slow release fertilizers, such as magnesium ammonium phosphate; pesticide sorption particles, such as activated carbon; liquid systemic pesticides emulsions, such as imidachloprid; and adhesives, such as a vinylidine chloride, 2-ethylhexyl acrylate and acrylic acid resin emulsion.

L4 ANSWER 12 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:470450 CAPLUS
DOCUMENT NUMBER: 133:90469
TITLE: Adhesive composition containing insecticides,

preservatives, termite repellents and bactericides for lignocellulosic material and its complex

INVENTOR(S): Jaesch, Thomas; Fushiki, Kiyoyuki; Saito, Takanobu; Katsusawa, Yoshinaga

PATENT ASSIGNEE(S): Bayer A.-G., Germany; Ohshika Shinko K. K.; Chemiholz K. K.

SOURCE: Jpn. Kokai Tokkyo Koho, 12 pp.
CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000192001	A2	20000711	JP 1998-376942	19981228
KR 2000048138	A	20000725	KR 1999-57526	19991214
EP 1018413	A1	20000712	EP 1999-124843	19991215
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
AU 9965409	A1	20010628	AU 1999-65409	19991222
NZ 502074	A	20020301	NZ 1999-502074	19991223
NO 9906479	A	20000629	NO 1999-6479	19991227
US 2001027217	A1	20011004	US 1999-472589	19991227
BR 9907435	A	20010320	BR 1999-7435	19991228

PRIORITY APPLN. INFO.:

JP 1998-376942 A 19981228

AB The composition, for preparation of wood products (e.g., plywood), comprises an adhesive, an organic phenolic composition, an insecticide, a preservative, a termite repellent and a bactericide. Thus, a composition was made from Oshika Resin PWP 60 containing a solution of imidacloprid 3, IPBC 20 and 2-phenylphenol 15, and a solvent 62%.

L4 ANSWER 13 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:467897 CAPLUS

DOCUMENT NUMBER: 133:85609

TITLE: Termiticidal baits comprising nonhygroscopic agents in hygroscopic containers

INVENTOR(S): Minakawa, Fumiyasu; Uchida, Yuki

PATENT ASSIGNEE(S): Yuko Chemical Industries Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000189031	A2	20000711	JP 1998-369335	19981225

PRIORITY APPLN. INFO.:

JP 1998-369335 19981225

AB A nonhygroscopic agent for controlling termites (e.g. diflubenzuron) is housed in a container which is made of an edible hygroscopic material (cellulosic cloth, polyvinyl alc. film). Thus, sulfluramid 0.001 and pine oil 1% (attractant) were dissolved in polyethylene glycol, and the solution was made to soaked into wood flour at a 25/100 weight ratio. The agent was heat sealed in an envelope (15 + 7 cm) made of nonwoven fabric of cellulose fibers with 1% by weight added pine oil to obtain a bait with satisfactory attractiveness to Reticulitermes.

L4 ANSWER 14 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

09886197

ACCESSION NUMBER: 2000:424169 CAPLUS
DOCUMENT NUMBER: 133:39440
TITLE: Efficacy of **imidacloprid** for cockroach control in a Gel Bait formulation
AUTHOR(S): Pospischil, R.; Schneider, U.; Bocker, T.; Junkersdorf, J.; Nentwig, G.; Smith, G.; Sonneck, R.
CORPORATE SOURCE: Geschäftsbereich Tiergesundheit, Landwirtschaftszentrum Monheim, Bayer AG, Leverkusen, D-51368, Germany
SOURCE: Pflanzenschutz-Nachrichten Bayer (German Edition) (1999), 52(3), 386-400
CODEN: PNBAT; ISSN: 0340-1723
PUBLISHER: Bayer AG
DOCUMENT TYPE: Journal
LANGUAGE: German

AB The active substance **imidacloprid** is the first of the chloronicotinyl class of compds. to be used in a gel bait formulation for cockroach control. Its high efficacy as an edible poison, lack of any contact activity against cockroaches, and lack of secondary effects via the feces and dead insects allow **imidacloprid** to be formulated as a gel bait that meets the high demands of an effective and safe cockroach control strategy. Extensive laboratory and field trials and initial market feedback have demonstrated the high efficacy of the **imidacloprid** cockroach gel against all economically important cockroach species. In preliminary laboratory tests, **imidacloprid** cockroach gel was also found to be active against other pests such as wood lice, house crickets, and ants. The **imidacloprid** gel still showed outstanding activity even 27 mo after deployment of the gel pellets under various conditions. No difference vs. freshly laid bait was observed

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 15 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:339593 CAPLUS
DOCUMENT NUMBER: 132:335994
TITLE: Wood-penetrable compositions for preservatives and termite-repellent chemicals
INVENTOR(S): Oda, Kunitaka; Nushida, Masanori; Ishida, Daisaku
PATENT ASSIGNEE(S): Fumakilla Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 7 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000141317	A2	20000523	JP 1998-314618	19981105
PRIORITY APPLN. INFO.:			JP 1998-314618	19981105

AB The compns. contain glycol ethers slightly-soluble in water, isoparaffin-type hydrocarbons, and aliphatic esters. Thus, a composition comprising hydrocarbon (IP 2028) 65, ethylene glycol ethylhexyl ether 25, and isooctanoic acid ester 10% could be easily penetrated into **wood** plates.

L4 ANSWER 16 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:267592 CAPLUS
DOCUMENT NUMBER: 132:261676
TITLE: Insecticidal fumigant containing **imidacloprid**

09886197

INVENTOR(S): Wang, Kaiyun; Jiang, Xingyin; Yi, Meiqin; Xue, Ming
PATENT ASSIGNEE(S): Shangdong Agricultural Univ., Peop. Rep. China
SOURCE: Faming Zhuanli Shenqing Gongkai Shuomingshu, 5 pp.
CODEN: CNXXEV
DOCUMENT TYPE: Patent
LANGUAGE: Chinese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
CN 1196179	A	19981021	CN 1997-105782	19970417
CN 1055820	B	20000830		

PRIORITY APPLN. INFO.: CN 1997-105782 19970417

AB The insecticidal fumigant comprises **imidacloprid**, dichlorvos, oxidant, fuel, and fire retardant. The ratio of **imidacloprid** : dichlorvos is 1:10-50. The oxidant is selected from NH₄NO₃ and KNO₃; the fuel from **wood** meal; and the fire retardant from saponite and clay.

L4 ANSWER 17 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:480946 CAPLUS
DOCUMENT NUMBER: 131:140842
TITLE: Insecticides and preservatives for lumber
INVENTOR(S): Ueno, Takahide; Yonetani, Koreyasu
PATENT ASSIGNEE(S): Yuko Chemical Industries Co., Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 14 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
JP 11207706	A2	19990803	JP 1998-16808	19980129

PRIORITY APPLN. INFO.: JP 1998-16808 19980129

AB A preservative, propiconazole, in combination with ≥ 1 insecticide selected from the group consisting of tralomethrin, bifenthrin, permethrin, **imidacloprid**, fenobucarb, fipronil, and pyriproxyfen with the ratio of insecticide/preservative being 1.0-15.0, is used for preserving lumber. The concentrate of the mixture in water contains ≥ 40 fold effective concentration of the mixture, and the preparation is diluted with water prior to application to lumber. The mixture is stable for a long period.

L4 ANSWER 18 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:125767 CAPLUS
DOCUMENT NUMBER: 130:178773
TITLE: Composition for the control of **wood**
-destroying insects, especially termites
INVENTOR(S): Anderson, John-phillip-evans; Keuken, Oliver
PATENT ASSIGNEE(S): Bayer A.-G., Germany
SOURCE: Eur. Pat. Appl., 21 pp.
CODEN: EPXXDW
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 896791	A2	19990217	EP 1998-114187	19980729
EP 896791	A3	20000112		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
DE 19734665	A1	19990218	DE 1997-19734665	19970811
TW 505500	B	20021011	TW 1998-87112592	19980731
US 6264968	B1	20010724	US 1998-128818	19980804
ZA 9807118	A	19990209	ZA 1998-7118	19980807
JP 11124302	A2	19990511	JP 1998-234861	19980807
AU 9879895	A1	19990218	AU 1998-79895	19980811
AU 768390	B2	20031211		
BR 9803138	A	19991221	BR 1998-3138	19980811

PRIORITY APPLN. INFO.: DE 1997-19734665 A 19970811
 AB The title compns. (no examples) comprise an insecticide, preferably **imidacloprid**, incorporated into an organic natural and/or synthetic carrier. Optional ingredients are insect attractants and microbicides.

L4 ANSWER 19 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1998:293323 CAPLUS

DOCUMENT NUMBER: 128:318352

TITLE: Wood preservatives for incorporation into binders, for plywood and chipboard manufacture
 INVENTOR(S): Buschhaus, Hans-Ulrich; Exner, Otto; Fushiki, Seiko
 PATENT ASSIGNEE(S): Bayer A.-G., Germany; Kemiholz Co. Ltd.; Buschhaus, Hans-Ulrich; Exner, Otto; Fushiki, Seiko

SOURCE: PCT Int. Appl., 21 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9818328	A1	19980507	WO 1997-EP5776	19971020
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
DE 19644008	A1	19980507	DE 1996-19644008	19961031
AU 9850511	A1	19980522	AU 1998-50511	19971020
AU 736300	B2	20010726		
EP 935416	A1	19990818	EP 1997-913163	19971020
R: AT, BE, CH, DE, DK, ES, FR, GB, IT, LI, NL, SE, FI				
CN 1235515	A	19991117	CN 1997-199283	19971020
NZ 335434	A	20001124	NZ 1997-335434	19971020
JP 2001508406	T2	20010626	JP 1998-519997	19971020
KR 2000049019	A	20000725	KR 1999-703085	19990409

PRIORITY APPLN. INFO.: DE 1996-19644008 A 19961031
 WO 1997-EP5776 W 19971020

OTHER SOURCE(S): MARPAT 128:318352

AB The invention relates to **wood** preservatives compatible with binders or adhesives, which can be employed for the manufacture of plywood, chipboard and timber materials. The **wood** preservatives are

09886197

RNACZ:XE [R = H, (un)substituted acyl, alkyl, aryl, aralkyl, heteroaryl and heteroarylalkyl; A = H, acyl, alkyl, aryl, or a bifunctional group which is linked to the radical Z; E = electron-withdrawal radical; X = CH or N; CH is optionally linked to Z instead of H; Z = alkyl, OR, SR, or a bifunctional group which is linked to A or X]. **Imidacloprid** is particularly preferred.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 20 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1998:183889 CAPLUS

DOCUMENT NUMBER: 128:240732

TITLE: Synergistic insecticidal and wood preservative compositions

INVENTOR(S): Asai, Takehito; Okumura, Kenya; Shizawa, Toshiyasu

PATENT ASSIGNEE(S): Sankyo Co., Ltd., Japan

SOURCE: Eur. Pat. Appl., 11 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 829203	A1	19980318	EP 1997-307024	19970910
EP 829203	B1	20021218		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
CA 2214952	AA	19980311	CA 1997-2214952	19970909
AU 9736872	A1	19980319	AU 1997-36872	19970909
AU 728200	B2	20010104		
US 5935943	A	19990810	US 1997-926372	19970909
JP 11029419	A2	19990202	JP 1997-244944	19970910
JP 3172698	B2	20010604		
ES 2187730	T3	20030616	ES 1997-307024	19970910
HK 1006215	A1	20030509	HK 1998-105511	19980617
US 6022881	A	20000208	US 1999-281712	19990330

PRIORITY APPLN. INFO.: JP 1996-240118 A 19960911

JP 1997-126988 A 19970516

US 1997-926372 A3 19970909

AB The presence of isobornyl thiocynoethyl ether exerts a synergistic effect on the insecticidal activity against harmful wood-eating insects of certain known insecticides, such as **imidacloprid**, phenylpyrazole derivs., pyrethroids and non-ester pyrethroid insecticides.

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 21 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1998:35959 CAPLUS

DOCUMENT NUMBER: 128:111913

TITLE: Wood preservatives and their use at ambient pressure

INVENTOR(S): Igarashi, Rei

PATENT ASSIGNEE(S): Takeda Chemical Industries, Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

09886197

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 10007502	A2	19980113	JP 1996-158363	19960619

PRIORITY APPLN. INFO.: JP 1996-158363 19960619

AB **Wood** preservatives contain water-immiscible fungicides, water-immiscible insecticides, water-immiscible liquid hydrocarbons with b.p. $\geq 220^\circ$ and flash point $\geq 100^\circ$, surfactants, and optional water. The preservatives are diluted with water and coated to **wood** at ambient pressure. A **wood** preservative emulsion was formulated containing IPBC, cyfluthrin, KMC 113 (dipropylnaphthalene) (sic), Newkalgen CP 80 (polyoxyalkylene styrylphenyl ether), and water.

L4 ANSWER 22 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1997:440126 CAPLUS

DOCUMENT NUMBER: 127:46479

TITLE: Water-based, solvent- and emulsifier-free microbicidal compositions.

INVENTOR(S): Buschhaus, Hans-Ulrich; Exner, Otto; Kugler, Martin; Nagano, Yukihiro

PATENT ASSIGNEE(S): Bayer A.-G., Germany

SOURCE: Ger. Offen., 12 pp.
CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19543477	A1	19970528	DE 1995-19543477	19951122
CA 2238033	AA	19970529	CA 1996-2238033	19961111
WO 9718713	A1	19970529	WO 1996-EP4919	19961111

W: AU, BB, BG, BR, BY, CA, CN, CZ, HU, JP, KR, KZ, LK, MX, NO, NZ, PL, RO, RU, SK, TR, UA, US

RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG

AU 9675694	A1	19970611	AU 1996-75694	19961111
EP 863709	A1	19980916	EP 1996-938169	19961111

R: AT, BE, CH, DE, DK, ES, FR, GB, IT, LI, NL

JP 2000500475	T2	20000118	JP 1997-519342	19961111
BR 9611746	A	20000328	BR 1996-11746	19961111

PRIORITY APPLN. INFO.: DE 1995-19543477 A 19951122
WO 1996-EP4919 W 19961111

OTHER SOURCE(S): MARPAT 127:46479

AB The title compns. comprise azole fungicide(s) (triadimefon, triadimenol, tebuconazole, hexaconazole, etc.), nitromethylene or related insecticide(s) and quaternary ammonium fungicide(s). The compns. are useful for the preservation of leather, **wood** and tech. materials.

L4 ANSWER 23 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1995:753647 CAPLUS

DOCUMENT NUMBER: 123:135913

TITLE: Synergistic combinations of ammonium salts for control of materials-destroying insects.

INVENTOR(S): Sagenmueller, Alfons; Schubert, Hans-Herbert; Uzawa, Shigeru; Saito, Kenichi

PATENT ASSIGNEE(S): Hoechst Schering AgrEvo GmbH, Germany

09886197

SOURCE: Eur. Pat. Appl., 11 pp.
CODEN: EPXXDW
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 664081	A2	19950726	EP 1995-100429	19950113
EP 664081	A3	19961002		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, NL, SE				
DE 4401542	A1	19950727	DE 1994-4401542	19940120
AU 9510286	A1	19950727	AU 1995-10286	19950118
CN 1111477	A	19951115	CN 1995-100978	19950118
US 5792755	A	19980811	US 1995-374309	19950118
CA 2140572	AA	19950721	CA 1995-2140572	19950119
ZA 9500425	A	19950926	ZA 1995-425	19950119
JP 07277906	A2	19951024	JP 1995-6607	19950119
US 5703132	A	19971230	US 1996-752582	19961121
PRIORITY APPLN. INFO.:			DE 1994-4401542	A 19940120
			US 1995-374309	A3 19950118

OTHER SOURCE(S): MARPAT 123:135913

AB The title compns. comprise a quaternary ammonium salt (Markush given) and a known insecticide, such as silafluofen, MTI-732, **imidacloprid**, ethofenprox, PP 682, etc. Thus, a mixture of Sanisol B-50 and silafluofen synergistically controlled *Reticulitermes speratus*.

L4 ANSWER 24 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1995:682581 CAPLUS

DOCUMENT NUMBER: 123:59251

TITLE: Wood preservative, concentrates and preservation of wood

INVENTOR(S): Heuer, Lutz; Kugler, Martin; Buschhaus, Hans-Ulrich; Schrage, Heinrich; Kunisch, Franz

PATENT ASSIGNEE(S): Bayer A.-G., Germany

SOURCE: PCT Int. Appl., 28 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9500303	A1	19950105	WO 1994-EP1868	19940608
W: AU, BB, BG, BR, BY, CA, CN, CZ, FI, HU, JP, KR, KZ, LK, NO, NZ, PL, RO, RU, SK, UA, US				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
DE 4320495	A1	19941222	DE 1993-4320495	19930621
DE 4406819	A1	19950907	DE 1994-4406819	19940302
AU 9471231	A1	19950117	AU 1994-71231	19940608
AU 689480	B2	19980402		
EP 705160	A1	19960410	EP 1994-920437	19940608
R: AT, BE, CH, DE, DK, ES, FR, GB, IT, LI, NL, PT, SE				
BR 9407120	A	19960903	BR 1994-7120	19940608
JP 08509437	T2	19961008	JP 1994-502383	19940608
NO 9505107	A	19951215	NO 1995-5107	19951215
US 5972971	A	19991026	US 1995-564249	19951215

09886197

FI 9506113 A 19951219 FI 1995-6113 19951219
 PRIORITY APPLN. INFO.: DE 1993-4320495 A 19930621
 DE 1994-4406819 A 19940302
 WO 1994-EP1868 W 19940608


AB Title combination contains α -butyl- α -(2,4-dichlorophenyl)-1H-1,2,4-triazol-1-ethanol (hexaconazole), and/or 5-[(4-chlorophenyl)methyl]-2,2-dimethyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol (metconazole) fungicides, and ≥ 1 supplementary synergistic insecticide. The addition of the synergistic insecticide to the azole fungicide does not impair the activity of the fungicide, the combinations have good stability, long term activity, a broad activity spectrum, and good penetrability in wood.

L4 ANSWER 25 OF 25 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1993:54353 CAPLUS
 DOCUMENT NUMBER: 118:54353
 TITLE: Imidozolidine derivatives and related compounds as industrial insecticides and wood preservatives
 INVENTOR(S): Tsuboi, Shinichi; Sone, Shinzaburo; Obinata, Toru; Exner, Otto; Schwamborn, Michael
 PATENT ASSIGNEE(S): Nihon Bayer Agrochem K. K., Japan
 SOURCE: Eur. Pat. Appl., 15 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 511541	A1	19921104	EP 1992-106384	19920414
EP 511541	B1	19960904		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
JP 05032505	A2	19930209	JP 1991-350751	19911212
JP 3162450	B2	20010425		
JP 2001031511	A2	20010206	JP 2000-233512	19911212
AU 9213908	A1	19921029	AU 1992-13908	19920330
AU 645672	B2	19940120		
AT 142077	E	19960915	AT 1992-106384	19920414
ES 2090400	T3	19961016	ES 1992-106384	19920414
BR 9201534	A	19921201	BR 1992-1534	19920427
US 6323224	B1	20011127	US 1995-543351	19951016
US 2001051643	A1	20011213	US 2001-886197	20010621
PRIORITY APPLN. INFO.:			JP 1991-125172	A 19910427
			JP 1991-350751	A 19911212
			US 1992-872279	B1 19920422
			US 1995-543351	A3 19951016

OTHER SOURCE(S): MARPAT 118:54353
 GI



09886197

AB The imidazolidine derivs. and related compds. I (X = NH, S; Y = CH, N; Z = 2-chloro-5-pyridyl, 2-chloro-5-thiazolyl; R = H, Me; n = 0, 1) are industrial insecticides and wood preservatives. Wood impregnated with 0.32 ppm imidacloprid was lethal to termites (Coptotermes formosanus) for ≥3 wk.

=>

=> logoff

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

LOGOFF? (Y)/N/HOLD:H

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION

FULL ESTIMATED COST

151.31	156.91
--------	--------

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION

CA SUBSCRIBER PRICE

-38.96	-38.96
--------	--------

SESSION WILL BE HELD FOR 60 MINUTES

STN INTERNATIONAL SESSION SUSPENDED AT 16:13:43 ON 21 JUL 2004

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssspta1202sxq

PASSWORD:

* * * * * RECONNECTED TO STN INTERNATIONAL * * * * *
SESSION RESUMED IN FILE 'CAPLUS' AT 16:15:44 ON 21 JUL 2004
FILE 'CAPLUS' ENTERED AT 16:15:44 ON 21 JUL 2004
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION

FULL ESTIMATED COST

151.31	156.91
--------	--------

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION

CA SUBSCRIBER PRICE

-38.96	-38.96
--------	--------

=> s termites

L6 2129 TERMITES

=> s insects

L7 29825 INSECTS

=> s l6 and l7

L8 274 L6 AND L7

=> s l8 and species

661358 SPECIES

L9 47 L8 AND SPECIES

=> d l9 1-5 ibib hitstr abs

8/4804

L9 ANSWER 1 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2004:542763 CAPLUS
 TITLE: Characterization of termite lipophorin and its involvement in hydrocarbon transport
 AUTHOR(S): Fan, Yongliang; Schal, Coby; Vargo, Edward L.; Bagneres, Anne-Genevieve
 CORPORATE SOURCE: Department of Entomology and W.M. Keck Center for Behavioral Biology, North Carolina State University, Box 7613, Raleigh, NC, 27695-7613, USA
 SOURCE: Journal of Insect Physiology (2004), 50(7), 609-620
 CODEN: JIPHAF; ISSN: 0022-1910
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The transport of lipids constitutes a vital function in **insects** and requires the plasma lipoprotein lipophorin. In all **insects** examined to date, cuticular hydrocarbons are also transported through the hemolymph by lipophorin, and in social **insects** they play important roles not only in water proofing the cuticle but also in nestmate recognition. High-d. lipophorin (HDLp), isolated from *Reticulitermes flavipes* plasma by KBr gradient ultracentrifugation, contains 66.2% protein and 33.8% lipids; hydrocarbons constitute its major neutral lipid (20.4% of total lipids). Anti-lipophorin serum was generated in rabbit and its specific association with lipophorin, and not with any other plasma proteins, was verified with Western blotting. Immunopptn. also confirmed that this antibody specifically recognizes lipophorin, because all hemolymph hydrocarbons of the **termites** *R. flavipes* and *R. lucifugus* and the cockroach *Supella longipalpa*, which associate only with lipophorin, were recovered in the immunopptd. protein. Cross-reactivity of the antiserum with lipophorin from related **species** was investigated by double immunodiffusion with 10 termite **species** in the genera *Reticulitermes*, *Coptotermes*, *Zootermopsis*, and *Kaloterme*s, and with five cockroach **species**. Involvement of lipophorin in hydrocarbon transport was shown by injecting HDLp antiserum into *Zootermopsis nevadensis* and then monitoring the de novo biosynthesis of hydrocarbons and their transport to the cuticular surface; the antiserum significantly disrupted hydrocarbon transport. ELISA revealed a gradual increase in the lipophorin titer in successively larger *R. flavipes* workers, and differences among castes in lipophorin titers were highest between nymphs and first instar larvae.

L9 ANSWER 2 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2004:169819 CAPLUS
 DOCUMENT NUMBER: 141:36378
 TITLE: Cuticular hydrocarbons and aggression in the termite *Macrotermes subhyalinus*
 AUTHOR(S): Kaib, Manfred; Jmhasly, Patrick; Wilfert, Lena; Durka, Walter; Franke, Stephan; Francke, Wittko; Leuthold, Reinhard H.; Brandl, Roland
 CORPORATE SOURCE: Department of Animal Physiology, University of Bayreuth, Bayreuth, D-95440, Germany
 SOURCE: Journal of Chemical Ecology (2004), 30(2), 365-385
 CODEN: JCECD8; ISSN: 0098-0331
 PUBLISHER: Kluwer Academic/Plenum Publishers
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Cuticular hydrocarbons are among the prime candidates for nestmate recognition in social **insects**. We analyzed the variation of cuticular hydrocarbons in the termite **species** *M. subhyalinus* in

West Africa (Comoe National Park) on a small spatial scale (<1 km). We found considerable variation in the composition of cuticular hydrocarbons among colonies, with four distinct chemical phenotypes. Different phenotypes occurred within each of the four habitats. The difference between these phenotypes is primarily due to unsatd. compds. A clear correlation between the difference of the hydrocarbon composition and the aggression between colonies was found. This correlation also holds in a multivariate anal. of genetic similarity (measured by AFLPs), morphometric distances (measured by Mahalanobis-distances), as well as geog. distances between colonies. In a more detailed anal. of the correlation between the composition of cuticular hydrocarbons and aggression, we found that no single compound is sufficient to explain variation in aggression between pairings of colonies. Thus, **termites** seem to use a bouquet of compds.

Multiple regression anal. suggested that many of these compds. are unsatd. hydrocarbons and, thus, may play a key role in colony recognition.

REFERENCE COUNT: 59 THERE ARE 59 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 3 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2004:116807 CAPLUS

DOCUMENT NUMBER: 140:284501

TITLE: The gut bacteria of **insects**: nonpathogenic interactions

AUTHOR(S): Dillon, R. J.; Dillon, V. M.

CORPORATE SOURCE: Department of Biology and Biochemistry, University of Bath, Bath, BA2 7AY, UK

SOURCE: Annual Review of Entomology (2004), 49, 71-92

CODEN: ARENAA; ISSN: 0066-4170

PUBLISHER: Annual Reviews Inc.

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB A review. The diversity of the Insecta is reflected in the large and varied microbial communities inhabiting the gut. Studies, particularly with **termites** and cockroaches, have focused on the nutritional contributions of gut bacteria in **insects** living on suboptimal diets. The indigenous gut bacteria, however, also play a role in withstanding the colonization of the gut by non-indigenous **species** including pathogens. Gut bacterial consortia adapt by the transfer of plasmids and transconjugation between bacterial strains, and some insect **species** provide ideal conditions for bacterial conjugation, which suggests that the gut is a hot spot for gene transfer. Genomic anal. provides new avenues for the study of the gut microbial community and will reveal the mol. foundations of the relationships between the insect and its microbiome. In this review the intestinal bacteria is discussed in the context of developing our understanding of symbiotic relationships, of multitrophic interactions between **insects** and plant or animal host, and in developing new strategies for controlling insect pests.

REFERENCE COUNT: 123 THERE ARE 123 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 4 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2004:8006 CAPLUS

DOCUMENT NUMBER: 140:232740

TITLE: Caste- and associated gene expression in a lower termite

AUTHOR(S): Scharf, Michael E.; Wu-Scharf, Dancia; Pittendrigh, Barry R.; Bennett, Gary W.

CORPORATE SOURCE: Department of Entomology, Purdue University, West Lafayette, IN, 47907-2089, USA

09886197

SOURCE: GenomeBiology (2003), 4(10), No pp. given
CODEN: GNBFW; ISSN: 1465-6914
URL: <http://genomebiology.com/content/pdf/gb-2003-4-10-r62.pdf>
PUBLISHER: BioMed Central Ltd.
DOCUMENT TYPE: Journal; (online computer file)
LANGUAGE: English

AB Social **insects** such as **termites** express dramatic polyphenism (the occurrence of multiple forms in a **species** on the basis of differential gene expression) both in association with caste differentiation and between castes after differentiation. We have used cDNA macroarrays to compare gene expression between polyphenic castes and intermediary developmental stages of the termite *Reticulitermes flavipes*. We identified differentially expressed genes from 9 ontogenic categories. Quant. PCR was used to quantify precise differences in gene expression between castes and between intermediary developmental stages. We found worker and nymph-biased expression of transcripts encoding termite and endosymbiont cellulases; presoldier-biased expression of transcripts encoding the storage/hormone-binding protein vitellogenin; and soldier-biased expression of gene transcripts encoding 2 transcription/translation factors, 2 signal transduction factors, and 4 cytoskeletal/muscle proteins. The 2 transcription/translation factors showed significant homol. to the bicaudal and bric-a-brac developmental genes of *Drosophila*. Our results show differential expression of regulatory, structural, and enzyme coding genes in association with termite castes and their developmental precursor stages. They also provide the 1st glimpse into how insect endosymbiont cellulase gene expression can vary in association with the caste of a host. These findings shed light on mol. processes associated with termite biol., polyphenism, caste differentiation, and development and highlight potentially interesting variations in developmental themes between **termites**, other **insects**, and higher animals.

REFERENCE COUNT: 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 5 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:453718 CAPLUS
DOCUMENT NUMBER: 139:114573
TITLE: Termite physiology in relation to wood degradation and termite control
AUTHOR(S): Shelton, Thomas G.; Grace, J. Kenneth
CORPORATE SOURCE: Department of Plant and Environmental Protection Sciences, University of Hawaii at Manoa, Honolulu, HI, 96822, USA
SOURCE: ACS Symposium Series (2003), 845(Wood Deterioration and Preservation), 242-252
CODEN: ACSMC8; ISSN: 0097-6156
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal; General Review
LANGUAGE: English

AB A review. The importance of **termites** (order Isoptera) in the degradation of wood (cellulose, hemicellulose, and lignin collectively) is discussed, and the relative contributions of termite enzymes and intestinal microfauna (protozoa and bacteria) are presented. We also provide an overview of the areas of cellulose degradation, and physiol. (enzymic and pheromonal) means of termite control. Discussion includes the currently known hormones and pheromones with application in control measures, and some reasons for their current use (or lack of use) in termite control. **Termites** are social **insects**, and hormonal/pheromonal control measures often do not have the same results as

are expected with solitary pest **species**. Finally, a short discussion of the current trends in research on feeding and foraging behavior of subterranean **termites** is presented.

REFERENCE COUNT: 63 THERE ARE 63 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d 19 40-47 ibib hitstr abs

L9 ANSWER 40 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1948:42746 CAPLUS
DOCUMENT NUMBER: 42:42746
ORIGINAL REFERENCE NO.: 42:8978b-f
TITLE: The influence of vitamin "T" on the form and habits of **insects**
AUTHOR(S): Goetsch, Wilhelm
CORPORATE SOURCE: Forschungsstelle Krumpendorf, Karnten, Carinthia, Austria
SOURCE: Osterr. zool. Z. (1947), 1(No. 3/4)
DOCUMENT TYPE: Journal
LANGUAGE: Unavailable

AB Vitamin "T" is present in the fat of **termites** and other **insects** and also in several of the fungi (hypomycetes, Penicillium, Torula, and others). It is soluble in H₂O and EtOH and is resistant to heat (can be heated up to 120°). Synonyms for vitamin "T" are termitin, insectine, hypomycin, penicin, mycoine. It is most easily prepared from Torula utilis by acidulating the nutrient substrate to pH 2 and extracting with ether. Prepns. were further purified by dialysis. Growths 5, 10, 15 and 20 days old were used; optimum yield was obtained from a 10-day growth. Heating removed the bacteriostatic factor. Extensive chemical tests would be necessary to establish the identity of vitamin "T." From biol. testing it appears to be distinct from any known vitamin. Vitamin "T" in concns. greater than threshold tends to produce giant forms in **insects**. The head and mandibles of **insects** of various **species** were stimulated to grow much larger in comparison with the body as a whole. If vitamin "T" was administered in excessive doses, dwarf **insects** were produced since the processes leading to maturity were stimulated more than those of growth. Giant forms with large heads could not be produced unless sufficient protein was in the insect diet and unless the vitamin "T" was fed before development had proceeded too far. The body proportions of cockroaches and ants approached those of the termite soldier cast. Flies (Drosophila) of the giant type were not only larger than normal but had larger eyes in proportion to the body than had the control flies. Vitamin "T" not only altered the body proportions but also the habits of ants. Those receiving vitamin "T" worked outside the nest while controls of the same age tended to work in the nest chamber. Vitamin "T" increased pigment formation in **insects**.

L9 ANSWER 41 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1948:23544 CAPLUS
DOCUMENT NUMBER: 42:23544
ORIGINAL REFERENCE NO.: 42:5091d-e
TITLE: Vitamin "T" a new growth factor
AUTHOR(S): Goetsch, Wilhelm
SOURCE: Experientia (1947), 3(No. 7), 1-5
CODEN: EXPEAM; ISSN: 0014-4754
DOCUMENT TYPE: Journal
LANGUAGE: Unavailable

AB cf. C.A. 42, 3472g. A vitamin complex has been extracted from

termites and other **insects**; also from *Penicillium*, *Hypomyces*, and some **species** of yeast. When fed to organisms ranging from vertebrates to yeasts, it stimulates assimilation (of protein in vertebrates), increases O consumption, and speeds the mobilization of reserve substances. Evidence is presented that the chief effect is to increase weight and general growth 10-20% even if the controls are fed the same or even less. The same holds if the diets are fortified with vitamins B1 and B2. **Insects** have been raised whose body proportions are greater than any found in nature (*Blattella germanica*, *Periplaneta orientalis*, *Tachycines asynomorus*, *Drosophila melanogaster*).

L9 ANSWER 42 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1948:23543 CAPLUS
DOCUMENT NUMBER: 42:23543
ORIGINAL REFERENCE NO.: 42:5091d-e
TITLE: Vitamin "T" a new growth factor
AUTHOR(S): Goetsch, Wilhelm
SOURCE: Oesterreichische Zoologische Zeitschrift (1947),
1(Nos. 1 and 2), 49-57
CODEN: OZZEAQ; ISSN: 0369-8084
DOCUMENT TYPE: Journal
LANGUAGE: Unavailable

AB cf. C.A. 42, 3472g. A vitamin complex has been extracted from **termites** and other **insects**; also from *Penicillium*, *Hypomyces*, and some **species** of yeast. When fed to organisms ranging from vertebrates to yeasts, it stimulates assimilation (of protein in vertebrates), increases O consumption, and speeds the mobilization of reserve substances. Evidence is presented that the chief effect is to increase weight and general growth 10-20% even if the controls are fed the same or even less. The same holds if the diets are fortified with vitamins B1 and B2. **Insects** have been raised whose body proportions are greater than any found in nature (*Blattella germanica*, *Periplaneta orientalis*, *Tachycines asynomorus*, *Drosophila melanogaster*).

L9 ANSWER 43 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1945:3526 CAPLUS
DOCUMENT NUMBER: 39:3526
ORIGINAL REFERENCE NO.: 39:567i,568a
TITLE: Protective value of asphalt-laminated paper against certain **insects**
AUTHOR(S): Sweetman, H. L.; Bourne, A. I.
SOURCE: Journal of Economic Entomology (1944), 37, 605-9
CODEN: JEENAI; ISSN: 0022-0493
DOCUMENT TYPE: Journal
LANGUAGE: Unavailable

AB A 2-ply asphalt-laminated paper (Kraft) sealed with an asphalt adhesive resisted feeding by 4 **species** of cockroaches, 3 **species** of thysanurans and partial resistance to 1 **species** of subterranean termite. Of the **species** tested, the firebrat (*Thermobia domestica*) and the termite (*Reticulitermes flavipes*) will probably penetrate wrapping paper most rapidly. Addition of pentachlorophenol to the adhesive for fungicidal purposes greatly reduced the attractiveness of the paper to cockroaches, and to all thysanurans except the firebrat. **Termites** built tubes over paper with 0.3-1.0% pentachlorophenol in the adhesive but did not damage the paper.

L9 ANSWER 44 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1944:35730 CAPLUS
DOCUMENT NUMBER: 38:35730
ORIGINAL REFERENCE NO.: 38:5331b-i,5332a-b

TITLE: Laboratory tests of DDT against various insect pests
 AUTHOR(S): Swingle, M. C.; Mayer, E. L.
 SOURCE: Journal of Economic Entomology (1944), 37, 141-2
 CODEN: JEENAI; ISSN: 0022-0493
 DOCUMENT TYPE: Journal
 LANGUAGE: Unavailable
 AB A dust containing 35% DDT, a spray containing 5% DDT and a wetting agent, and DDT

concentrate were tested on 20 **species** of **insects**. Methods are described. American cockroaches (*Periplaneta americana*) were more susceptible to DDT dusts than to NaF or pyrethrum. Bean leaf rollers (*Urbanus proteus*) were controlled by 3% DDT dust and by DDT spray (4 lb. DDT per 100 gal. water); the pyrethrum standard (1.2% pyrethrum) was about as efficient as the DDT dust. Against blister beetles (*Epicauta lemniscata*) 3% DDT dust killed 100% of the **insects** in 3 days; undild. BaSiF₆ killed 96% in this time. Suspensions were less toxic but were more effective than cryolite suspensions. Cabbage looper (*Autographa brassicae*), 100% kill by 3% dust in 2 days; pyrethrum standard killed 96% in the same time. Colorado potato beetle (*Leptinotarsa decemlineata*) 100% kill by spray (1 lb./100 gal. water). Corn leaf hopper (*Peregrinus maidis*) 100% kill by 5% dust in 1 day; pyrethrum dust (10% pyrethrum standard in talc) gave similar results. Cow-pea weevil (*Callosobruchus maculatus*) 100% kill in 2 days when 3% dust was mixed (1:10,000 by weight) with peas containing the weevils. Derris standard (4.8% rotenone) gave similar results. Cross-striped cabbage worm (*Evergestis rimosalis*), 100% kill by 3% dust in 2 days; 90 and 100% kill from spray (4 lb. per 100 gal. water) in 2 and 4 days, resp. Pyrethrum standard resembled the dust in effectiveness but derris standard in a spray (8 lb. per 100 gal. water) was less effective. Garden flea hopper (*Halticus bracteatus*), excellent control by 3% dust and by a spray containing 8 lb. of 5% DDT in 100 gal. of water. A com. pyrethrum dust was relatively ineffective. Harlequin bug, (*Murgantia histrionica*), 90% kill by 1% dust in 2 days; pyrethrum standard gave similar results. Imported cabbage worm (*Pieris rapae*) 100% kill by 5% dust in 2 days. The derris standard gave the same results. A **species** of looper (*Autographa ro.acte.gationis*) suffered 90-100% kill by 1% dusts in 2 days, and 100% kill by 0.6% dust in 3 days; derris dust (0.96% rotenone) killed 89% in 3 days. A spray (4 lb. per 100 gal. water) killed 62% in 6 days vs. PbHasO₄ spray (8 lb. per 100 gal. water) 100% kill in 4 days and derris spray (4 lb. derris standard per 100 gal. water) 46% kill in 6 days. Melon worm (*Diaphania hyalinata*), 90-100% kill by 0.6% dusts in 2 days. Derris dust (0.96% rotenone) killed only 36% in 2 days. Suspensions and sprays of DDT are highly toxic and better than derris spray for this insect. Pickleworm (*Diaphania nitidalis*) 100% kill by 0.6% dust vs. 24% kill by derris dust (0.96% rotenone) in the same time. DDT sprays equaled derris sprays in effectiveness against this insect. Red flour beetle (*Tribolium castaneum*) 100% kill by a 3% dust diluted 1:10,000 in wheat in 2 days. With the same dust at the same dilution 50% and 82% of rice weevils (*Sitophilus oryza*) were killed in 3 and 5 days, resp. Derris 1:200 killed 36% and 100% of the weevils in 4 and 7 days, resp. Southern armyworm (*Prodenia eridania*) 100% kill by 3% dust vs. 48% by pyrethrum in 2 days. A suspension (8 lb. of 5% DDT per 100 gal. water) killed 100% vs. PbHasO₄ spray (8 lb. in 100 gal. water) 97% kill in 2 days. The DDT-sprayed foliage was toxic to the larvae 8 days after application. Spirea aphid (*Aphis spireacolis*) 100% kill by 3% dust; but 0.25% nicotine sulfate spray gave a quicker kill. Squash bug (*Anasa tristis*) 100% kill of 1st and 4th instar nymphs by 3% dust in 2 days; same result for the pyrethrum standard. **Termites** (*Reticulitermes* sp.) no mortality from 3% dust diluted with sand 1:10,000, but the mixture was repellent to the **termites**. Phytotoxicity tests showed that 2 applications of DDT spray (8 lb. of 5% DDT per 100 gal. water) applied 7

days apart did not injure young bean, pea, pumpkin, Swiss chard and collard plants; and 1% of an aqueous suspension of DDT caused no injury to young bean, pumpkin, Swiss chard, potato and collard plants.

L9 ANSWER 45 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1942:37993 CAPLUS
DOCUMENT NUMBER: 36:37993
ORIGINAL REFERENCE NO.: 36:5946b-f
TITLE: Nicotine as an insect fumigant
AUTHOR(S): Richardson, Henry H.; Casanges, A. H.
SOURCE: Journal of Economic Entomology (1942), 35, 242-6
CODEN: JEENAI; ISSN: 0022-0493
DOCUMENT TYPE: Journal
LANGUAGE: Unavailable

AB Laboratory expts. were made with nicotine vapor on 37 **species** of **insects**. Wide specific variations in resistance were noted. Among the susceptible **species** (those showing complete mortality at a concentration of 0.025 mg./l. in 30 min. at 25°) are included various **species** of aphids, 3 **species** of thrips, *Trialeurodes packardii*, *Empoasca fabae*, *Carpocapsa pomonella* (adults and 1st instar larvae), *Bombyx mori* larvae, *Aphidius phorodontis* adults and *Reticulitermes flavipes* workers. Highly resistant (0.06-0.278 mg. per l.) were most of the beetles and the adult honeybee (*Apis mellifera*). The resistance of the aphid, *Myzus persicae*, varied greatly with respect to the host plant from which it was taken. Late instar larvae of *Prodenia eridamia* and *Heliothis armigera* were much more resistant than young larvae, but there was little difference in the effect of toxicity with age of *Bombyx mori* larvae. The formula $CT = K$, in which C is the gas concentration,

T exposure time and K a constant, holds for some **insects** and fumigants but varied greatly for nicotine. The product CT was smallest for the shortest exposures and increased greatly with longer exposures. Gas concentration had a greater effect on toxic efficiency than exposure time. Comparative ratings at the 95% mortality concns. differed sometimes from those made on the basis of 50% mortality concns. Nicotine is more toxic in the laboratory to some **insects** than is HCN. The exposure time for some **species** is as low as 1 min.

L9 ANSWER 46 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1942:1348 CAPLUS
DOCUMENT NUMBER: 36:1348
ORIGINAL REFERENCE NO.: 36:210d-g
TITLE: Phthalonitrile as an insecticide
AUTHOR(S): Swingle, M. C.; Gahan, J. B.; Phillips, A. M.
SOURCE: U. S. Dept. Agr., Bur. Entomol. Plant Quarantine (1941), E-548, 12 pp.
DOCUMENT TYPE: Journal
LANGUAGE: Unavailable

AB Phthalonitrile (o-dicyanobenzene) was tested on 9 **species** of leaf-eating **insects** in comparison with standard insecticides against the resp. **species**. When used in preliminary tests as a dust on foliage, this material was in general superior to the standard insecticide with which it was compared. Fumigation tests with phthalonitrile in closed Petri dishes gave no mortality; this shows the compound to be either a stomach or a contact poison. Sprays made up with various wetting and dispersing agents showed considerable variation in effectiveness. The most satisfactory spray used on cruciferous plants was made by dissolving the phthalonitrile in acetone and adding the solution to water containing saponin. In cage tests with various concns. of spray, phthalonitrile was effective when used at 2 lb./100 gal. At very dilute

concns. it was not so effective as the standard insecticides. Small field plots of collard and pumpkin plants were sprayed with an 8:8:100 concentration of

phthalonitrile with bentonite, and leaf samples taken from the plots every 2 days were fed to **insects** in the laboratory. The leaves were toxic to larvae for the 1st 96 hrs. after spraying but were almost nontoxic thereafter. An 8:100 concentration of spray applied to several varieties of truckcrop plants caused no injury in 24 days. Phthalonitrile was effective against **termites** when applied as a soil treatment at a concentration of 1:3000.

L9 ANSWER 47 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1934:20157 CAPLUS
DOCUMENT NUMBER: 28:20157
ORIGINAL REFERENCE NO.: 28:2420i,2421a-c
TITLE: The digestion of wood by insect larvae
AUTHOR(S): Mansour, K.; Mansour, J. J.
SOURCE: Proc. Acad. Sci. Amsterdam (1933), 36, 795-9
DOCUMENT TYPE: Journal
LANGUAGE: Unavailable

AB The larvae of 2 wood-feeding Coleoptera were free from associated microorganisms, either intracellularly or in the alimentary canal; nevertheless both digest wood rapidly. Larvae of *Macrotoma palmata* yielded a stomach juice rich in cellulose-splitting enzyme; in 48 hrs., at pH 6.3, 0.5 cc. of juice hydrolyzed 28% of the 6 mg. purified filter paper added. The animal lives on the wood of *Morus alba*, which contains less than 0.5% of total sugar and starch; it is, therefore, dependent for nutrition on the hydrolysis of cellulose. Larvae of *Xystrocera globosa* yielded no cellulose-splitting enzyme, but only an active amylase. Correspondingly it is found only in the sapwood (6.2% sugar and 3.9% starch), and not the heartwood, of its host (*Albizia lebbek*). The animal, therefore, depends on these constituents of the wood, and correspondingly the amount of its excreta is very large. Wood-feeding **insects** thus belong to 3 types: those like the 1st above, which contain a cellulase; those like the 2nd, which do not, and are therefore restricted to woods with a high content of starch or sugar; and finally those like **termites**, which harbor various microorganisms able to digest cellulose. A number of the **species** recorded in the literature are assigned to these 3 groups.

=> d 19 30-39 ibib hitstr abs


L9 ANSWER 30 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1979:505471 CAPLUS
DOCUMENT NUMBER: 91:105471
TITLE: Some effects of juvenile hormone analogs on laboratory groups of *Kaloterms flavicollis* and *Coptotermes amanii* (Isoptera: Kalotermitidae, Rhinotermitidae) at different levels of nutrition
AUTHOR(S): Lenz, Michael
CORPORATE SOURCE: Bundesanst. Materialpruef., Berlin, D-1000.45, Fed. Rep. Ger.
SOURCE: Beihefte zu Material und Organismen (1976), 3 (Org. Holz), 377-92
CODEN: MOBHAK; ISSN: 0375-9318
DOCUMENT TYPE: Journal
LANGUAGE: German

AB After feeding on juvenile hormone analog (JHA) on filter paper, an average of 27% (12-61%) of *K. flavicollis* pseudergates molted into presoldiers (PS),

even though only a limited number of **insects** are normally competent to do so. In *C. amanii*, 50% (42-57%) of the workers changed into PS compared with only 4% in the controls. Comparing pine wood with and without decay (brown rot), both termite **species** showed higher mortality on the latter. Despite twice as much consumption (JHA ingestion) of decayed wood by *K. flavicollis*, formation of PS was similar in both series. When the groups were reexamd. 1 yr later, a difference of survival rates had continued, but all had recommenced egg production. These PS and soldiers which formed under the influence of JHA were easily distinguishable even after 1 yr, because PS had not molted further. During the expts., many PS formed in *C. amanii* groups died during the molt and thus fewer PS were observed on the decayed wood than on the decayed series. However, in these latter groups, the occurrence of soldier/worker intercastes increased with increasing concns. of JHA. When the **termites** were given a choice between decayed wood, with and without the addition of JHA, *K. flavicollis* formed similar nos. of PS, whereas *C. amanii* formed only a few more than in untreated controls. This may be due, not only to differences in termite behavior, but primarily to the variations in test conditions which exposed the *K. flavicollis* to greater amts. of JHA vapor than *C. amanii*. Apparently, the extent to which **termites** were affected by JHA was dependent on the quality of the food available.

L9 ANSWER 31 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1976:587432 CAPLUS
 DOCUMENT NUMBER: 85:187432
 TITLE: The effect of dieldrin coverspraying on populations of night-flying **insects**
 AUTHOR(S): Van Ark, H.
 CORPORATE SOURCE: Plant Prot. Res. Inst., Pretoria, S. Afr.
 SOURCE: Phytophylactica (1976), 8(2), 31-6
 CODEN: PPPMA9; ISSN: 0370-1263
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB Most of some 70 groups or insect **species** studied were not affected by the application of dieldrin (I) [60-57-1] at 93-161 g/ha to natural vegetation. A few were to some extent reduced in nos., but these effects disappeared within 2 months after treatment. The only **species** that was seriously reduced was *Pseudohippopsis filiformis*. The I treatment may have been responsible for the fact that population ds. of *Grammodes euclidioides* and *Platymetopus figuratus* were larger in the treated than the adjacent control areas. The method of assessment was not particularly sensitive because the effects of the insecticide could not be separated from the effects of other environmental factors. Moreover, immigration of **insects** from outside the exptl. areas was one of

Germany
 SOURCE: Holzforschung (1964), 18(1-2), 38-47
 CODEN: HOLZAZ; ISSN: 0018-3830

DOCUMENT TYPE: Journal
 LANGUAGE: Unavailable

AB The woods examined were: *Staudtia stipitata*; *Ocotea rodiaei*; *Afzelia*; *Mansonina altissima*; *Chlorophora excelsa*; and *Sequoia sempervirens*. The origin, com. uses, ds., and important extractives of the above woods are given. For the 1st time, the problem of possible changes in the natural durability of these **species** was studied systematically. To determine the reasons for their durability and its extent, the respective sawdusts and their extractives were exposed to the action of **termites** (*Reticulitermes lucifugus*). Small wood samples were also subjected to artificial weathering in a "Garner-rad" apparatus, which is illustrated and the operation of which is explained. The weathering periods were increased gradually, following a geometric progression. It had been shown initially that all of the 6 **species** were more or less toxic to **insects** and that this was responsible for their durability. Very similar results had been obtained previously. The present work indicated that this native durability did not have a constant value, that it decreased with the duration of the weathering period, and that it also depended on the **species** studied. B. and A prefer the term "resistance behavior." Thus, the decrease in resistance behavior was far greater in the case of Redwood and Iroko wood than for *Afzelia*, *S. stipitata*, *O. rodii*, and *M. altissima*. 29 references.

L9 ANSWER 36 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1964:55816 CAPLUS
 DOCUMENT NUMBER: 60:55816
 ORIGINAL REFERENCE NO.: 60:9840h,9841a-b
 TITLE: Toxicity of dieldrin-concrete mixtures to **termites**
 AUTHOR(S): Allen, T. C.; Esenther, G. R.; Lichtenstein, E. P.
 CORPORATE SOURCE: Univ. of Wisconsin, Madison
 SOURCE: Journal of Economic Entomology (1964), 57(1), 26-9
 CODEN: JEENAI; ISSN: 0022-0493
 DOCUMENT TYPE: Journal
 LANGUAGE: Unavailable

AB Exposure of *Reticulitermes flavipes* on the surface of 0.1% and 1.6% dieldrin-concrete mixture resulted in knockdown and death. Laboratory aging of mixts. caused an initial reduction in surface toxicity leaving a stable residual toxicity. After 16 months the residual toxicity caused knockdown. Newly cracked surfaces of laboratory-aged mixture were equivalent in toxicity to the original surface of new mixture. After 22 months dieldrin-concrete posts set in clay loam soil at Madison, Wisconsin, have shown no reduction in surface toxicity to *R. flavipes*. Dieldrin concentration in the soil, 11/2 years after a 1.6% dieldrin-concrete post had been put in the ground, was 2.49 p.p.m. at 2 in. and 0.49 p.p.m. at 6 in. from the post. Ten **species** of **termites** exposed to 0.1% and 1.6% dieldrin-concrete blocks were affected. The dieldrin concentration in the 0.1% blocks was 542 ± 7 p.p.m. The results of exposure indicated that 4 subterranean **species** were more sensitive to dieldrin-concrete poisoning than were 5 nonsubterranean **species** (not including 1 **species** in which only debilitated **insects** were available).

L9 ANSWER 37 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1961:114019 CAPLUS

DOCUMENT NUMBER: 55:114019
 ORIGINAL REFERENCE NO.: 55:21452c-f
 TITLE: Relation of lipide adsorptivity of powders to their suitability as insecticide diluents
 AUTHOR(S): Ebeling, Walter; Wagner, Robert E.
 CORPORATE SOURCE: Univ. of California, Los Angeles
 SOURCE: Hilgardia (1961), 30(No. 18), 565-86
 CODEN: HILGA4; ISSN: 0073-2230
 DOCUMENT TYPE: Journal
 LANGUAGE: Unavailable

AB Pretreatment of *Drosophila pseudoobscura* with sorptive powders (Olantha clay, Pikes Peak clay, and Santocel C) before treatment with toxicants resulted in a period 3.3 times longer to bring about knockdown than when pretreated with nonsorptive powders (walnut shell flour, Mississippi Diluent, and blue talc). In the pretreatment of 2 **species** of **termites** and 2 **species** of cockroaches, the effect was just the opposite. When used as diluents for the toxicants, the sorptive powders were less effective than the nonsorptive powders. Pretreatment of *D. pseudoobscura* with sorptive powders decreased the toxic action of organic P compds., but increased that of lindane and Sevin; with the German cockroach the toxic action of all toxicants was increased. When used as diluents, the sorptive powders decreased the toxic action of all organic P compds. against both **insects**. With lindane, there was no difference between the 2 powders. With Sevin, the sorptive diluents increased the toxic action against the cockroaches but not against *D. pseudoobscura*. Chlordan and dieldrin resulted in a more rapid knockdown when diluted with nonsorptive powders. Sorptive diluents had the most deleterious effect when used with toxicants in a liquid state. Dibrom, DDVP, and Dylox were more adversely affected than parathion and malathion. Pyrophyllite mixture was superior as an insecticide when organic P toxicants were used. Residues of DDVP in pyrophyllite and Pikes Peak clay, allowed to age indoors for 3 months, resulted in an increased knockdown period.

L9 ANSWER 38 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1954:5042 CAPLUS
 DOCUMENT NUMBER: 48:5042
 ORIGINAL REFERENCE NO.: 48:940e-h
 TITLE: Attempts to control subterranean pests
 AUTHOR(S): Bertels, Andre
 CORPORATE SOURCE: Inst. Agron. Sul, Pelotas, Rio Grande do Sul, Brasil, Arg.
 SOURCE: Agros (1951), 4, 140-9
 DOCUMENT TYPE: Journal
 LANGUAGE: Unavailable

AB Three types of insecticides, dusts, solns., and gases, were used. Toxicants in the dusts were either DDT or benzene hexachloride, the DDT being used at 3%, 5%, and 10%, resp., and the benzene hexachloride at 15% and 25%, resp. Solns. were either DDT or thiophosphate formulations. Gases included vaporized DDT, CH₃Br, CS₂, and CCl₄. Depth of 10 cm. was the optimum placement for dusts. Formulations with each of the toxicants in dust were effective but benzene hexachloride could not be used on edible root crops. In all types of treatment use of toxicant before seeding was preferable to after seeding. The period after treatment when seeding was safe differed with the host plant. Thus for benzene hexachloride used at a level of 30 g. per planting hole (Gammexane 15 and Gammexane 25), 94% of potato plants were killed when planted 14 days after treatment, 53.5% of cucumber seedlings were killed, and only 6.3% of transplanted cabbage. In the CH₃Br treatment, lettuce, carrots, red beets, peas, potatoes, and sugar cane suffered great damage. Turnips were killed. Some of the Compositae and Graminaceae were resistant to CH₃Br.

09886197

Neither CS2 nor CCl4 killed plants. Types of **insects** found included **termites**, beetles (larvae and adults), caterpillars, and ants. Dosages for individual **species** are not indicated.

L9 ANSWER 39 OF 47 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1953:63648 CAPLUS

DOCUMENT NUMBER: 47:63648

ORIGINAL REFERENCE NO.: 47:10797h-i

TITLE: The control of powder-post beetles in buildings

AUTHOR(S): Tooke, F. G. C.

SOURCE: Farming in S. Africa (1953), 28, 79-83

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

AB The wood-destroying **species** found in the coastal belt are: West Indian drywood termite (*Cryptotermes brevis*), European house borer (*Hylotrupes bajulus*), furniture beetles (*Anobium punctatum* and *Nicobium castaneum*) and powder-post beetle (*Lyctus brunneus*). Control of these **insects** is obtained by treatment with pentachlorophenol (5%) and DDT (5%) in white spirits.

=> d his

(FILE 'HOME' ENTERED AT 16:06:53 ON 21 JUL 2004)

FILE 'STNGUIDE' ENTERED AT 16:07:06 ON 21 JUL 2004

FILE 'REGISTRY' ENTERED AT 16:08:17 ON 21 JUL 2004

L1 0 S IMIDALCLOPRID

FILE 'CAPLUS' ENTERED AT 16:09:11 ON 21 JUL 2004

L2 1376 S IMIDACLOPRID

L3 22 S L2 AND TERMITES

L4 25 S L2 AND WOOD

L5 6 S L3 AND L4

L6 2129 S TERMITES

L7 29825 S INSECTS

L8 274 S L6 AND L7

L9 47 S L8 AND SPECIES

=> s l2 and l8

L10 4 L2 AND L8

=> d l10 ibib hitstr abs

L10 ANSWER 1 OF 4 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:799698 CAPLUS

DOCUMENT NUMBER: 132:9953

TITLE: Termite control

INVENTOR(S): De Villiers, Vivian; Van der Westhuizen, M. C.;

Robbertse, Ernest

PATENT ASSIGNEE(S): Bayer A.-G., Germany

SOURCE: S. African, 16 pp.

CODEN: SFXXAB

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.

KIND DATE

APPLICATION NO. DATE

```

-----
ZA 9711701      A    19980706      ZA 1997-11701    19971230
AP 1174         A    20030630      AP 1998-1424     19981228
W:  BW, GH, GM, KE, LS, MW, SD, SZ, UG, ZM, ZW
BR 9805735     A    20010424      BR 1998-5735     19981229

```

PRIORITY APPLN. INFO.: ZA 1997-11701 A 19971230

AB Agonists or antagonists of nicotinerbic acetylcholine receptors of **insects** are used for the control of harvester **termites**, i.e. Hodotermidae. **Imidacloprid** is the preferred active ingredient. The bait formulations comprise lucerne or grass particles.

=> d 110 2-4 ibib hitstr abs

L10 ANSWER 2 OF 4 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:125767 CAPLUS

DOCUMENT NUMBER: 130:178773

TITLE: Composition for the control of wood-destroying **insects**, especially **termites**

INVENTOR(S): Anderson, John-phillip-evans; Keuken, Oliver

PATENT ASSIGNEE(S): Bayer A.-G., Germany

SOURCE: Eur. Pat. Appl., 21 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 896791	A2	19990217	EP 1998-114187	19980729
EP 896791	A3	20000112		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
DE 19734665	A1	19990218	DE 1997-19734665	19970811
TW 505500	B	20021011	TW 1998-87112592	19980731
US 6264968	B1	20010724	US 1998-128818	19980804
ZA 9807118	A	19990209	ZA 1998-7118	19980807
JP 11124302	A2	19990511	JP 1998-234861	19980807
AU 9879895	A1	19990218	AU 1998-79895	19980811
AU 768390	B2	20031211		
BR 9803138	A	19991221	BR 1998-3138	19980811

PRIORITY APPLN. INFO.: DE 1997-19734665 A 19970811

AB The title comps. (no examples) comprise an insecticide, preferably **imidacloprid**, incorporated into an organic natural and/or synthetic carrier. Optional ingredients are insect attractants and microbicides.

L10 ANSWER 3 OF 4 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1996:411657 CAPLUS

TITLE: **Imidacloprid** - chemical synergist for microbial control agents of **termites**.

AUTHOR(S): Boucias, D. G.

CORPORATE SOURCE: Department Entomology & Nematology, University Florida, Gainesville, FL, 32611-0620, USA

SOURCE: Book of Abstracts, 212th ACS National Meeting, Orlando, FL, August 25-29 (1996), AGRO-019. American Chemical Society: Washington, D. C.

CODEN: 63BFAF

DOCUMENT TYPE: Conference; Meeting Abstract

LANGUAGE: English

AB Our research has determined that the neurotoxin, **imidacloprid**, at sublethal concns., can significantly alter the behavioral patterns of **insects**. For example, the subterranean termite, *Reticulotermis flavipes* possesses social behaviors (grooming, tunnel construction) which serve as the primary line of defense against pathogenic and opportunistic microorganisms. These behaviors, in combination with the resident microflora, confer a high degree of disease resistance upon these social **insects**. Exposure to low dosages of **imidacloprid** produces a long term disruption of these social behaviors resulting in the onset of epizootics initiated by either resident or introduced microbes. Related studies on other nonsocial **insects** (cockroaches, weevils) have supported the results found with **termites**. At sublethal concns., **imidacloprid** acted as a behavioral modifying agent significantly increasing the host **insects** susceptibility to microbial control agents.

L10 ANSWER 4 OF 4 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1995:187187 CAPLUS

DOCUMENT NUMBER: 122:25815

TITLE: **Imidacloprid** - a new systemic insecticide.

AUTHOR(S): Elbert, A.; Becker, B.; Hartwig, J.; Erdelen, C.

CORPORATE SOURCE: Geschäftsbereich Pflanzenschutz
Entwicklung/Insektizide, Bayer AG, Leverkusen, 5090,
Germany

SOURCE: Pflanzenschutz-Nachrichten Bayer (German Edition)
(1991), 44(2), 113-36

CODEN: PNBAT; ISSN: 0340-1723

PUBLISHER: Bayer AG

DOCUMENT TYPE: Journal

LANGUAGE: German

AB The biol. profile of **Imidacloprid** (I) was defined on the basis of the results of exhaustive laboratory expts. and greenhouse trials. I is extremely effective against sucking **insects**, such as rice leafhoppers, aphids, thrips and mealybugs, and very effective against whitefly. It is also effective against some species of biting **insects**, such as paddy stem borers and Colorado beetle, but it has no effect on nematodes or spider mites. At comparatively high doses it kills adult **insects** and has ovicidal effects. I is a nicotinic acetylcholine receptor stimulator. Its mechanism of action differs from that of conventional insecticides. It therefore gives excellent control of all resistant populations investigated hitherto. I has a pos. temperature coefficient After foliar application, it has a good residual action, it is highly photostable and it shows satisfactory resistance to rain. I is active after oral ingestion and by direct contact, but it is not active in the vapor phase. The LD95 after oral ingestion by *Myzus persicae* is .apprx.2 pg/aphid. After topical application it is .apprx.160 pg/aphid. It has not been possible to demonstrate recovery of injured aphids, or antifeeding effects. I has a faster action against aphids than oxydemeton-Me. After foliar application, I shows good translaminar and acropetal translocation, so it is also likely to provide effective control of pests with a furtive lifestyle, and protect the parts of the plant which regenerate after treatment. By virtue of its good contact action and powerful systemic action after uptake through the root system, I can be applied to soil and used as a seed dressing. It gives excellent control of pests such as onion maggots, *Diabrotica*, wire worms, **termites** and fire ants which live in the soil, and of **insects** such as aphids which live above ground level. It has a good residual action after application to the soil and when it is used as a seed dressing. The compatibility of I with plants is good after use as a seed dressing, as a soil treatment and after foliar application. By

09886197

virtue of its biol. properties, I is likely to have a wide range of uses for controlling economically important pests of rice, cotton, cereals, maize, sugar beet, potatoes, vegetables, citrus fruit, pome and stone fruit and other crops.

=>

=> d his

(FILE 'HOME' ENTERED AT 16:06:53 ON 21 JUL 2004)

FILE 'STNGUIDE' ENTERED AT 16:07:06 ON 21 JUL 2004

FILE 'REGISTRY' ENTERED AT 16:08:17 ON 21 JUL 2004

L1 0 S IMIDACLOPRID

FILE 'CAPLUS' ENTERED AT 16:09:11 ON 21 JUL 2004

L2 1376 S IMIDACLOPRID
L3 22 S L2 AND TERMITES
L4 25 S L2 AND WOOD
L5 6 S L3 AND L4
L6 2129 S TERMITES
L7 29825 S INSECTS
L8 274 S L6 AND L7
L9 47 S L8 AND SPECIES
L10 4 S L2 AND L8

=> s l9 and l2

L11 1 L9 AND L2

=> d l11 ibib hitstr abs

L11 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1995:187187 CAPLUS

DOCUMENT NUMBER: 122:25815

TITLE: Imidacloprid - a new systemic insecticide.

AUTHOR(S): Elbert, A.; Becker, B.; Hartwig, J.; Erdelen, C.

CORPORATE SOURCE: Geschäftsbereich Pflanzenschutz
Entwicklung/Insektizide, Bayer AG, Leverkusen, 5090,
Germany

SOURCE: Pflanzenschutz-Nachrichten Bayer (German Edition)
(1991), 44(2), 113-36

CODEN: PNBAT; ISSN: 0340-1723

PUBLISHER: Bayer AG

DOCUMENT TYPE: Journal

LANGUAGE: German

AB The biol. profile of **Imidacloprid** (I) was defined on the basis of the results of exhaustive laboratory expts. and greenhouse trials. I is extremely effective against sucking **insects**, such as rice leafhoppers, aphids, thrips and mealybugs, and very effective against whitefly. It is also effective against some **species** of biting **insects**, such as paddy stem borers and Colorado beetle, but it has no effect on nematodes or spider mites. At comparatively high doses it kills adult **insects** and has ovicidal effects. I is a nicotinic acetylcholine receptor stimulator. Its mechanism of action differs from that of conventional insecticides. It therefore gives excellent control of all resistant populations investigated hitherto. I has a pos. temperature coefficient After foliar application, it has a good residual action, it is highly photostable and it shows satisfactory resistance to rain. I is active after oral ingestion and by direct contact, but it is not active in

the vapor phase. The LD95 after oral ingestion by Myzus persicae is .apprx.2 pg/aphid. After topical application it is .apprx.160 pg/aphid. It has not been possible to demonstrate recovery of injured aphids, or antifeeding effects. I has a faster action against aphids than oxydemeton-Me. After foliar application, I shows good translaminar and acropetal translocation, so it is also likely to provide effective control of pests with a furtive lifestyle, and protect the parts of the plant which regenerate after treatment. By virtue of its good contact action and powerful systemic action after uptake through the root system, I can be applied to soil and used as a seed dressing. It gives excellent control of pests such as onion maggots, Diabrotica, wire worms, **termites** and fire ants which live in the soil, and of **insects** such as aphids which live above ground level. It has a good residual action after application to the soil and when it is used as a seed dressing. The compatibility of I with plants is good after use as a seed dressing, as a soil treatment and after foliar application. By virtue of its biol. properties, I is likely to have a wide range of uses for controlling economically important pests of rice, cotton, cereals, maize, sugar beet, potatoes, vegetables, citrus fruit, pome and stone fruit and other crops.

=>

=> logoff

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

LOGOFF? (Y)/N/HOLD:H

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

238.44

244.04

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-59.54

-59.54

SESSION WILL BE HELD FOR 60 MINUTES

STN INTERNATIONAL SESSION SUSPENDED AT 16:24:10 ON 21 JUL 2004

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssspta1202sxq

PASSWORD:

* * * * * RECONNECTED TO STN INTERNATIONAL * * * * *

SESSION RESUMED IN FILE 'CAPLUS' AT 16:25:08 ON 21 JUL 2004

FILE 'CAPLUS' ENTERED AT 16:25:08 ON 21 JUL 2004

COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

238.44

244.04

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-59.54

-59.54

09886197

=> s isoptera

L12 570 ISOPTERA

=> s l12 and l2

L13 13 L12 AND L2

=> d l13 1-13 ibib hitstr abs

L13 ANSWER 1 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2004:453202 CAPLUS

DOCUMENT NUMBER: 141:23526

TITLE: Novel pyrazole-based anthranilamide insecticides and their preparation, compositions, and use

INVENTOR(S): Hughes, Kenneth Andrew; Lahm, George Philip; Selby, Thomas Paul

PATENT ASSIGNEE(S): E.I. Du Pont De Nemours and Company, USA

SOURCE: PCT Int. Appl., 96 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1


PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004046129	A2	20040603	WO 2003-US36167	20031112
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			

PRIORITY APPLN. INFO.:

US 2002-426693P P 20021115

GI



AB The invention provides title compds. I and their N-oxides and suitable

salts [wherein: Y, V = N or CR_{4a}; W = N, CH, or CR₆; R₁ = H, (un)substituted alkyl, alkenyl, alkynyl or cycloalkyl, alkylcarbonyl, alkoxy carbonyl, (di)alkylaminocarbonyl; R₂ = H, alkyl, alkenyl, alkynyl, cycloalkyl, alkoxy, (di)alkylamino, cycloalkylamino, alkoxy carbonyl, or alkylcarbonyl; R₃ = H, G, (un)substituted alkyl, alkenyl, alkynyl or cycloalkyl; or NR₂R₃ = (un)substituted heterocyclic (N/O/S) ring; G = (un)substituted 5- or 6-membered non-aromatic carbo- or heterocyclic ring; R_{4a}, R_{4b} = H, various carbon and heteroat. substituents; R₅ = alk(en/yn)yl, various derivs. of OH, SH, and NH₂; R₆ = (halo)alk(en/yn)yl, OH and derivs. or thio analogs, halo, cyano, CO₂H, (di)alkylamino, (un)substituted Ph, PhCH₂, PhCO, PhO, etc.; n = 0-4]. The invention also pertains to compns. for controlling invertebrate pests, comprising a biol. effective amount of I, their N-oxides, or their agronomically or nonagronomically suitable salts, and at least one addnl. component selected from surfactants, solid diluents, and liquid diluents, and optionally further comprising an effective amount of at least one addnl. biol. active compound or agent. Also disclosed are methods for controlling invertebrate pests by contact of the pests or their environment with said compds. Eighteen compds. I were prepared and tested. For instance, 3-chloro-2-hydrazinopyridine was cyclocondensed with di-Et maleate to give 55% Et 1-(3-chloro-2-pyridinyl)-3-pyrazolidinone-5-carboxylate, which was oxidized to a dihydropyrazolone, saponified to an acid, cyclized with dichloroanthranilic acid to give a benzoxazinone, O-mesylated at the pyrazolone, and ring-opened with MeNH₂, to give invention compound II. In a test of larval *Plutella xylostella* on radish plants, II at 50 ppm (spray) reduced feeding damage by 80% or more. Compds. I were also effective against *Spodoptera frugiperda*, *Myzus persicae*, and *Empoasca fabae*.

L13 ANSWER 2 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:336622 CAPLUS

DOCUMENT NUMBER: 139:48626

TITLE: Effects of exposure duration on transfer of nonrepellent termiticides among workers of *Coptotermes formosanus* Shiraki (**Isoptera**: Rhinotermitidae)

AUTHOR(S): Shelton, Thomas G.; Grace, J. Kenneth

CORPORATE SOURCE: Department of Plant & Environmental Protection Sciences, University of Hawaii, Honolulu, HI, 96822-2271, USA

SOURCE: Journal of Economic Entomology (2003), 96(2), 456-460
CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The potential for transfer of nonrepellent termiticide toxicants between workers of the Formosan subterranean termite, *Coptotermes formosanus* Shiraki, was examined using two com. available pesticide formulations and a simple donor-recipient model modified from current methods in the literature. Pesticides used were **imidacloprid**, formulated as Premise 75 WP, and fipronil, formulated as Termidor SC, in concns. of 1, 10, and 100 ppm (weight of active ingredient/weight of sand) in sand. A significant increase was shown in recipient mortality over control mortality when donor workers were treated with 100 ppm **imidacloprid** or 100 ppm fipronil. Although all three colonies studied were affected, one colony (colony 3) was affected to a significantly greater extent than the other colonies. This effect was not correlated with termite body size (dry mass). In a second study, recipient mortality was evaluated after exposure of donors to 1 ppm insecticide for 3, 6, 12, or 24 h. Recipient mortality indicated that these exposures did not consistently lead to lethal transfer of the

09886197

insecticides.

REFERENCE COUNT: 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 3 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:177184 CAPLUS

DOCUMENT NUMBER: 138:333176

TITLE: Effect of **imidacloprid** tree treatments on the occurrence of formosan subterranean termites, *Coptotermes formosanus* Shiraki (**Isoptera**: Rhinotermitidae), in independent monitors

AUTHOR(S): Osbrink, Weste L. A.; Lax, Alan R.

CORPORATE SOURCE: Southern Regional Research Center, USDA-ARS, New Orleans, LA, 70124, USA

SOURCE: Journal of Economic Entomology (2003), 96(1), 117-125
CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Periodic sampling of 87 independent monitors, initially active with the Formosan subterranean termite, *Coptotermes formosanus* Shiraki, was conducted. Monitors, located in eight sectors adjacent to seven buildings, were various distances (1-46 m) from 57 trees treated with 0.1% **imidacloprid** foam. Termites collected from six of the eight sectors showed latent mortality attributed to **imidacloprid** intoxication at all monitor-tree distances. Approx. 6 mo after treatment, termite populations had recovered in these sectors. Another sector showed termite population suppression for ≈ 15 mo, followed by recovery. **Imidacloprid** tree treatments did not control *C. formosanus* populations in independent monitors adjacent to the treatments.

REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 4 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:777603 CAPLUS

DOCUMENT NUMBER: 137:274431

TITLE: Insecticide compositions containing amino acids,

INVENTOR(S): Sandeman, Richard Mark; Chandler, David Spencer; Duncan, Ann Maree; Hay, Phillip Maxwell

PATENT ASSIGNEE(S): Nufarm Limited, Australia; La Trobe University

SOURCE: PCT Int. Appl., 62 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002078448	A1	20021010	WO 2002-AU389	20020328
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			

09886197

EP 1385379 A1 20040204 EP 2002-712624 20020328

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR

PRIORITY APPLN. INFO.: AU 2001-4069 A 20010329
WO 2002-AU389 W 20020328

OTHER SOURCE(S): MARPAT 137:274431

AB Insecticides of formula $R_3N(R_2)AC(R_1)(:O)$ and the agriculturally acceptable salts thereof ($R_1 = OR_5$ wherein $R_5 = H$, (un)substituted alkyl, (un)substituted aryl, (un)substituted cycloalkyl, (un)substituted heterocyclic; NR_6OH wherein $R_6 = H$, (un)substituted alkyl, (un)substituted aryl, (un)substituted carbocyclic; NR_7R_8 wherein R_7 and $R_8 = H$, (un)substituted alkyl, (un)substituted aryl and carbocyclic; and wherein R_1 is linked to R_2 to form a diradical bridging group; R_2 and $R_3 = H$, (un)substituted alkyl, (un)substituted carbocyclic, (un)substituted aryl, and (un)substituted acyl; and $A =$ diradical linking group, which has a mol. weight of preferably less than 200 and more preferably less than 100) are used to control insect species selected from the orders Lepidoptera, Hemiptera, Orthoptera, Coleoptera, Psocoptera, **Isoptera**, Thysanoptera and Homoptera on cotton.

REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 5 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:720924 CAPLUS

DOCUMENT NUMBER: 135:340463

TITLE: Chemical prevention of colony foundation by
Cryptotermes brevis (**Isoptera**:
Kalotermitidae) in attic modules

AUTHOR(S): Scheffrahn, Rudolf H.; Busey, Philip; Edwards, Jeffrey
K.; Krecek, Jan; Maharajh, Boudanath; Su, Nan-Yao

CORPORATE SOURCE: Ft. Lauderdale Research and Education Center,
University of Florida, Fort Lauderdale, FL, 33314, USA

SOURCE: Journal of Economic Entomology (2001), 94(4), 915-919
CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Disodium octaborate tetrahydrate (DOT) dust, DOT aqueous solution, **imidacloprid** dust, and amorphous silica gel dust with synergized 1% pyrethrins were applied on wood surfaces to simulated attic modules. Modules (30 by 30 cm) with and without fiberglass insulation were exposed to dispersal flights of *Cryptotermes brevis* (Walker) in May and June of 1998 and 1999. Six months after flights, modules were disassembled and inspected for nuptial chamber location and contents. During both years, air and water control treatments contained 22.2 ± 9.94 (mean \pm SD) nuptial chambers, 7.5 ± 5.7 live imagoes, and 2.0 ± 1.4 chambers with brood. This survivorship indicated that the attic modules performed well as a colonizing platform for *C. brevis*. *C. brevis* dealates preferred constructing nuptial chambers in the crevices at the bases or tops of the modules instead of internal crevices. Modules treated in 1998 and 1999 with DOT or silica dusts contained no live termites, whereas zero of five modules treated with **imidacloprid** dust in 1998 and two of 20 modules treated with **imidacloprid** dust in 1999 contained single live incipient colonies. In 1998, 15% DOT solution, applied as a postconstruction treatment, yielded significantly fewer chambers and live termites than controls, but was not as effective as dusts in preventing successful colonization. In 1999, the DOT solution, applied as a construction-phase treatment, was equally as effective in preventing colonization as the dust treatments during that year. Results indicate that dust formulations of DOT, silica gel, and **imidacloprid** can

be used to prevent drywood termite colonization in existing building voids and attics. Where the entire wood framing is exposed to treatment, such as during building construction, aqueous DOT solution can be equally effective as

dusts in preventing colonization by *C. brevis*.

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 6 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:336305 CAPLUS

DOCUMENT NUMBER: 135:1645

TITLE: Effects of sublethal exposure to **imidacloprid** on subsequent behavior of subterranean termite *Reticulitermes virginicus* (**Isoptera**: Rhinotermitidae)

AUTHOR(S): Thorne, Barbara L.; Breisch, Nancy L.

CORPORATE SOURCE: Department of Entomology, University of Maryland, College Park, MD, 20742, USA

SOURCE: Journal of Economic Entomology (2001), 94(2), 492-498
CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Expts. were conducted to determine whether subterranean termites, *Reticulitermes virginicus* (Banks), previously exposed to sublethal doses of **imidacloprid** (Premise), and allowed to recover for 1 wk, demonstrated behavioral aversion to a subsequent exposure. Worker termites experiencing a previous sublethal but debilitating exposure to **imidacloprid**-treated sand (either 10 or 100 ppm for 4 h) showed no apparent aversion to a second encounter with **imidacloprid** -treated sand under conditions of this experiment If these laboratory results hold

in the field and termites traveling through a zone of soil treated with **imidacloprid** are impaired but subsequently recover, they will be just as likely as their naive nestmates to reenter the treated area if their travels take them through the nonrepellent application a second time. Thus, a sublethal exposure to **imidacloprid** can affect termite tunneling behavior. Many worker termites that received an initial 4-h exposure to 100 ppm **imidacloprid**-treated sand died, but those that survived tunneled significantly less than did their naive nestmates, as did some termites exposed to 10 ppm **imidacloprid**.

REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 7 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:666543 CAPLUS

DOCUMENT NUMBER: 133:248390

TITLE: Synergistic insecticidal compositions containing a neuronal sodium channel antagonist and another insecticide

INVENTOR(S): Treacy, Michael Frank; Borysewicz, Raymond Frank; Schwinghammer, Kurt Allen; Rensner, Paul Erich; Oloumi-Sadeghi, Hassan

PATENT ASSIGNEE(S): American Cyanamid Company, USA

SOURCE: PCT Int. Appl., 30 pp.

CODEN: PIXXD2


DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000054591	A2	20000921	WO 2000-US5879	20000307
WO 2000054591	A3	20010118		
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 2000036175	A5	20001004	AU 2000-36175	20000307
AU 765767	B2	20030925		
BR 2000008930	A	20011218	BR 2000-8930	20000307
EP 1198170	A2	20020424	EP 2000-914839	20000307
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL				
NZ 514000	A	20030429	NZ 2000-514000	20000307
JP 2003517455	T2	20030527	JP 2000-604685	20000307
US 6479543	B1	20021112	US 2000-521987	20000309
ZA 2001007484	A	20021201	ZA 2001-7484	20010911
US 2002177597	A1	20021128	US 2002-145784	20020516
PRIORITY APPLN. INFO.:			US 1999-124306P	P 19990312
			US 1999-158201P	P 19991007
			WO 2000-US5879	W 20000307
			US 2000-521987	A3 20000309
OTHER SOURCE(S):			MARPAT 133:248390	
GI				



AB A synergistic insecticidal composition comprises a neuronal sodium channel antagonist such as I (X, Y, Z = H, halo, OH, CN, NO₂, alkyl, etc.; W = O or S; m, p, q = 1, 2, 3, 4, or 5; n = 0, 1, or 2; R, R₁, R₂, R₃ = alkyl) in combination with one or more pyrethroids, pyrethroid-type compds., recombinant nucleopolyhedroviruses expressing an insect toxin, organophosphates, carbamates, formamidines, macrocyclic lactones, amidinohydrazones, GABA antagonists and acetylcholine receptor ligands.

L13 ANSWER 8 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:573349 CAPLUS

DOCUMENT NUMBER: 133:248356

TITLE: Feeding inhibition and mortality in Reticulitermes flavipes (Isoptera: Rhinotermitidae) after exposure to imidacloprid-treated soils

AUTHOR(S): Ramakrishnan, Rathna; Suiter, Daniel R.; Nakatsu, Cindy H.; Bennett, Gary W.

09886197

CORPORATE SOURCE: Center for Urban & Industrial Pest Management,
Department of Entomology, Purdue University, West
Lafayette, IN, 47907, USA
SOURCE: Journal of Economic Entomology (2000), 93(2), 422-428
CODEN: JEENAI; ISSN: 0022-0493
PUBLISHER: Entomological Society of America
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Feeding inhibition and mortality of *Reticulitermes flavipes* (Kollar) exposed to sand, sandy loam, loam, and silty clay loam soils treated with several concns. of **imidacloprid** were studied using bioassay techniques under laboratory conditions. Termite workers stopped feeding after exposure to treated soils. Differences in feeding reduction varied among the soil types. Based on the magnitude of the F-statistics, the effect of **imidacloprid** on the reduction of termite feeding was greatest in sand followed by sandy loam, loam, and silty clay loam soils. Soil properties such as organic matter content, silt and clay proportions, pH, and cation exchange capacity were suggested to affect the bioavailability of **imidacloprid**. Similar soil effects on mortality were observed in termites continuously exposed to treated soil for 21 days. In 3 of 4 soils tested, susceptibility to **imidacloprid** was not affected by the source of the termites tested.

REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 9 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:470450 CAPLUS
DOCUMENT NUMBER: 133:90469
TITLE: Adhesive composition containing insecticides, preservatives, termite repellents and bactericides for lignocellulosic material and its complex
INVENTOR(S): Jaesch, Tohmas; Fushiki, Kiyoyuki; Saito, Takanobu; Katsusawa, Yoshinaga
PATENT ASSIGNEE(S): Bayer A.-G., Germany; Ohshika Shinko K. K.; Chemiholz K. K.
SOURCE: Jpn. Kokai Tokkyo Koho, 12 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000192001	A2	20000711	JP 1998-376942	19981228
KR 2000048138	A	20000725	KR 1999-57526	19991214
EP 1018413	A1	20000712	EP 1999-124843	19991215
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
AU 9965409	A1	20010628	AU 1999-65409	19991222
NZ 502074	A	20020301	NZ 1999-502074	19991223
NO 9906479	A	20000629	NO 1999-6479	19991227
US 2001027217	A1	20011004	US 1999-472589	19991227
BR 9907435	A	20010320	BR 1999-7435	19991228

PRIORITY APPLN. INFO.: JP 1998-376942 A 19981228

AB The composition, for preparation of wood products (e.g., plywood), comprises an adhesive, an organic phenolic composition, an insecticide, a preservative, a termite repellent and a bactericide. Thus, a composition was made from Oshika Resin PWP 60 containing a solution of **imidacloprid** 3, IPBC 20 and 2-phenylphenol 15, and a solvent 62%.

09886197

L13 ANSWER 10 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:797191 CAPLUS

DOCUMENT NUMBER: 132:60446

TITLE: **Imidacloprid**-enhanced *Reticulitermes flavipes* (**Isoptera**: Rhinotermitidae) susceptibility to the entomopathogen *Metarhizium anisopliae*

AUTHOR(S): Ramakrishnan, Rathna; Suiter, Daniel R.; Nakatsu, Cindy H.; Humber, Richard A.; Bennett, Gary W.

CORPORATE SOURCE: Center for Urban & Industrial Pest Management, Department of Entomology, Purdue University, West Lafayette, IN, 47907, USA

SOURCE: Journal of Economic Entomology (1999), 92(5), 1125-1132

CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The effects of **imidacloprid** and the entomopathogen *Metarhizium anisopliae* (Metsch.) Sorokin on the eastern subterranean termite, *Reticulitermes flavipes* (Kollar), were evaluated in a 4 + 3 factorial experiment in both sterile and nonsterile loam soil. Termites were not susceptible to *M. anisopliae* when assays were conducted in nonsterile soil, but were highly susceptible in sterile soil. Termite mortality after 21 days of continuous exposure to 104 conidia per g soil was 0 and 41.6% in nonsterile and sterile soil, resp. Termites were significantly more susceptible to sterile soil containing 107 conidia per g than to the same soil containing 104 conidia per g. In continuous exposure assays, termites were highly susceptible to **imidacloprid**-treated (5,10, and 20 ppm) nonsterile and sterile soil containing no exptl. introduced *M. anisopliae*. Exposure of termites to **imidacloprid** enhanced their susceptibility to introduced *M. anisopliae* in nonsterile and sterile soil. Native entomopathogens recovered from termites exposed to **imidacloprid**-treated, nonsterile soil (i.e., no introduced *M. anisopliae*) included *Conidiobolus coronatus* (Constantin) Batko, *Cunninghamella echinulata* Thaxter, *Fusarium* spp., *Aspergillus* spp., and a naturally occurring strain of *M. anisopliae* variety majus.

REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 11 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1998:631799 CAPLUS

DOCUMENT NUMBER: 129:246360

TITLE: Ant-repellent thermoplastic foam molding compositions containing chloropyridines for thermal insulators

INVENTOR(S): Toyonaga, Yoshihiro; Kanzaki, Masahiro

PATENT ASSIGNEE(S): Shinto Paint Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 10259270	A2	19980929	JP 1997-86053	19970319
PRIORITY APPLN. INFO.:			JP 1997-86053	19970319

AB Title compns. contain acetamiprid (I) or **imidacloprid** as an ant

repellent. Thus, prefoamed polystyrene was coated with a solution containing I and adhesive, mixed with uncoated polystyrene, and heated to give a molding containing 0.2% I. The molding showed bending strength 3.8 kg/cm² and no damage by termite for ≥21 days.

L13 ANSWER 12 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1998:631792 CAPLUS
 DOCUMENT NUMBER: 129:317341
 TITLE: Termite-repellent polyurethane foam molding compositions containing chloropyridines for thermal insulators
 INVENTOR(S): Toyonaga, Yoshihiro; Okuta, Kazuo
 PATENT ASSIGNEE(S): Shinto Paint Co., Ltd., Japan
 SOURCE: Jpn. Kokai Tokkyo Koho, 4 pp.
 CODEN: JKXXAF
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 10259263	A2	19980929	JP 1997-86055	19970319
PRIORITY APPLN. INFO.:			JP 1997-86055	19970319

AB Title comps. contain acetamiprid (I) or **imidacloprid** as termite repellents. Thus, HCFC-containing Polyol GB, I, and polyphenyl-type polyisocyanates were molded to give 0.1% I-containing polyurethane foams showing bending strength 4.5 kg/cm² and no damage by termite for ≥21 days.

L13 ANSWER 13 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN
 ACCESSION NUMBER: 1995:881641 CAPLUS
 DOCUMENT NUMBER: 123:278677
 TITLE: Field tests for control of the mound-building termite *Cornitermes cumulans* (Kollar, 1832) (**Isoptera**, Termitidae)
 AUTHOR(S): Mariconi, F.A.M.; Galan, V.B.; Rocha, M.T.; Maule, R.F.; Passos, H.R.; Silva, R.A.A.
 CORPORATE SOURCE: ESALQ, USP, Piracicaba, 13418-900, Brazil
 SOURCE: Scientia Agricola (Piracicaba, Brazil) (1994), 51(3), 505-8
 CODEN: SGRIEF; ISSN: 0103-9016
 PUBLISHER: Universidade de Sao Paulo, Campus de Piracicaba
 DOCUMENT TYPE: Journal
 LANGUAGE: Portuguese

AB Two field tests were carried out to evaluate the performance of several pesticides for the control of the mound termite pest in pastures. Experiment I: 60 mounds were selected and measured outside. There were 6 treatments with 10 replications: A) abamectin (50 cm³ 1.8% EC); B) silafluofen (200 cm³ 80% EC); C) silafluofen (400 cm³ 80% EC); D) fipronil (15g 2% G); E) fipronil (20g 2% G); F) chlorpyrifos (30g 0.125% G). In A,B,C, the quantities between parenthesis are of the com. formulation in 100 L of water. In D,E,F, are of granular insecticides per mound. One liter of the liqs. was used per nest. Demolition of the mounds were made 103 days after the application. The most efficient were abamectin and fipronil. Experiment II: Also 60 nests, with 6 treatments and 10 replications: A) fipronil (10g 2% G); B) fipronil (15g 2% G); C) bendiocarb (20g 0.1% G); D) bendiocarb (20g 0.5% G); E) **imidacloprid** (0.15g 70% G); F) **imidacloprid** (0.30g 70% G). In A,B,C,D, the quantities of granular insecticides are by nest. In E,F, of dispersible granule in 1 L

09886197

of water, by nest. The demolition of the nests was made 148 and 149 days after the application. The most efficient were fipronil and imidacloprid.

=>

=> logoff

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

LOGOFF? (Y)/N/HOLD:H

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

276.54

282.14

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-69.09

-69.09

SESSION WILL BE HELD FOR 60 MINUTES

STN INTERNATIONAL SESSION SUSPENDED AT 16:27:00 ON 21 JUL 2004

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssspta1202sxq

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 May 12 EXTEND option available in structure searching
NEWS 4 May 12 Polymer links for the POLYLINK command completed in REGISTRY
NEWS 5 May 27 New UPM (Update Code Maximum) field for more efficient patent SDIs in Caplus
NEWS 6 May 27 Caplus super roles and document types searchable in REGISTRY
NEWS 7 Jun 28 Additional enzyme-catalyzed reactions added to CASREACT
NEWS 8 Jun 28 ANTE, AQUALINE, BIOENG, CIVILENG, ENVIROENG, MECHENG, and WATER from CSA now available on STN(R)
NEWS 9 Jul 12 BEILSTEIN enhanced with new display and select options, resulting in a closer connection to BABS
NEWS 10 Jul 30 BEILSTEIN on STN workshop to be held August 24 in conjunction with the 228th ACS National Meeting
NEWS 11 AUG 02 IFIPAT/IFIUDB/IFICDB reloaded with new search and display fields
NEWS 12 AUG 02 Caplus and CA patent records enhanced with European and Japan Patent Office Classifications
NEWS 13 AUG 02 STN User Update to be held August 22 in conjunction with the 228th ACS National Meeting
NEWS 14 AUG 02 The Analysis Edition of STN Express with Discover! (Version 7.01 for Windows) now available

NEWS EXPRESS JULY 30 CURRENT WINDOWS VERSION IS V7.01, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),

L10 ANSWER 4 OF 4 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1995:187187 CAPLUS

DOCUMENT NUMBER: 122:25815

TITLE: **Imidacloprid** - a new systemic insecticide.

AUTHOR(S): Elbert, A.; Becker, B.; Hartwig, J.; Erdelen, C.

CORPORATE SOURCE: Geschäftsbereich Pflanzenschutz
Entwicklung/Insektizide, Bayer AG, Leverkusen, 5090,
GermanySOURCE: Pflanzenschutz-Nachrichten Bayer (German Edition)
(1991), 44(2), 113-36

CODEN: PNBAT; ISSN: 0340-1723

PUBLISHER: Bayer AG

DOCUMENT TYPE: Journal

LANGUAGE: German

AB The biol. profile of **Imidacloprid** (I) was defined on the basis of the results of exhaustive laboratory expts. and greenhouse trials. I is extremely effective against sucking **insects**, such as rice leafhoppers, aphids, thrips and mealybugs, and very effective against whitefly. It is also effective against some species of biting **insects**, such as paddy stem borers and Colorado beetle, but it has no effect on nematodes or spider mites. At comparatively high doses it kills adult **insects** and has ovicidal effects. I is a nicotinic acetylcholine receptor stimulator. Its mechanism of action differs from that of conventional insecticides. It therefore gives excellent control of all resistant populations investigated hitherto. I has a pos. temperature coefficient. After foliar application, it has a good residual action, it is highly photostable and it shows satisfactory resistance to rain. I is active after oral ingestion and by direct contact, but it is not active in the vapor phase. The LD95 after oral ingestion by *Myzus persicae* is .apprx.2 pg/aphid. After topical application it is .apprx.160 pg/aphid. It has not been possible to demonstrate recovery of injured aphids, or antifeeding effects. I has a faster action against aphids than oxydemeton-Me. After foliar application, I shows good translaminar and acropetal translocation, so it is also likely to provide effective control of pests with a furtive lifestyle, and protect the parts of the plant which regenerate after treatment. By virtue of its good contact action and powerful systemic action after uptake through the root system, I can be applied to soil and used as a seed dressing. It gives excellent control of pests such as onion maggots, *Diabrotica*, wire worms, **termites** and fire ants which live in the soil, and of **insects** such as aphids which live above ground level. It has a good residual action after application to the soil and when it is used as a seed dressing. The compatibility of I with plants is good after use as a seed dressing, as a soil treatment and after foliar application. By virtue of its biol. properties, I is likely to have a wide range of uses for controlling economically important pests of rice, cotton, cereals, maize, sugar beet, potatoes, vegetables, citrus fruit, pome and stone fruit and other crops.

=>

09886197

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssspta1202sxq

PASSWORD:

* * * * * RECONNECTED TO STN INTERNATIONAL * * * * *
SESSION RESUMED IN FILE 'CAPLUS' AT 16:25:08 ON 21 JUL 2004
FILE 'CAPLUS' ENTERED AT 16:25:08 ON 21 JUL 2004
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	238.44	244.04
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-59.54	-59.54

=> s isoptera
L12 570 ISOPTERA

=> s l12 and l2
L13 13 L12 AND L2

=> d l13 1-13 ibib hitstr abs

L13 ANSWER 1 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 2004:453202 CAPLUS
DOCUMENT NUMBER: 141:23526
TITLE: Novel pyrazole-based anthranilamide insecticides and
their preparation, compositions, and use
INVENTOR(S): Hughes, Kenneth Andrew; Lahm, George Philip; Selby,
Thomas Paul
PATENT ASSIGNEE(S): E.I. Du Pont De Nemours and Company, USA
SOURCE: PCT Int. Appl., 96 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004046129	A2	20040603	WO 2003-US36167	20031112
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,			


7/21/04

NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.:

US 2002-426693P P 20021115

GI



AB The invention provides title compds. I and their N-oxides and suitable salts [wherein: Y, V = N or CR_{4a}; W = N, CH, or CR₆; R₁ = H, (un)substituted alkyl, alkenyl, alkynyl or cycloalkyl, alkylcarbonyl, alkoxy, (di)alkylaminocarbonyl; R₂ = H, alkyl, alkenyl, alkynyl, cycloalkyl, alkoxy, (di)alkylamino, cycloalkylamino, alkoxy, alkylcarbonyl, or alkylcarbonyl; R₃ = H, G, (un)substituted alkyl, alkenyl, alkynyl or cycloalkyl; or NR₂R₃ = (un)substituted heterocyclic (N/O/S) ring; G = (un)substituted 5- or 6-membered non-aromatic carbo- or heterocyclic ring; R_{4a}, R_{4b} = H, various carbon and heteroat. substituents; R₅ = alk(en/yn)yl, various derivs. of OH, SH, and NH₂; R₆ = (halo)alk(en/yn)yl, OH and derivs. or thio analogs, halo, cyano, CO₂H, (di)alkylamino, (un)substituted Ph, PhCH₂, PhCO, PhO, etc.; n = 0-4]. The invention also pertains to compns. for controlling invertebrate pests, comprising a biol. effective amount of I, their N-oxides, or their agronomically or nonagronomically suitable salts, and at least one addnl. component selected from surfactants, solid diluents, and liquid diluents, and optionally further comprising an effective amount of at least one addnl. biol. active compound or agent. Also disclosed are methods for controlling invertebrate pests by contact of the pests or their environment with said compds. Eighteen compds. I were prepared and tested. For instance, 3-chloro-2-hydrazinopyridine was cyclocondensed with di-Et maleate to give 55% Et 1-(3-chloro-2-pyridinyl)-3-pyrazolidinone-5-carboxylate, which was oxidized to a dihydropyrazolone, saponified to an acid, cyclized with dichloroanthranilic acid to give a benzoxazinone, O-mesylated at the pyrazolone, and ring-opened with MeNH₂, to give invention compound II. In a test of larval *Plutella xylostella* on radish plants, II at 50 ppm (spray) reduced feeding damage by 80% or more. Compds. I were also effective against *Spodoptera frugiperda*, *Myzus persicae*, and *Empoasca fabae*.

L13 ANSWER 2 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:336622 CAPLUS

DOCUMENT NUMBER: 139:48626

TITLE: Effects of exposure duration on transfer of
nonrepellent termiticides among workers of *Coptotermes*
formosanus Shiraki (Isoptera:

09886197

Rhinotermitidae)
AUTHOR(S): Shelton, Thomas G.; Grace, J. Kenneth
CORPORATE SOURCE: Department of Plant & Environmental Protection
Sciences, University of Hawaii, Honolulu, HI,
96822-2271, USA
SOURCE: Journal of Economic Entomology (2003), 96(2), 456-460
CODEN: JEENAI; ISSN: 0022-0493
PUBLISHER: Entomological Society of America
DOCUMENT TYPE: Journal
LANGUAGE: English
AB The potential for transfer of nonrepellent termiticide toxicants between
workers of the Formosan subterranean termite, *Coptotermes formosanus*
Shiraki, was examined using two com. available pesticide formulations and a
simple donor-recipient model modified from current methods in the
literature. Pesticides used were **imidacloprid**, formulated as
Premise 75 WP, and fipronil, formulated as Termidor SC, in concns. of 1,
10, and 100 ppm (weight of active ingredient/weight of sand) in sand. A
significant increase was shown in recipient mortality over control
mortality when donor workers were treated with 100 ppm
imidacloprid or 100 ppm fipronil. Although all three colonies
studied were affected, one colony (colony 3) was affected to a
significantly greater extent than the other colonies. This effect was not
correlated with termite body size (dry mass). In a second study,
recipient mortality was evaluated after exposure of donors to 1 ppm
insecticide for 3, 6, 12, or 24 h. Recipient mortality indicated that
these exposures did not consistently lead to lethal transfer of the
insecticides.
REFERENCE COUNT: 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 3 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 2003:177184 CAPLUS
DOCUMENT NUMBER: 138:333176
TITLE: Effect of **imidacloprid** tree treatments on
the occurrence of formosan subterranean termites,
Coptotermes formosanus Shiraki (**Isoptera**:
Rhinotermitidae), in independent monitors
AUTHOR(S): Osbrink, Weste L. A.; Lax, Alan R.
CORPORATE SOURCE: Southern Regional Research Center, USDA-ARS, New
Orleans, LA, 70124, USA
SOURCE: Journal of Economic Entomology (2003), 96(1), 117-125
CODEN: JEENAI; ISSN: 0022-0493
PUBLISHER: Entomological Society of America
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Periodic sampling of 87 independent monitors, initially active with the
Formosan subterranean termite, *Coptotermes formosanus* Shiraki, was
conducted. Monitors, located in eight sectors adjacent to seven
buildings, were various distances (1-46 m) from 57 trees treated with 0.1%
imidacloprid foam. Termites collected from six of the eight
sectors showed latent mortality attributed to **imidacloprid**
intoxication at all monitor-tree distances. Approx. 6 mo after treatment,
termite populations had recovered in these sectors. Another sector showed
termite population suppression for ≈15 mo, followed by recovery.
Imidacloprid tree treatments did not control *C. formosanus*
populations in independent monitors adjacent to the treatments.
REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

7/21/04

09886197

L13 ANSWER 4 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:777603 CAPLUS
DOCUMENT NUMBER: 137:274431
TITLE: Insecticide compositions containing amino acids
INVENTOR(S): Sandeman, Richard Mark; Chandler, David Spencer;
Duncan, Ann Maree; Hay, Phillip Maxwell
PATENT ASSIGNEE(S): Nufarm Limited, Australia; La Trobe University
SOURCE: PCT Int. Appl., 62 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002078448	A1	20021010	WO 2002-AU389	20020328
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1385379	A1	20040204	EP 2002-712624	20020328
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
PRIORITY APPLN. INFO.:			AU 2001-4069	A 20010329
			WO 2002-AU389	W 20020328

OTHER SOURCE(S): MARPAT 137:274431

AB Insecticides of formula R3N(R2)AC(R1)(:O) and the agriculturally acceptable salts thereof (R1 = OR5 wherein R5 = H, (un)substituted alkyl, (un)substituted aryl, (un)substituted cycloalkyl, (un)substituted heterocyclic; NR6OH wherein R6 = H, (un)substituted alkyl, (un)substituted aryl, (un)substituted carbocyclic; NR7R8 wherein R7 and R8 = H, (un)substituted alkyl, (un)substituted aryl and carbocyclic; and wherein R1 is linked to R2 to form a diradical bridging group; R2 and R3 = H, (un)substituted alkyl, (un)substituted carbocyclic, (un)substituted aryl, and (un)substituted acyl; and A = diradical linking group, which has a mol. weight of preferably less than 200 and more preferably less than 100) are used to control insect species selected from the orders Lepidoptera, Hemiptera, Orthoptera, Coleoptera, Psocoptera, **Isoptera**, Thysanoptera and Homoptera on cotton.

REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 5 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:720924 CAPLUS
DOCUMENT NUMBER: 135:340463
TITLE: Chemical prevention of colony foundation by
Cryptotermes brevis (**Isoptera**:
Kalotermitidae) in attic modules
AUTHOR(S): Scheffrahn, Rudolf H.; Busey, Philip; Edwards, Jeffrey
K.; Krecek, Jan; Maharajh, Boudanath; Su, Nan-Yao
CORPORATE SOURCE: Ft. Lauderdale Research and Education Center,
University of Florida, Fort Lauderdale, FL, 33314, USA

09886197

SOURCE: Journal of Economic Entomology (2001), 94(4), 915-919
CODEN: JEENAI; ISSN: 0022-0493
PUBLISHER: Entomological Society of America
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Disodium octaborate tetrahydrate (DOT) dust, DOT aqueous solution, **imidacloprid** dust, and amorphous silica gel dust with synergized 1% pyrethrins were applied on wood surfaces to simulated attic modules. Modules (30 by 30 cm) with and without fiberglass insulation were exposed to dispersal flights of *Cryptotermes brevis* (Walker) in May and June of 1998 and 1999. Six months after flights, modules were disassembled and inspected for nuptial chamber location and contents. During both years, air and water control treatments contained 22.2 ± 9.94 (mean \pm SD) nuptial chambers, 7.5 ± 5.7 live imagos, and 2.0 ± 1.4 chambers with brood. This survivorship indicated that the attic modules performed well as a colonizing platform for *C. brevis*. *C. brevis* dealates preferred constructing nuptial chambers in the crevices at the bases or tops of the modules instead of internal crevices. Modules treated in 1998 and 1999 with DOT or silica dusts contained no live termites, whereas zero of five modules treated with **imidacloprid** dust in 1998 and two of 20 modules treated with **imidacloprid** dust in 1999 contained single live incipient colonies. In 1998, 15% DOT solution, applied as a postconstruction treatment, yielded significantly fewer chambers and live termites than controls, but was not as effective as dusts in preventing successful colonization. In 1999, the DOT solution, applied as a construction-phase treatment, was equally as effective in preventing colonization as the dust treatments during that year. Results indicate that dust formulations of DOT, silica gel, and **imidacloprid** can be used to prevent drywood termite colonization in existing building voids and attics. Where the entire wood framing is exposed to treatment, such as during building construction, aqueous DOT solution can be equally effective as dusts in preventing colonization by *C. brevis*.

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 6 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:336305 CAPLUS

DOCUMENT NUMBER: 135:1645

TITLE: Effects of sublethal exposure to **imidacloprid** on subsequent behavior of subterranean termite *Reticulitermes virginicus* (**Isoptera**: Rhinotermitidae)

AUTHOR(S): Thorne, Barbara L.; Breisch, Nancy L.

CORPORATE SOURCE: Department of Entomology, University of Maryland, College Park, MD, 20742, USA

SOURCE: Journal of Economic Entomology (2001), 94(2), 492-498
CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Expts. were conducted to determine whether subterranean termites, *Reticulitermes virginicus* (Banks), previously exposed to sublethal doses of **imidacloprid** (Premise), and allowed to recover for 1 wk, demonstrated behavioral aversion to a subsequent exposure. Worker termites experiencing a previous sublethal but debilitating exposure to **imidacloprid**-treated sand (either 10 or 100 ppm for 4 h) showed no apparent aversion to a second encounter with **imidacloprid**-treated sand under conditions of this experiment If these laboratory results hold

in the field and termites traveling through a zone of soil treated with **imidacloprid** are impaired but subsequently recover, they will be just as likely as their naive nestmates to reenter the treated area if their travels take them through the nonrepellent application a second time. Thus, a sublethal exposure to **imidacloprid** can affect termite tunneling behavior. Many worker termites that received an initial 4-h exposure to 100 ppm **imidacloprid**-treated sand died, but those that survived tunneled significantly less than did their naive nestmates, as did some termites exposed to 10 ppm **imidacloprid**.

REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 7 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:666543 CAPLUS

DOCUMENT NUMBER: 133:248390

TITLE: Synergistic insecticidal compositions containing a neuronal sodium channel antagonist and another insecticide

INVENTOR(S): Treacy, Michael Frank; Borysewicz, Raymond Frank; Schwinghammer, Kurt Allen; Rensner, Paul Erich; Oloumi-Sadeghi, Hassan

PATENT ASSIGNEE(S): American Cyanamid Company, USA

SOURCE: PCT Int. Appl., 30 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000054591	A2	20000921	WO 2000-US5879	20000307
WO 2000054591	A3	20010118		
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
AU 2000036175	A5	20001004	AU 2000-36175	20000307
AU 765767	B2	20030925		
BR 2000008930	A	20011218	BR 2000-8930	20000307
EP 1198170	A2	20020424	EP 2000-914839	20000307
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL			
NZ 514000	A	20030429	NZ 2000-514000	20000307
JP 2003517455	T2	20030527	JP 2000-604685	20000307
US 6479543	B1	20021112	US 2000-521987	20000309
ZA 2001007484	A	20021201	ZA 2001-7484	20010911
US 2002177597	A1	20021128	US 2002-145784	20020516

PRIORITY APPLN. INFO.:

US 1999-124306P	P	19990312
US 1999-158201P	P	19991007
WO 2000-US5879	W	20000307
US 2000-521987	A3	20000309

OTHER SOURCE(S): MARPAT 133:248390

GI



AB A synergistic insecticidal composition comprises a neuronal sodium channel antagonist such as I (X, Y, Z = H, halo, OH, CN, NO₂, alkyl, etc.; W = O or S; m, p, q = 1, 2, 3, 4, or 5; n = 0, 1, or 2; R, R₁, R₂, R₃ = alkyl) in combination with one or more pyrethroids, pyrethroid-type compds., recombinant nucleopolyhedroviruses expressing an insect toxin, organophosphates, carbamates, formamidines, macrocyclic lactones, amidinohydrazones, GABA antagonists and acetylcholine receptor ligands.

L13 ANSWER 8 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:573349 CAPLUS

DOCUMENT NUMBER: 133:248356

TITLE: Feeding inhibition and mortality in *Reticulitermes flavipes* (Isoptera: Rhinotermitidae) after exposure to **imidacloprid**-treated soils

AUTHOR(S): Ramakrishnan, Rathna; Suiter, Daniel R.; Nakatsu, Cindy H.; Bennett, Gary W.

CORPORATE SOURCE: Center for Urban & Industrial Pest Management, Department of Entomology, Purdue University, West Lafayette, IN, 47907, USA

SOURCE: Journal of Economic Entomology (2000), 93(2), 422-428
CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Feeding inhibition and mortality of *Reticulitermes flavipes* (Kollar) exposed to sand, sandy loam, loam, and silty clay loam soils treated with several concns. of **imidacloprid** were studied using bioassay techniques under laboratory conditions. Termite workers stopped feeding after exposure to treated soils. Differences in feeding reduction varied among the soil types. Based on the magnitude of the F-statistics, the effect of **imidacloprid** on the reduction of termite feeding was greatest in sand followed by sandy loam, loam, and silty clay loam soils. Soil properties such as organic matter content, silt and clay proportions, pH, and cation exchange capacity were suggested to affect the bioavailability of **imidacloprid**. Similar soil effects on mortality were observed in termites continuously exposed to treated soil for 21 days. In 3 of 4 soils tested, susceptibility to **imidacloprid** was not affected by the source of the termites tested.

REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 9 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:470450 CAPLUS

DOCUMENT NUMBER: 133:90469

TITLE: Adhesive composition containing insecticides, preservatives, termite repellents and bactericides for

INVENTOR(S): lignocellulosic material and it complex
Jaesch, Tohmas; Fushiki, Kiyoyuki; Saito, Takanobu;
Katsusawa, Yoshinaga
PATENT ASSIGNEE(S): Bayer A.-G., Germany; Ohshika Shinko K. K.; Chemiholz
K. K.
SOURCE: Jpn. Kokai Tokkyo Koho, 12 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000192001	A2	20000711	JP 1998-376942	19981228
KR 2000048138	A	20000725	KR 1999-57526	19991214
EP 1018413	A1	20000712	EP 1999-124843	19991215
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
AU 9965409	A1	20010628	AU 1999-65409	19991222
NZ 502074	A	20020301	NZ 1999-502074	19991223
NO 9906479	A	20000629	NO 1999-6479	19991227
US 2001027217	A1	20011004	US 1999-472589	19991227
BR 9907435	A	20010320	BR 1999-7435	19991228
PRIORITY APPLN. INFO.:		JP 1998-376942	A	19981228

AB The composition, for preparation of wood products (e.g., plywood), comprises an adhesive, an organic phenolic composition, an insecticide, a preservative, a termite repellent and a bactericide. Thus, a composition was made from Oshika Resin PWP 60 containing a solution of **imidacloprid** 3, IPBC 20 and 2-phenylphenol 15, and a solvent 62%.

L13 ANSWER 10 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:797191 CAPLUS

DOCUMENT NUMBER: 132:60446

TITLE: **Imidacloprid**-enhanced Reticulitermes flavipes (**Isoptera**: Rhinotermitidae) susceptibility to the entomopathogen Metarhizium anisopliae

AUTHOR(S): Ramakrishnan, Rathna; Suiter, Daniel R.; Nakatsu, Cindy H.; Humber, Richard A.; Bennett, Gary W.
CORPORATE SOURCE: Center for Urban & Industrial Pest Management, Department of Entomology, Purdue University, West Lafayette, IN, 47907, USA

SOURCE: Journal of Economic Entomology (1999), 92(5), 1125-1132
CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The effects of **imidacloprid** and the entomopathogen Metarhizium anisopliae (Metsch.) Sorokin on the eastern subterranean termite, Reticulitermes flavipes (Kollar), were evaluated in a 4 + 3 factorial experiment in both sterile and nonsterile loam soil. Termites were not susceptible to M. anisopliae when assays were conducted in nonsterile soil, but were highly susceptible in sterile soil. Termite mortality after 21 days of continuous exposure to 104 conidia per g soil was 0 and 41.6% in nonsterile and sterile soil, resp. Termites were significantly more susceptible to sterile soil containing 107 conidia per g than to the same soil containing 104 conidia per g. In continuous exposure assays, termites

09886197

were highly susceptible to **imidacloprid**-treated (5,10, and 20 ppm) nonsterile and sterile soil containing no exptl. introduced *M. anisopliae*. Exposure of termites to **imidacloprid** enhanced their susceptibility to introduced *M. anisopliae* in nonsterile and sterile soil. Native entomopathogens recovered from termites exposed to **imidacloprid**-treated, nonsterile soil (i.e., no introduced *M. anisopliae*) included *Conidiobolus coronatus* (Constantin) Batko, *Cunninghamella echinulata* Thaxter, *Fusarium* spp., *Aspergillus* spp., and a naturally occurring strain of *M. anisopliae* variety *majus*.

REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 11 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1998:631799 CAPLUS

DOCUMENT NUMBER: 129:246360

TITLE: Ant-repellent thermoplastic foam molding compositions containing chloropyridines for thermal insulators

INVENTOR(S): Toyonaga, Yoshihiro; kanzaki, Masahiro

PATENT ASSIGNEE(S): Shinto Paint Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 10259270	A2	19980929	JP 1997-86053	19970319

PRIORITY APPLN. INFO.: JP 1997-86053 19970319

AB Title compns. contain acetamiprid (I) or **imidacloprid** as an ant repellent. Thus, prefoamed polystyrene was coated with a solution containing I and adhesive, mixed with uncoated polystyrene, and heated to give a molding containing 0.2% I. The molding showed bending strength 3.8 kg/cm2 and no damage by termite for ≥ 21 days.

L13 ANSWER 12 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1998:631792 CAPLUS

DOCUMENT NUMBER: 129:317341

TITLE: Termite-repellent polyurethane foam molding compositions containing chloropyridines for thermal insulators

INVENTOR(S): Toyonaga, Yoshihiro; Okuta, Kazuo

PATENT ASSIGNEE(S): Shinto Paint Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 4 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 10259263	A2	19980929	JP 1997-86055	19970319

PRIORITY APPLN. INFO.: JP 1997-86055 19970319

AB Title compns. contain acetamiprid (I) or **imidacloprid** as termite repellents. Thus, HCFC-containing Polyol GB, I, and polyphenyl-type polyisocyanates were molded to give 0.1% I-containing polyurethane foams showing bending strength 4.5 kg/cm2 and no damage by termite for

≥21 days.

L13 ANSWER 13 OF 13 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1995:881641 CAPLUS

DOCUMENT NUMBER: 123:278677

TITLE: Field tests for control of the mound-building termite
Cornitermes cumulans (Kollar, 1832) (**Isoptera**
, **Termitidae**)AUTHOR(S): Mariconi, F.A.M.; Galan, V.B.; Rocha, M.T.; Maule,
R.F.; Passos, H.R.; Silva, R.A.A.

CORPORATE SOURCE: ESALQ, USP, Piracicaba, 13418-900, Brazil

SOURCE: Scientia Agricola (Piracicaba, Brazil) (1994), 51(3),
505-8

CODEN: SGRIEF; ISSN: 0103-9016

PUBLISHER: Universidade de Sao Paulo, Campus de Piracicaba

DOCUMENT TYPE: Journal

LANGUAGE: Portuguese

AB Two field tests were carried out to evaluate the performance of several pesticides for the control of the mound termite pest in pastures. Experiment I: 60 mounds were selected and measured outside. There were 6 treatments with 10 replications: A) abamectin (50 cm³ 1.8% EC); B) silafluofen (200 cm³ 80% EC); C) silafluofen (400 cm³ 80% EC); D) fipronil (15g 2% G); E) fipronil (20g 2% G); F) chlorpyrifos (30g 0.125% G). In A,B,C, the quantities between parenthesis are of the com. formulation in 100 L of water. In D,E,F, are of granular insecticides per mound. One liter of the liqs. was used per nest. Demolition of the mounds were made 103 days after the application. The most efficient were abamectin and fipronil. Experiment II: Also 60 nests, with 6 treatments and 10 replications: A) fipronil (10g 2% G); B) fipronil (15g 2% G); C) bendiocarb (20g 0.1% G); D) bendiocarb (20g 0.5% G); E) **imidacloprid** (0.15g 70% G); F) **imidacloprid** (0.30g 70% G). In A,B,C,D, the quantities of granular insecticides are by nest. In E,F, of dispersible granule in 1 L of water, by nest. The demolition of the nests was made 148 and 149 days after the application. The most efficient were fipronil and **imidacloprid**.

09886197

=> d his

(FILE 'HOME' ENTERED AT 16:06:53 ON 21 JUL 2004)

FILE 'STNGUIDE' ENTERED AT 16:07:06 ON 21 JUL 2004

FILE 'REGISTRY' ENTERED AT 16:08:17 ON 21 JUL 2004

L1 0 S IMIDALCLOPRID

FILE 'CAPLUS' ENTERED AT 16:09:11 ON 21 JUL 2004

L2 1376 S IMIDACLOPRID
L3 22 S L2 AND TERMITES
L4 25 S L2 AND WOOD
L5 6 S L3 AND L4
L6 2129 S TERMITES
L7 29825 S INSECTS
L8 274 S L6 AND L7
L9 47 S L8 AND SPECIES
L10 4 S L2 AND L8

=> s l9 and l2

L11 1 L9 AND L2

=> d l11 ibib hitstr abs

L11 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1995:187187 CAPLUS

DOCUMENT NUMBER: 122:25815

TITLE: Imidacloprid - a new systemic insecticide.

AUTHOR(S): Elbert, A.; Becker, B.; Hartwig, J.; Erdelen, C.

CORPORATE SOURCE: Geschäftsbereich Pflanzenschutz
Entwicklung/Insektizide, Bayer AG, Leverkusen, 5090,
Germany

SOURCE: Pflanzenschutz-Nachrichten Bayer (German Edition)
(1991), 44(2), 113-36
CODEN: PNBYPAT; ISSN: 0340-1723

PUBLISHER: Bayer AG

DOCUMENT TYPE: Journal

LANGUAGE: German

AB The biol. profile of **Imidacloprid** (I) was defined on the basis of the results of exhaustive laboratory expts. and greenhouse trials. I is extremely effective against sucking **insects**, such as rice leafhoppers, aphids, thrips and mealybugs, and very effective against whitefly. It is also effective against some **species** of biting **insects**, such as paddy stem borers and Colorado beetle, but it has no effect on nematodes or spider mites. At comparatively high doses it kills adult **insects** and has ovicidal effects. I is a nicotinic acetylcholine receptor stimulator. Its mechanism of action differs from that of conventional insecticides. It therefore gives excellent control of all resistant populations investigated hitherto. I has a pos. temperature coefficient. After foliar application, it has a good residual action, it is highly photostable and it shows satisfactory resistance to rain. I is active after oral ingestion and by direct contact, but it is not active in the vapor phase. The LD95 after oral ingestion by *Myzus persicae* is .apprx.2 pg/aphid. After topical application it is .apprx.160 pg/aphid. It has not been possible to demonstrate recovery of injured aphids, or antifeeding effects. I has a faster action against aphids than oxydemeton-Me. After foliar application, I shows good translaminar and

acropetal translocation, so it is also likely to provide effective control of pests with a furtive lifestyle, and protect the parts of the plant which regenerate after treatment. By virtue of its good contact action and powerful systemic action after uptake through the root system, I can be applied to soil and used as a seed dressing. It gives excellent control of pests such as onion maggots, Diabrotica, wire worms, **termites** and fire ants which live in the soil, and of **insects** such as aphids which live above ground level. It has a good residual action after application to the soil and when it is used as a seed dressing. The compatibility of I with plants is good after use as a seed dressing, as a soil treatment and after foliar application. By virtue of its biol. properties, I is likely to have a wide range of uses for controlling economically important pests of rice, cotton, cereals, maize, sugar beet, potatoes, vegetables, citrus fruit, pome and stone fruit and other crops.

=>

09886197

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal202sxq

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

*Terminates
order Isoptera*

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 May 12 EXTEND option available in structure searching
NEWS 4 May 12 Polymer links for the POLYLINK command completed in REGISTRY
NEWS 5 May 27 New UPM (Update Code Maximum) field for more efficient patent
SDIs in Caplus
NEWS 6 May 27 Caplus super roles and document types searchable in REGISTRY
NEWS 7 Jun 22 STN Patent Forums to be held July 19-22, 2004
NEWS 8 Jun 28 Additional enzyme-catalyzed reactions added to CASREACT
NEWS 9 Jun 28 ANTE, AQUALINE, BIOENG, CIVILENG, ENVIROENG, MECHENG,
and WATER from CSA now available on STN(R)
NEWS 10 Jul 12 BEILSTEIN enhanced with new display and select options,
resulting in a closer connection to BABS

NEWS EXPRESS MARCH 31 CURRENT WINDOWS VERSION IS V7.00A, CURRENT
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 26 APRIL 2004
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 16:06:53 ON 21 JUL 2004

=> FIL STNGUIDE

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'STNGUIDE' ENTERED AT 16:07:06 ON 21 JUL 2004

USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT

7/21/04

09886197

COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE
AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: Jul 16, 2004 (20040716/UP).

```
=> file reg
COST IN U.S. DOLLARS          SINCE FILE      TOTAL
                               ENTRY      SESSION
FULL ESTIMATED COST          0.12        0.33
```

FILE 'REGISTRY' ENTERED AT 16:08:17 ON 21 JUL 2004
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2004 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.

STRUCTURE FILE UPDATES: 20 JUL 2004 HIGHEST RN 713489-00-0
DICTIONARY FILE UPDATES: 20 JUL 2004 HIGHEST RN 713489-00-0

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 6, 2004

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more
information enter HELP PROP at an arrow prompt in the file or refer
to the file summary sheet on the web at:
<http://www.cas.org/ONLINE/DBSS/registryss.html>

```
=> s imidalcloprid
L1          0 IMIDALCLOPRID
```

```
=> file caplus
COST IN U.S. DOLLARS          SINCE FILE      TOTAL
                               ENTRY      SESSION
FULL ESTIMATED COST          5.27        5.60
```

FILE 'CAPLUS' ENTERED AT 16:09:11 ON 21 JUL 2004
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is
held by the publishers listed in the PUBLISHER (PB) field (available
for records published or updated in Chemical Abstracts after December
26, 1996), unless otherwise indicated in the original publications.
The CA Lexicon is the copyrighted intellectual property of the
American Chemical Society and is provided to assist you in searching
databases on STN. Any dissemination, distribution, copying, or storing
of this information, without the prior written consent of CAS, is
strictly prohibited.

FILE COVERS 1907 - 21 Jul 2004 VOL 141 ISS 4

7/21/04

09886197

FILE LAST UPDATED: 20 Jul 2004 (20040720/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s imidacloprid
L2 1376 IMIDACLOPRID

=> s l2 and termites
2129 TERMITES
L3 22 L2 AND TERMITES

=> s l2 and wood
145525 WOOD
L4 25 L2 AND WOOD

=> s l3 and l4
L5 6 L3 AND L4

=> d l5 1-6 ibib hitstr abs

L5 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 2002:695680 CAPLUS
DOCUMENT NUMBER: 137:228094
TITLE: Termiticidal baits for eliminating termite colonies
INVENTOR(S): Brode, Philip Frederick, III; Garrett, Garry Steven;
Laughlin, Leo Timothy; Matthews, Randall Stryker;
Barker, Dale Edwin; Kinne, Daniel James; Miller,
Christopher Miles; Probst, Timothy Robert; McKibben,
Gary Eugene
PATENT ASSIGNEE(S): The Procter & Gamble Company, USA
SOURCE: PCT Int. Appl., 61 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002069704	A2	20020912	WO 2002-US6200	20020301
WO 2002069704	A3	20021114		
WO 2002069704	C1	20031231		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 2002172658	A1	20021121	US 2001-799184	20010305
US 6716421	B2	20040406		
US 2003017187	A1	20030123	US 2002-172855	20020617
US 2003124166	A1	20030703	US 2002-173527	20020617
US 2003124164	A1	20030703	US 2002-268356	20021010
WO 2003105580	A1	20031224	WO 2003-US17713	20030605

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

WO 2003106395 A1 20031224 WO 2003-US17714 20030605

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

WO 2004032625 A2 20040422 WO 2003-US32092 20031007


W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.:

US 2001-799184 A 20010305
US 2002-172855 A 20020617
US 2002-173527 A 20020617
US 2002-268356 A 20021010

OTHER SOURCE(S): MARPAT 137:228094
GI



AB This invention relates to devices, kits, and methods for eliminating termite colonies. The kits, devices, and methods employ a termiticidal bait matrix contain (a) a termiticide (I, X = nil, a hydrocarbon group, O or NR₈, R₉ where R₈ and R₉ are H or a hydrocarbon group; X₁ = CH, a carbon atom or a heteroatom; R, R₁, R₂, R₃ = H or OH and if R₄ and R₅ are O and R₆ and R₇ are H then R, R₁, R₂ and R₃ may be C₁-6; R₄ and R₅ are H, O or N; R₉ and R₁₀ are nil, C₁-6, and amides) selected such that the termiticide causes death to about 50 to about 100% of **termites** within about

24 to about 84 days after the **termites** begin to ingest the termiticide or the bait matrix comprising the termiticide, (b) a cellulose containing material, and (c) water. The termiticidal bait matrix can be used in a bait station installed in the ground. The kits are suitable to be used by consumers in their homes.

L5 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:547214 CAPLUS

DOCUMENT NUMBER: 137:105178

TITLE: Termiticides containing 2-pyridinethiol-1-oxide salts and **wood** and polymers containing the termiticides

INVENTOR(S): Nishimoto, Koichi; Sato, Toshio; Suga, Mamoru

PATENT ASSIGNEE(S): Yoshitomi Fine Chemicals Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002205906	A2	20020723	JP 2001-337124	20010926
PRIORITY APPLN. INFO.:			JP 2000-381082	A 20001108

AB The termiticides, which are effective on **termites**, bark beetles, etc., and environmentally safe, contain (a) ≥ 1 selected from Cu, Zn, and Na salts of 2-pyridine-1-oxide and optionally (b) ≥ 1 selected from pyrethroids, nicotinoids, organophosphorus compds., isocyanuric acid compds., carbamates, acetamiprid, and inorg. boric acid compds. **Wood** and polymers containing the termiticides are also claimed. A **wood** block was coated with DMSO solution containing Cu pyrithione and **imidacloprid** and dried at room temperature for ≥ 20 days. The **wood** block.

L5 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:720924 CAPLUS

DOCUMENT NUMBER: 135:340463

TITLE: Chemical prevention of colony foundation by *Cryptotermes brevis* (Isoptera: Kalotermitidae) in attic modules

AUTHOR(S): Scheffrahn, Rudolf H.; Busey, Philip; Edwards, Jeffrey K.; Krecek, Jan; Maharajh, Boudanath; Su, Nan-Yao

CORPORATE SOURCE: Ft. Lauderdale Research and Education Center, University of Florida, Fort Lauderdale, FL, 33314, USA

SOURCE: Journal of Economic Entomology (2001), 94(4), 915-919

CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Disodium octaborate tetrahydrate (DOT) dust, DOT aqueous solution, **imidacloprid** dust, and amorphous silica gel dust with synergized 1% pyrethrins were applied on **wood** surfaces to simulated attic modules. Modules (30 by 30 cm) with and without fiberglass insulation were exposed to dispersal flights of *Cryptotermes brevis* (Walker) in May and June of 1998 and 1999. Six months after flights, modules were disassembled and inspected for nuptial chamber location and contents. During both years, air and water control treatments contained 22.2 ± 9.94 (mean \pm SD) nuptial chambers, 7.5 ± 5.7 live imagos, and 2.0

± 1.4 chambers with brood. This survivorship indicated that the attic modules performed well as a colonizing platform for *C. brevis*. *C. brevis* dealates preferred constructing nuptial chambers in the crevices at the bases or tops of the modules instead of internal crevices. Modules treated in 1998 and 1999 with DOT or silica dusts contained no live **termites**, whereas zero of five modules treated with **imidacloprid** dust in 1998 and two of 20 modules treated with **imidacloprid** dust in 1999 contained single live incipient colonies. In 1998, 15% DOT solution, applied as a postconstruction treatment, yielded significantly fewer chambers and live **termites** than controls, but was not as effective as dusts in preventing successful colonization. In 1999, the DOT solution, applied as a construction-phase treatment, was equally as effective in preventing colonization as the dust treatments during that year. Results indicate that dust formulations of DOT, silica gel, and **imidacloprid** can be used to prevent drywood termite colonization in existing building voids and attics. Where the entire **wood** framing is exposed to treatment, such as during building construction, aqueous DOT solution can be equally effective as dusts

in

preventing colonization by *C. brevis*.

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:467897 CAPLUS

DOCUMENT NUMBER: 133:85609

TITLE: Termiticidal baits comprising nonhygroscopic agents in hygroscopic containers

INVENTOR(S): Minakawa, Fumiyasu; Uchida, Yuki

PATENT ASSIGNEE(S): Yuko Chemical Industries Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000189031	A2	20000711	JP 1998-369335	19981225
PRIORITY APPLN. INFO.:			JP 1998-369335	19981225

AB A nonhygroscopic agent for controlling **termites** (e.g. diflubenzuron) is housed in a container which is made of an edible hygroscopic material (cellulosic cloth, polyvinyl alc. film). Thus, sulfluramid 0.001 and pine oil 1% (attractant) were dissolved in polyethylene glycol, and the solution was made to soaked into **wood** flour at a 25/100 weight ratio. The agent was heat sealed in an envelope (15 + 7 cm) made of nonwoven fabric of cellulose fibers with 1% by weight added pine oil to obtain a bait with satisfactory attractiveness to *Reticulitermes*.

L5 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:125767 CAPLUS

DOCUMENT NUMBER: 130:178773

TITLE: Composition for the control of **wood** -destroying insects, especially **termites**

INVENTOR(S): Anderson, John-phillip-evans; Keuken, Oliver

PATENT ASSIGNEE(S): Bayer A.-G., Germany

SOURCE: Eur. Pat. Appl., 21 pp.

09886197

DOCUMENT TYPE: CODEN: EPXXDW
LANGUAGE: Patent
FAMILY ACC. NUM. COUNT: 1 German
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 896791	A2	19990217	EP 1998-114187	19980729
EP 896791	A3	20000112		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
DE 19734665	A1	19990218	DE 1997-19734665	19970811
TW 505500	B	20021011	TW 1998-87112592	19980731
US 6264968	B1	20010724	US 1998-128818	19980804
ZA 9807118	A	19990209	ZA 1998-7118	19980807
JP 11124302	A2	19990511	JP 1998-234861	19980807
AU 9879895	A1	19990218	AU 1998-79895	19980811
AU 768390	B2	20031211		
BR 9803138	A	19991221	BR 1998-3138	19980811

PRIORITY APPLN. INFO.: DE 1997-19734665 A 19970811
AB The title compns. (no examples) comprise an insecticide, preferably **imidacloprid**, incorporated into an organic natural and/or synthetic carrier. Optional ingredients are insect attractants and microbicides.

L5 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1993:54353 CAPLUS

DOCUMENT NUMBER: 118:54353

TITLE: Imidozolidine derivatives and related compounds as industrial insecticides and wood preservatives

INVENTOR(S): Tsuboi, Shinichi; Sone, Shinzaburo; Obinata, Toru; Exner, Otto; Schwamborn, Michael

PATENT ASSIGNEE(S): Nihon Bayer Agrochem K. K., Japan

SOURCE: Eur. Pat. Appl., 15 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 511541	A1	19921104	EP 1992-106384	19920414
EP 511541	B1	19960904		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
JP 05032505	A2	19930209	JP 1991-350751	19911212
JP 3162450	B2	20010425		
JP 2001031511	A2	20010206	JP 2000-233512	19911212
AU 9213908	A1	19921029	AU 1992-13908	19920330
AU 645672	B2	19940120		
AT 142077	E	19960915	AT 1992-106384	19920414
ES 2090400	T3	19961016	ES 1992-106384	19920414
BR 9201534	A	19921201	BR 1992-1534	19920427
US 6323224	B1	20011127	US 1995-543351	19951016
US 2001051643	A1	20011213	US 2001-886197	20010621


PRIORITY APPLN. INFO.: JP 1991-125172 A 19910427
JP 1991-350751 A 19911212
US 1992-872279 B1 19920422

09886197

US 1995-543351 A3 19951016

OTHER SOURCE(S):
GI

MARPAT 118:54353



AB The imidazolidine derivs. and related compds. I (X = NH, S; Y = CH, N; Z = 2-chloro-5-pyridyl, 2-chloro-5-thiazolyl; R = H, Me; n = 0, 1) are industrial insecticides and wood preservatives. Wood impregnated with 0.32 ppm imidacloprid was lethal to termites (Coptotermes formosanus) for ≥3 wk.

=> d his

(FILE 'HOME' ENTERED AT 16:06:53 ON 21 JUL 2004)

FILE 'STNGUIDE' ENTERED AT 16:07:06 ON 21 JUL 2004

FILE 'REGISTRY' ENTERED AT 16:08:17 ON 21 JUL 2004

L1 0 S IMIDALCLOPRID

FILE 'CAPLUS' ENTERED AT 16:09:11 ON 21 JUL 2004

L2 1376 S IMIDACLOPRID

L3 22 S L2 AND TERMITES

L4 25 S L2 AND WOOD

L5 6 S L3 AND L4

=> d l3 1-22 ibib hitstr abs

L3 ANSWER 1 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:892507 CAPLUS

DOCUMENT NUMBER: 139:360411

TITLE: Naphthalenic compounds as termite bait toxicants

INVENTOR(S): Rojas, Maria Guadalupe; Morales-Ramos, Juan A.; Green, Frederick, III

PATENT ASSIGNEE(S): The United States of America, as Represented by the Secretary of Agriculture, USA

SOURCE: PCT Int. Appl., 17 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003092375	A2	20031113	WO 2003-US13457	20030430
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,				

7/21/04

LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH,
 PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ,
 UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU,
 TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
 CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
 NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
 GW, ML, MR, NE, SN, TD, TG

US 6691453 B1 20040217 US 2002-135224 20020430

PRIORITY APPLN. INFO.: US 2002-135224 A 20020430

AB A matrix suitable to be used as baits and attractants for **termites** comprises cellulose, naphthalenic compds., water, and potentially other termite-preferred nutrients. Methods of monitoring the presence of **termites** using such matrixes and methods of controlling **termites** using such matrixes to deliver termite toxicants (e.g., streptomycin sulfate or **imidacloprid**) are also provided. Thus, N-hydroxynaphthalimide sodium salt (I) was incorporated into a bait matrix containing lecithin, ergosterol, EtOH, yeast hydrolyzate, and cellulose. I at 500 ppm was sufficient to induce mortality of Formosan subterranean termite (*Coptotermes formosanus*) within .apprx.2 mo without any repellency to **termites**.

L3 ANSWER 2 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:690211 CAPLUS

DOCUMENT NUMBER: 139:334278

TITLE: Evaluation of chemical control measures for **termites** in maize

AUTHOR(S): Riekert, H. F.; Van den Berg, J.

CORPORATE SOURCE: ARC-Grain Crops Institute, Potchefstroom, 2520, S. Afr.

SOURCE: South African Journal of Plant and Soil (2003), 20(1), 1-5

CODEN: SAJSEV; ISSN: 0257-1862

PUBLISHER: Forum Press International

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Field trials were conducted from the 1994/95 to 2000/2001 growing seasons to evaluate various insecticides for preventative and corrective control of the fungus-growing **termites**, *Microtermes* sp., *Odontotermes* sp. and *Allodotermes* sp. in maize. The incidence of lodged maize plants was used as criteria for insecticide efficacy. Carbofuran GR, **imidacloprid** WS, chlorpyrifos GR and fipronil GR were evaluated as preventative treatments. Corrective treatments in the form of spray applications of the systemic insecticides carbosulfan EC, benfuracarb EC and **imidacloprid** SL were also evaluated. Treatments were applied to the basal 25 cm of maize stems and to the soil surface surrounding plants. **Imidacloprid** spray applications generally provided good control of **termites**. The optimum plant growth stage for **imidacloprid** application was during the pre-flowering stage, 6 to 10 wk after plant emergence. Pre-flowering applications were usually more effective in limiting damage than post-flowering applications. The granular insecticide, fipronil, showed promise for termite control. Chlorpyrifos granules, applied as a side dressing four weeks after plant emergence, significantly reduced lodging. Two novel control methods (fishmeal and diesel fuel) on the soil surface resulted in suppression of termite damage and subsequent reduction in lodging of plants. In the majority of trials total yields (lodged and upright plants) did not differ over insecticide treatments. However, the proportion of the total yield that had to be hand-harvested from lodged plants ranged from 0 to

41%, and was significantly higher in ineffective treatments. This resulted in increased production costs and uneconomic maize production

REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 3 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:177184 CAPLUS
 DOCUMENT NUMBER: 138:333176
 TITLE: Effect of **imidacloprid** tree treatments on the occurrence of formosan subterranean **termites**, *Coptotermes formosanus* Shiraki (Isoptera: Rhinotermitidae), in independent monitors
 AUTHOR(S): Osbrink, Weste L. A.; Lax, Alan R.
 CORPORATE SOURCE: Southern Regional Research Center, USDA-ARS, New Orleans, LA, 70124, USA
 SOURCE: Journal of Economic Entomology (2003), 96(1), 117-125
 CODEN: JEENAI; ISSN: 0022-0493
 PUBLISHER: Entomological Society of America
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Periodic sampling of 87 independent monitors, initially active with the Formosan subterranean termite, *Coptotermes formosanus* Shiraki, was conducted. Monitors, located in eight sectors adjacent to seven buildings, were various distances (1-46 m) from 57 trees treated with 0.1% **imidacloprid** foam. **Termites** collected from six of the eight sectors showed latent mortality attributed to **imidacloprid** intoxication at all monitor-tree distances. Approx. 6 mo after treatment, termite populations had recovered in these sectors. Another sector showed termite population suppression for ≈15 mo, followed by recovery. **Imidacloprid** tree treatments did not control *C. formosanus* populations in independent monitors adjacent to the treatments.

REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 4 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:54728 CAPLUS
 DOCUMENT NUMBER: 138:333162
 TITLE: Comparative evaluation of chemical and botanical insecticides against **termites**
 AUTHOR(S): Singh, S. K.; Singh, G.
 CORPORATE SOURCE: Indian Institute of Pulses Research, Kanpur, 208024, India
 SOURCE: Entomon (2002), 27(2), 153-160
 CODEN: ENTOD5; ISSN: 0377-9335
 PUBLISHER: Association for Advancement of Entomology
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Insecticides viz., **Imidacloprid** 17.8 SL, chlorpyrifos 20 EC, lindane 20 EC, endosulfan 35 EC, cypermethrin 10 EC and phorate 10G and neem manure were tested against **termites** in pots. **Imidacloprid** 0.012% was effective up to 3 mo but at 0.008 and 0.004% were effective up to 2 mo only. Chlorpyrifos at 0.04% was effective up to 2 mo but at 0.02 and 0.03% were effective up to one month only. Lindane at 0.03 and 0.04% and endosulfan at 0.08% were effective up to one month. All the above insecticides gave above 50% corrected mortality. Lindane 0.02%, endosulfan 0.07%, neem manure 50 g per pot, phorate 0.1 g a.i. per pot and cypermethrin 0.0025% were found least effective. Among botanical insecticides, Nimbecidine and Nemactin were effective up to two months while Rakshak, Multineem, Neemgourd and Vanguard were effective for

09886197

short time up to one month. Field trial was conducted in mango orchards of Upeda, Ghaziabad and Rohenda, Bulandshahar, Uttar Pradesh, India. **Imidacloprid** 0.012%, chlorpyrifos 0.04% and lindane 0.04% were found most effective and gave 100% reduction in termite population up to five months. **Imidacloprid** 0.004%, chlorpyrifos 0.02%, lindane 0.02%, lindane 1.3% dust @ 100 g per tree and neem manure 500 g per tree were found less effective.

REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 5 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:14144 CAPLUS

DOCUMENT NUMBER: 138:40461

TITLE: Manufacture of additive-containing prefoamed polymer particles

INVENTOR(S): Maeda, Tadanobu

PATENT ASSIGNEE(S): Mitsubishi Chemical Foam Plastic Corp., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 7 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2003001627	A2	20030108	JP 2001-193245	20010626
PRIORITY APPLN. INFO.:			JP 2001-193245	20010626

AB The particles are manufactured by heating expandable polymer particles in a prefoaming apparatus under stirring and adding plastic additives to the prefoaming polymer particles. Thus, 600 g Styropor JF 200 (polystyrene expandable particle) was prefoamed, mixed with 0.1 part **imidacloprid** at expansion ratio 2, and further expanded to give prefoamed particles (expansion ratio 50), which were molded to give a plastic foam molding with compressive strength at 5% strain (JIS A 9511) 118 kPa, bending strength 273 kPa, d. 20.3 g/L, and reduced damage caused by **termites**.

L3 ANSWER 6 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:695680 CAPLUS

DOCUMENT NUMBER: 137:228094

TITLE: Termiticidal baits for eliminating termite colonies
INVENTOR(S): Brode, Philip Frederick, III; Garrett, Garry Steven; Laughlin, Leo Timothy; Matthews, Randall Stryker; Barker, Dale Edwin; Kinne, Daniel James; Miller, Christopher Miles; Probst, Timothy Robert; McKibben, Gary Eugene

PATENT ASSIGNEE(S): The Procter & Gamble Company, USA

SOURCE: PCT Int. Appl., 61 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002069704	A2	20020912	WO 2002-US6200	20020301
WO 2002069704	A3	20021114		

7/21/04

09886197

WO 2002069704 C1 20031231

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,
UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH,
CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR,
BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

US 2002172658 A1 20021121 US 2001-799184 20010305

US 6716421 B2 20040406

US 2003017187 A1 20030123 US 2002-172855 20020617

US 2003124166 A1 20030703 US 2002-173527 20020617

US 2003124164 A1 20030703 US 2002-268356 20021010

WO 2003105580 A1 20031224 WO 2003-US17713 20030605

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, OM, PH, PL, PT, RO,
RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ,
VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG

WO 2003106395 A1 20031224 WO 2003-US17714 20030605

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, OM, PH, PL, PT, RO,
RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ,
VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG

WO 2004032625 A2 20040422 WO 2003-US32092 20031007

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,
PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ,
UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU,
TJ, TM
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.:

US 2001-799184 A 20010305

US 2002-172855 A 20020617


US 2002-173527 A 20020617

US 2002-268356 A 20021010

OTHER SOURCE(S): MARPAT 137:228094

GI

7/21/04



AB This invention relates to devices, kits, and methods for eliminating termite colonies. The kits, devices, and methods employ a termiticidal bait matrix contain (a) a termiticide (I, X = nil, a hydrocarbon group, O or NR₈,R₉ where R₈ and R₉ are H or a hydrocarbon group; X₁ = CH, a carbon atom or a heteroatom; R,R₁,R₂,R₃ = H or OH and if R₄ and R₅ are O and R₆ and R₇ are H then R,R₁,R₂ and R₃ may be C₁-6; R₄ and R₅ are H, O or N; R₉ and R₁₀ are nil, C₁-6, and amides) selected such that the termiticide causes death to about 50 to about 100% of **termites** within about 24 to about 84 days after the **termites** begin to ingest the termiticide or the bait matrix comprising the termiticide, (b) a cellulose containing material, and (c) water. The termiticidal bait matrix can be used in a bait station installed in the ground. The kits are suitable to be used by consumers in their homes.

L3 ANSWER 7 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:547214 CAPLUS

DOCUMENT NUMBER: 137:105178

TITLE: Termiticides containing 2-pyridinethiol-1-oxide salts and wood and polymers containing the termiticides

INVENTOR(S): Nishimoto, Koichi; Sato, Toshio; Suga, Mamoru

PATENT ASSIGNEE(S): Yoshitomi Fine Chemicals Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002205906	A2	20020723	JP 2001-337124	20010926
PRIORITY APPLN. INFO.: JP 2000-381082 A 20001108				

AB The termiticides, which are effective on **termites**, bark beetles, etc., and environmentally safe, contain (a) ≥1 selected from Cu, Zn, and Na salts of 2-pyridine-1-oxide and optionally (b) ≥1 selected from pyrethroids, nicotinoids, organophosphorus compds., isocyanuric acid compds., carbamates, acetamiprid, and inorg. boric acid compds. Wood and polymers containing the termiticides are also claimed. A wood block was coated with DMSO solution containing Cu pyrithione and **imidacloprid** and dried at room temperature for ≥20 days. The wood block.

L3 ANSWER 8 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:317186 CAPLUS

DOCUMENT NUMBER: 136:365273

TITLE: Effect of insecticide treatments against **termites** on yield and quality of sugarcane

AUTHOR(S): Singh, Manager; Singh, N. B.

CORPORATE SOURCE: Sugarcane Research Institute, Shahjahanpur, 242 001,

India
 SOURCE: Sugar Cane International (2002), (March/April), 27-29
 CODEN: SCINFQ; ISSN: 1468-6031
 PUBLISHER: Agra Europe (London) Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB In a field experiment in 1995-97 at three sites in Uttar Pradesh sugarcane cv. Cos 767 setts were treated with several insecticides for the control of **termites**. Mean cane yields were highest with treatment with 0.20% solution **imidacloprid** 70 ws (77.8 t/ha), 2.5 kg ai/ha phorate 10 G (76.1 t), 2.5 kg ai/ha chlorpyrifos 15 G (73.9 t) and 1 kg ai/ha chlorpyrifos 20 EC (73.5 t) compared with the control yield of 54.4 t. Cane juice sucrose content was highest with 0.20% solution **imidacloprid** 70 WS (17.53%) compared with the control of 14.96%.
 REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 9 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:720924 CAPLUS
 DOCUMENT NUMBER: 135:340463
 TITLE: Chemical prevention of colony foundation by *Cryptotermes brevis* (Isoptera: Kalotermitidae) in attic modules
 AUTHOR(S): Scheffrahn, Rudolf H.; Busey, Philip; Edwards, Jeffrey K.; Krecek, Jan; Maharajh, Boudanath; Su, Nan-Yao
 CORPORATE SOURCE: Ft. Lauderdale Research and Education Center, University of Florida, Fort Lauderdale, FL, 33314, USA
 SOURCE: Journal of Economic Entomology (2001), 94(4), 915-919
 CODEN: JEENAI; ISSN: 0022-0493
 PUBLISHER: Entomological Society of America
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Disodium octaborate tetrahydrate (DOT) dust, DOT aqueous solution, **imidacloprid** dust, and amorphous silica gel dust with synergized 1% pyrethrins were applied on wood surfaces to simulated attic modules. Modules (30 by 30 cm) with and without fiberglass insulation were exposed to dispersal flights of *Cryptotermes brevis* (Walker) in May and June of 1998 and 1999. Six months after flights, modules were disassembled and inspected for nuptial chamber location and contents. During both years, air and water control treatments contained 22.2 ± 9.94 (mean \pm SD) nuptial chambers, 7.5 ± 5.7 live imagos, and 2.0 ± 1.4 chambers with brood. This survivorship indicated that the attic modules performed well as a colonizing platform for *C. brevis*. *C. brevis* dealates preferred constructing nuptial chambers in the crevices at the bases or tops of the modules instead of internal crevices. Modules treated in 1998 and 1999 with DOT or silica dusts contained no live **termites**, whereas zero of five modules treated with **imidacloprid** dust in 1998 and two of 20 modules treated with **imidacloprid** dust in 1999 contained single live incipient colonies. In 1998, 15% DOT solution, applied as a postconstruction treatment, yielded significantly fewer chambers and live **termites** than controls, but was not as effective as dusts in preventing successful colonization. In 1999, the DOT solution, applied as a construction-phase treatment, was equally as effective in preventing colonization as the dust treatments during that year. Results indicate that dust formulations of DOT, silica gel, and **imidacloprid** can be used to prevent drywood termite colonization in existing building voids and attics. Where the entire wood framing is exposed to treatment, such as during building construction, aqueous DOT solution can be equally effective
 as

09886197

dusts in preventing colonization by *C. brevis*.

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 10 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:336305 CAPLUS

DOCUMENT NUMBER: 135:1645

TITLE: Effects of sublethal exposure to **imidacloprid**
on subsequent behavior of subterranean termite
Reticulitermes virginicus (Isoptera: Rhinotermitidae)

AUTHOR(S): Thorne, Barbara L.; Breisch, Nancy L.

CORPORATE SOURCE: Department of Entomology, University of Maryland,
College Park, MD, 20742, USA

SOURCE: Journal of Economic Entomology (2001), 94(2), 492-498
CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Expts. were conducted to determine whether subterranean **termites**,
Reticulitermes virginicus (Banks), previously exposed to sublethal doses
of **imidacloprid** (Premise), and allowed to recover for 1 wk,
demonstrated behavioral aversion to a subsequent exposure. Worker
termites experiencing a previous sublethal but debilitating
exposure to **imidacloprid**-treated sand (either 10 or 100 ppm for
4 h) showed no apparent aversion to a second encounter with
imidacloprid-treated sand under conditions of this experiment. If these
laboratory results hold in the field and **termites** traveling through a
zone of soil treated with **imidacloprid** are impaired but
subsequently recover, they will be just as likely as their naive nestmates
to reenter the treated area if their travels take them through the
nonrepellent application a second time. Thus, a sublethal exposure to
imidacloprid can affect termite tunneling behavior. Many worker
termites that received an initial 4-h exposure to 100 ppm
imidacloprid-treated sand died, but those that survived tunneled
significantly less than did their naive nestmates, as did some
termites exposed to 10 ppm **imidacloprid**.

REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 11 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:283714 CAPLUS

DOCUMENT NUMBER: 134:276894

TITLE: Nonedible foraging matrix insert for subterranean
termite control

INVENTOR(S): Koehler, Philip G.; Oi, Faith M.

PATENT ASSIGNEE(S): University of Florida, USA; United States of America,
as Represented by the Secretary of Agriculture

SOURCE: PCT Int. Appl., 30 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001026456	A1	20010419	WO 2000-US6591	20000314
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS,				

JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK,
 MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ,
 TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD,
 RU, TJ, TM

RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE,
 DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF,
 CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

AU 758489 B2 20030320 AU 2000-37432 20000314

AU 2000037432 A5 20010423

JP 2004500043 T2 20040108 JP 2001-529256 20000314

PRIORITY APPLN. INFO.:

US 1999-159266P P 19991013

WO 2000-US6591 W 20000314

AB A several step process starts with taking a tube with a removal cap at one end, such as a two to four inch PVC tube, and filling the inner chamber with a food source such as rolled cardboard. The tube is then placed with its open end adjacent to a termite population, so that live **termites** can then enter the entrance/exit of the tube to reach the food source. Once **termites** are inside the tube, the cap is removed from the tube, and a nonedible foraging matrix, such as a disk of loose soil and or sand that is treated with a slow acting and nonrepellent toxicant, is placed between the food source in the chamber and the termite entrance/exit of the chamber. Slow-acting and non-repellent toxicants can be fipronil, chlorfenapyr, **imidacloprid**, and chlorpyrifos. The **termites** are then forced to pass through and disperse the slow-acting and non-repellent toxicant on soil particles or other nonedible foraging matrixes through their tunnels and living space in order to kill **termites**. **Termites** that contact tunnels and living space contaminated with the treated nonedible foraging matrix particles die over time.

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 12 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:573349 CAPLUS

DOCUMENT NUMBER: 133:248356

TITLE: Feeding inhibition and mortality in *Reticulitermes flavipes* (Isoptera: Rhinotermitidae) after exposure to **imidacloprid**-treated soils

AUTHOR(S): Ramakrishnan, Rathna; Suiter, Daniel R.; Nakatsu, Cindy H.; Bennett, Gary W.

CORPORATE SOURCE: Center for Urban & Industrial Pest Management, Department of Entomology, Purdue University, West Lafayette, IN, 47907, USA

SOURCE: Journal of Economic Entomology (2000), 93(2), 422-428
 CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Feeding inhibition and mortality of *Reticulitermes flavipes* (Kollar) exposed to sand, sandy loam, loam, and silty clay loam soils treated with several concns. of **imidacloprid** were studied using bioassay techniques under laboratory conditions. Termite workers stopped feeding after exposure to treated soils. Differences in feeding reduction varied among the soil types. Based on the magnitude of the F-statistics, the effect of **imidacloprid** on the reduction of termite feeding was greatest in sand followed by sandy loam, loam, and silty clay loam soils. Soil properties such as organic matter content, silt and clay proportions, pH, and cation exchange capacity were suggested to affect the bioavailability of **imidacloprid**. Similar soil effects on mortality were observed in

termites continuously exposed to treated soil for 21 days. In 3 of 4 soils tested, susceptibility to **imidacloprid** was not affected by the source of the **termites** tested.

REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 13 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:467897 CAPLUS

DOCUMENT NUMBER: 133:85609

TITLE: Termiticidal baits comprising nonhygroscopic agents in hygroscopic containers

INVENTOR(S): Minakawa, Fumiyasu; Uchida, Yuki

PATENT ASSIGNEE(S): Yuko Chemical Industries Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2000189031	A2	20000711	JP 1998-369335	19981225
PRIORITY APPLN. INFO.:			JP 1998-369335	19981225

AB A nonhygroscopic agent for controlling **termites** (e.g. diflubenzuron) is housed in a container which is made of an edible hygroscopic material (cellulosic cloth, polyvinyl alc. film). Thus, sulfluramid 0.001 and pine oil 1% (attractant) were dissolved in polyethylene glycol, and the solution was made to soaked into wood flour at a 25/100 weight ratio. The agent was heat sealed in an envelope (15 + 7 cm) made of nonwoven fabric of cellulose fibers with 1% by weight added pine oil to obtain a bait with satisfactory attractiveness to *Reticulitermes*.

L3 ANSWER 14 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:52160 CAPLUS

DOCUMENT NUMBER: 132:133596

TITLE: Degradation of bifenthrin, chlorpyrifos and **imidacloprid** in soil and bedding materials at termiticidal application rates

AUTHOR(S): Baskaran, Sundaram; Kookana, Rai S.; Naidu, Ravendra

CORPORATE SOURCE: CSIRO Land and Water, Glen Osmond, 5064, Australia

SOURCE: Pesticide Science (1999), 55(12), 1222-1228

CODEN: PSSCBG; ISSN: 0031-613X

PUBLISHER: John Wiley & Sons Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Organophosphorus, pyrethroid and chloronicotinyl insecticides have been used to control **termites** in building structures in recent years. The degradation behavior of three insecticides (bifenthrin, chlorpyrifos and **imidacloprid**) at termiticidal application rates was studied under standard laboratory conditions (25°C, 60% field moisture capacity and darkness) for 24 mo. The study was carried out on one soil and two bedding materials (sand-dolomite and quarry sand), which are commonly used under housing in Australia. Expts. were also conducted to examine the effect of soil moisture on the degradation of these insecticides. Insecticide residues in the samples collected at different days after application were measured by HPLC. The rate of degradation of bifenthrin and **imidacloprid** insecticides was adequately described by a first-order kinetic model ($r^2=0.93-0.97$). However, chlorpyrifos degradation

09886197

was biphasic, showing an initial faster degradation followed by a slower rate. Therefore, the degradation data during the slower phase only (after a two-month period) followed the first-order law ($r^2=0.95$). Soil moisture had little effect on degradation of **imidacloprid** and bifenthrin. Among the three insecticides, bifenthrin and **imidacloprid** were most stable and chlorpyrifos the least. Chlorpyrifos showed a major loss (75-90%) of residue during the 24 mo incubation period. In the bedding materials, simultaneous accumulation of the primary metabolite of chlorpyrifos, TCP (3,5,6-trichloro-2-pyridinol) was observed. Hydrolysis appeared to have caused the observed rapid loss of chlorpyrifos, especially in the

highly alkaline bedding materials (sand-dolomite and quarry sand).

REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 15 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:799698 CAPLUS

DOCUMENT NUMBER: 132:9953

TITLE: Termite control

INVENTOR(S): De Villiers, Vivian; Van der Westhuizen, M. C.; Robbertse, Ernest

PATENT ASSIGNEE(S): Bayer A.-G., Germany

SOURCE: S. African, 16 pp.

CODEN: SFXXAB

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
ZA 9711701	A	19980706	ZA 1997-11701	19971230
AP 1174	A	20030630	AP 1998-1424	19981228
W: BW, GH, GM, KE, LS, MW, SD, SZ, UG, ZM, ZW				
BR 9805735	A	20010424	BR 1998-5735	19981229

PRIORITY APPLN. INFO.: ZA 1997-11701 A 19971230

AB Agonists or antagonists of nicotinergerg acetylcholine receptors of insects are used for the control of harvester **termites**, i.e. *Hodotermitidae*. **Imidacloprid** is the preferred active ingredient. The bait formulations comprise lucerne or grass particles.

L3 ANSWER 16 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:797191 CAPLUS

DOCUMENT NUMBER: 132:60446

TITLE: **Imidacloprid**-enhanced *Reticulitermes flavipes* (Isoptera: Rhinotermitidae) susceptibility to the entomopathogen *Metarhizium anisopliae*

AUTHOR(S): Ramakrishnan, Rathna; Suiter, Daniel R.; Nakatsu, Cindy H.; Humber, Richard A.; Bennett, Gary W.

CORPORATE SOURCE: Center for Urban & Industrial Pest Management, Department of Entomology, Purdue University, West Lafayette, IN, 47907, USA

SOURCE: Journal of Economic Entomology (1999), 92(5), 1125-1132

CODEN: JEENAI; ISSN: 0022-0493

PUBLISHER: Entomological Society of America

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The effects of **imidacloprid** and the entomopathogen *Metarhizium*

anisopliae (Metsch.) Sorokin on the eastern subterranean termite, *Reticulitermes flavipes* (Kollar), were evaluated in a 4 + 3 factorial experiment in both sterile and nonsterile loam soil. **Termites** were not susceptible to *M. anisopliae* when assays were conducted in nonsterile soil, but were highly susceptible in sterile soil. Termite mortality after 21 days of continuous exposure to 104 conidia per g soil was 0 and 41.6% in nonsterile and sterile soil, resp. **Termites** were significantly more susceptible to sterile soil containing 107 conidia per g than to the same soil containing 104 conidia per

g.

In continuous exposure assays, **termites** were highly susceptible to **imidacloprid**-treated (5,10, and 20 ppm) nonsterile and sterile soil containing no exptl. introduced *M. anisopliae*. Exposure of **termites** to **imidacloprid** enhanced their susceptibility to introduced *M. anisopliae* in nonsterile and sterile soil. Native entomopathogens recovered from **termites** exposed to **imidacloprid**-treated, nonsterile soil (i.e., no introduced *M. anisopliae*) included *Conidiobolus coronatus* (Constantin) Batko, *Cunninghamella echinulata* Thaxter, *Fusarium* spp., *Aspergillus* spp., and a naturally occurring strain of *M. anisopliae* variety majus.

REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 17 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:125767 CAPLUS

DOCUMENT NUMBER: 130:178773

TITLE: Composition for the control of wood-destroying insects, especially **termites**

INVENTOR(S): Anderson, John-phillip-evans; Keuken, Oliver

PATENT ASSIGNEE(S): Bayer A.-G., Germany

SOURCE: Eur. Pat. Appl., 21 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 896791	A2	19990217	EP 1998-114187	19980729
EP 896791	A3	20000112		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
DE 19734665	A1	19990218	DE 1997-19734665	19970811
TW 505500	B	20021011	TW 1998-87112592	19980731
US 6264968	B1	20010724	US 1998-128818	19980804
ZA 9807118	A	19990209	ZA 1998-7118	19980807
JP 11124302	A2	19990511	JP 1998-234861	19980807
AU 9879895	A1	19990218	AU 1998-79895	19980811
AU 768390	B2	20031211		
BR 9803138	A	19991221	BR 1998-3138	19980811

PRIORITY APPLN. INFO.: DE 1997-19734665 A 19970811

AB The title comps. (no examples) comprise an insecticide, preferably **imidacloprid**, incorporated into an organic natural and/or synthetic carrier. Optional ingredients are insect attractants and microbicides.

L3 ANSWER 18 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:54407 CAPLUS

DOCUMENT NUMBER: 130:206253

09886197

TITLE: Control of the termite *Heterotermes tenuis* (Hagen) using Termitrap baits impregnated with insecticides associated with the entomopathogenic fungus *Beauveria bassiana* (Bals.) Vuill.

AUTHOR(S): Almeida, Jose E. M.; Alves, Sergio B.; Moino, Alcides, Jr.; Lopes, E. Rogerio B.

CORPORATE SOURCE: Laboratorio de Controle Biologico, Centro Experimental, Instituto Biologico, Campinas, 13001-970, Brazil

SOURCE: Anais da Sociedade Entomologica do Brasil (1998), 27(4), 639-644

CODEN: ASENBI; ISSN: 0301-8059

PUBLISHER: Sociedade Entomologica do Brasil

DOCUMENT TYPE: Journal

LANGUAGE: Portuguese

AB The control of *H. tenuis* was evaluated using the bait/trap Termitrap impregnated with insecticides in low concns., associated to *B. bassiana* isolate 634 (from *Solenopsis invicta*), in sugarcane (*Saccharum officinarum*). The treatments consisted of: **imidacloprid** 0,01%; **imidacloprid** 0,01% + *B. bassiana*; WG 0,003%; WG 0,003% + *B. bassiana*; *B. bassiana*; and untreated control. Each treatment was replicated five times. The insecticides were impregnated on baits by immersion in water, their concns. being calculated according to the weight of the bait, and the *B. bassiana* was applied as pure conidia (109 conidia/bait). The evaluations were made after 15, 30, 41, 63, 86 e 136 days, by assigning indexes to populations levels. All treatments significantly reduced termite populations when compared to the control. It took longer for *B. bassiana* alone to reduced *H. tenuis* population. The treatments with **imidacloprid** and WG were the most efficient in the control of **termites** in sugarcane. The baits/traps did not repel the **termites**.

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 19 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1996:411657 CAPLUS

TITLE: **Imidacloprid** - chemical synergist for microbial control agents of **termites**.

AUTHOR(S): Boucias, D. G.

CORPORATE SOURCE: Department Entomology & Nematology, University Florida, Gainesville, FL, 32611-0620, USA

SOURCE: Book of Abstracts, 212th ACS National Meeting, Orlando, FL, August 25-29 (1996), AGRO-019. American Chemical Society: Washington, D. C.

CODEN: 63BFAF

DOCUMENT TYPE: Conference; Meeting Abstract

LANGUAGE: English

AB Our research has determined that the neurotoxin, **imidacloprid**, at sublethal concns., can significantly alter the behavioral patterns of insects. For example, the subterranean termite, *Reticulotermis flavipes* possesses social behaviors (grooming, tunnel construction) which serve as the primary line of defense against pathogenic and opportunistic microorganisms. These behaviors, in combination with the resident microflora, confer a high degree of disease resistance upon these social insects. Exposure to low dosages of **imidacloprid** produces a long term disruption of these social behaviors resulting in the onset of epizootics initiated by either resident or introduced microbes. Related studies on other nonsocial insects (cockroaches, weevils) have supported


09886197

the results found with **termites**. At sublethal concns., **imidacloprid** acted as a behavioral modifying agent significantly increasing the host insects susceptibility to microbial control agents.

L3 ANSWER 20 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 1995:648220 CAPLUS
DOCUMENT NUMBER: 123:27832
TITLE: Odorless insect repellents against **termites**
INVENTOR(S): Ueda, Masayoshi; Muto, Yutaka
PATENT ASSIGNEE(S): Japan Carlit Co Ltd, Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 07089803	A2	19950404	JP 1993-258961	19930924
PRIORITY APPLN. INFO.:			JP 1993-258961	19930924
OTHER SOURCE(S):	MARPAT 123:27832			

GI



AB An odorless insect repellent contains a repellent, a solvent and surfactant, or preservative; the solvent being I (R1, R2 = H, C1-2 alkyl; R3 = C1-3 alkyl). The active repellent may be chlorpyrifos, phoxim, pyridaphenthion, allethrin, carbaril, **imidacloprid**, etc. For example, an odorless emulsion was prepared by combining dimethylpropylnaphthalene, chlorpyrifos, Sorpol-3006K and Sorpol-3008K.

L3 ANSWER 21 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN
ACCESSION NUMBER: 1995:187187 CAPLUS
DOCUMENT NUMBER: 122:25815
TITLE: **Imidacloprid** - a new systemic insecticide.
AUTHOR(S): Elbert, A.; Becker, B.; Hartwig, J.; Erdelen, C.
CORPORATE SOURCE: Geschäftsbereich Pflanzenschutz
Entwicklung/Insektizide, Bayer AG, Leverkusen, 5090, Germany
SOURCE: Pflanzenschutz-Nachrichten Bayer (German Edition)
(1991), 44(2), 113-36
CODEN: PNBAT; ISSN: 0340-1723
PUBLISHER: Bayer AG
DOCUMENT TYPE: Journal
LANGUAGE: German

AB The biol. profile of **Imidacloprid** (I) was defined on the basis of the results of exhaustive laboratory expts. and greenhouse trials. I is extremely effective against sucking insects, such as rice leafhoppers, aphids, thrips and mealybugs, and very effective against whitefly. It is also effective against some species of biting insects, such as paddy stem

borers and Colorado beetle, but it has no effect on nematodes or spider mites. At comparatively high doses it kills adult insects and has ovicidal effects. I is a nicotinic acetylcholine receptor stimulator. Its mechanism of action differs from that of conventional insecticides. It therefore gives excellent control of all resistant populations investigated hitherto. I has a pos. temperature coefficient After foliar application, it has a good residual action, it is highly photostable and it shows satisfactory resistance to rain. I is active after oral ingestion and by direct contact, but it is not active in the vapor phase. The LD95 after oral ingestion by *Myzus persicae* is .apprx.2 pg/aphid. After topical application it is .apprx.160 pg/aphid. It has not been possible to demonstrate recovery of injured aphids, or antifeeding effects. I has a faster action against aphids than oxydemeton-Me. After foliar application, I shows good translaminar and acropetal translocation, so it is also likely to provide effective control of pests with a furtive lifestyle, and protect the parts of the plant which regenerate after treatment. By virtue of its good contact action and powerful systemic action after uptake through the root system, I can be applied to soil and used as a seed dressing. It gives excellent control of pests such as onion maggots, *Diabrotica*, wire worms, **termites** and fire ants which live in the soil, and of insects such as aphids which live above ground level. It has a good residual action after application to the soil and when it is used as a seed dressing. The compatibility of I with plants is good after use as a seed dressing, as a soil treatment and after foliar application. By virtue of its biol. properties, I is likely to have a wide range of uses for controlling economically important pests of rice, cotton, cereals, maize, sugar beet, potatoes, vegetables, citrus fruit, pome and stone fruit and other crops.

L3 ANSWER 22 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1993:54353 CAPLUS
 DOCUMENT NUMBER: 118:54353
 TITLE: Imidozolidine derivatives and related compounds as industrial insecticides and wood preservatives
 INVENTOR(S): Tsuboi, Shinichi; Sone, Shinzaburo; Obinata, Toru; Exner, Otto; Schwamborn, Michael
 PATENT ASSIGNEE(S): Nihon Bayer Agrochem K. K., Japan
 SOURCE: Eur. Pat. Appl., 15 pp.
 CODEN: EPXXDW
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 511541	A1	19921104	EP 1992-106384	19920414
EP 511541	B1	19960904		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
JP 05032505	A2	19930209	JP 1991-350751	19911212
JP 3162450	B2	20010425		
JP 2001031511	A2	20010206	JP 2000-233512	19911212
AU 9213908	A1	19921029	AU 1992-13908	19920330
AU 645672	B2	19940120		
AT 142077	E	19960915	AT 1992-106384	19920414
ES 2090400	T3	19961016	ES 1992-106384	19920414
BR 9201534	A	19921201	BR 1992-1534	19920427
US 6323224	B1	20011127	US 1995-543351	19951016
US 2001051643	A1	20011213	US 2001-886197	20010621


09886197

PRIORITY APPLN. INFO.:

JP 1991-125172	A 19910427
JP 1991-350751	A 19911212
US 1992-872279	B1 19920422
US 1995-543351	A3 19951016

OTHER SOURCE(S):
GI

MARPAT 118:54353



AB The imidazolidine derivs. and related compds. I (X = NH, S; Y = CH, N; Z = 2-chloro-5-pyridyl, 2-chloro-5-thiazolyl; R = H, Me; n = 0, 1) are industrial insecticides and wood preservatives. Wood impregnated with 0.32 ppm **imidacloprid** was lethal to **termites** (*Coptotermes formosanus*) for ≥ 3 wk.